

**ALUNBRIG™ APPROVED PACKAGE INSERT**  
**PROFESSIONAL INFORMATION**

**SCHEDULING STATUS**

S4

**1. NAME OF THE MEDICINE**

**ALUNBRIG™ 30 mg film-coated tablets**

**ALUNBRIG™ 90 mg film-coated tablets**

**ALUNBRIG™ 180 mg film-coated tablets**

**2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

**ALUNBRIG™ 30 mg film-coated tablets**

Each film-coated tablet contains 30 mg of brigatinib.

*Excipient with known effect*

Each film-coated tablet contains 56 mg of lactose monohydrate.

**ALUNBRIG™ 90 mg film-coated tablets**

Each film-coated tablet contains 90 mg of brigatinib.

*Excipient with known effect*

Each film-coated tablet contains 168 mg of lactose monohydrate.

**ALUNBRIG™ 180 mg film-coated tablets**

Each film-coated tablet contains 180 mg of brigatinib.

*Excipient with known effect*

Each film-coated tablet contains 336 mg of lactose monohydrate.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

#### **ALUNBRIG™** 30 mg film-coated tablets

Round, white to off-white film-coated tablet of approximately 7 mm in diameter with debossed “U3” on one side and plain on the other side.

#### **ALUNBRIG™** 90 mg film-coated tablets

Oval, white to off-white film-coated tablet of approximately 15 mm in length with debossed “U7” on one side and plain on the other side.

#### **ALUNBRIG™** 180 mg film-coated tablets

Oval, white to off-white film-coated tablet of approximately 19 mm in length with debossed “U13” on one side and plain on the other side.

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

**ALUNBRIG™** is indicated as monotherapy for the treatment of adult patients with anaplastic lymphoma kinase (ALK) positive advanced non small cell lung cancer (NSCLC) of the adenocarcinoma cell type previously not treated with an ALK inhibitor.

**ALUNBRIG™** is indicated as monotherapy for the treatment of adult patients with anaplastic lymphoma kinase (ALK)-positive advanced non-small cell lung cancer (NSCLC) of the adenocarcinoma cell type, who have progressed on or are intolerant to crizotinib.

#### 4.2 Posology and method of administration

Treatment with **ALUNBRIG™** should be initiated and supervised by a medical practitioner experienced in the use of anticancer medicines.

ALK-positive NSCLC status should be known prior to initiation of **ALUNBRIG™** therapy. A validated ALK assay is necessary for the selection of ALK-positive NSCLC patients (see section 5.1).

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

Assessment for ALK-positive NSCLC should be performed by laboratories with demonstrated proficiency in the specific technology being utilised.

### Posology

The recommended starting dose of **ALUNBRIG™** is 90 mg once daily for the first 7 days, then 180 mg once daily.

If **ALUNBRIG™** is interrupted for 14 days or longer for reasons other than adverse reactions, treatment should be resumed at 90 mg once daily for 7 days before increasing to the previously tolerated dose.

If a dose is missed or vomiting occurs after taking a dose, an additional dose should not be administered and the next dose should be taken at the scheduled time.

Treatment should continue as long as clinical benefit is observed.

### Dose adjustments

Dosing interruption and/or dose reduction may be required based on individual safety and tolerability.

**ALUNBRIG™** dose modification levels are summarised in Table 1.

**Table 1: Recommended ALUNBRIG™ dose reduction levels**

Dose	Dose reduction levels		
	First	Second	Third
90 mg once daily (first 7 days)	reduce to 60 mg once daily	permanently discontinue	not applicable
180 mg once daily	reduce to 120 mg once daily	reduce to 90 mg once daily	reduce to 60 mg once daily

**ALUNBRIG™** should be permanently discontinued if patient is unable to tolerate the 60 mg once daily dose.

Recommendations for dose modifications of ALUNBRIG™ for the management of adverse reactions are summarised in Table 2.

**Table 2: Recommended ALUNBRIG™ dose modifications for adverse reactions**

Adverse reaction	Severity*	Dose modification
Interstitial lung disease (ILD)/pneumonitis	Grade 1	<ul style="list-style-type: none"> <li>If event occurs during the first 7 days of treatment, <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at same dose level and not escalated to 180 mg once daily.</li> <li>If ILD/pneumonitis occurs after the first 7 days of treatment, <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at same dose level.</li> <li>If Interstitial lung disease ILD/pneumonitis recurs, <b>ALUNBRIG™</b> should be permanently discontinued.</li> </ul>
	Grade 2	<ul style="list-style-type: none"> <li>If ILD/pneumonitis occurs during the first 7 days of treatment, <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at next lower dose level as described in Table 1 and not escalated to 180 mg once daily.</li> <li>If ILD/pneumonitis occurs after the first 7 days of treatment, <b>ALUNBRIG™</b> should be withheld until recovery to baseline. <b>ALUNBRIG™</b> should be resumed at next lower dose level as described in Table 1.</li> <li>If ILD/pneumonitis recurs, <b>ALUNBRIG™</b> should be permanently discontinued.</li> </ul>
	Grade 3 or 4	<ul style="list-style-type: none"> <li><b>ALUNBRIG™</b> should be permanently discontinued.</li> </ul>
Hypertension	Grade 3 hypertension (SBP ≥ 160 mmHg or	<ul style="list-style-type: none"> <li><b>ALUNBRIG™</b> should be withheld until hypertension has recovered to Grade ≤ 1</li> </ul>

	<p>DBP ≥ 100 mmHg, medical intervention indicated, more than one anti-hypertensive medicine, or more intensive therapy than previously used indicated)</p>	<p>(SBP &lt; 140 mmHg and DBP &lt; 90 mmHg), then resumed at same dose.</p> <ul style="list-style-type: none"> <li>If Grade 3 hypertension recurs, <b>ALUNBRIG™</b> should be withheld until hypertension has recovered to Grade ≤ 1 then resumed at the next lower dose level per Table 1 or permanently discontinued</li> </ul>
	<p>Grade 4 hypertension (life threatening consequences, urgent intervention indicated)</p>	<ul style="list-style-type: none"> <li><b>ALUNBRIG™</b> should be withheld until hypertension has recovered to Grade ≤ 1 (SBP &lt; 140 mmHg and DBP &lt; 90 mmHg), then resumed at the next lower dose level per Table 1 or permanently discontinued.</li> <li>If Grade 4 hypertension recurs, <b>ALUNBRIG™</b> should be permanently discontinued.</li> </ul>
<p>Bradycardia (Heart Rate less than 60 bpm)</p>	<p>Symptomatic bradycardia</p>	<ul style="list-style-type: none"> <li><b>ALUNBRIG™</b> should be withheld until recovery to asymptomatic bradycardia or to a resting heart rate of 60 bpm or above.</li> <li>If a concomitant medicine known to cause bradycardia is identified and discontinued, or its dose is adjusted, <b>ALUNBRIG™</b> should be resumed at same dose upon recovery to asymptomatic bradycardia or to a resting heart rate of 60 bpm or above.</li> <li>If no concomitant medicine known to cause bradycardia is identified, or if contributing concomitant medications are not discontinued or dose modified, <b>ALUNBRIG™</b> should be resumed at the next lower dose level per Table 1 upon recovery to asymptomatic bradycardia or to a resting heart rate of 60 bpm or above.</li> </ul>
	<p>Bradycardia with life-threatening</p>	<ul style="list-style-type: none"> <li>If contributing concomitant medicine is identified and discontinued, or its dose is</li> </ul>

	consequences, urgent intervention indicated	<p>adjusted, <b>ALUNBRIG™</b> should be resumed at the next lower dose level per Table 1 upon recovery to asymptomatic bradycardia or to a resting heart rate of 60 bpm or above, with frequent monitoring as clinically indicated.</p> <ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be permanently discontinued if no contributing concomitant medicine is identified.</li> <li>• <b>ALUNBRIG™</b> should be permanently discontinued in case of recurrence.</li> </ul>
Elevation of CPK	Grade 3 or 4 elevation of CPK ( $> 5 \times \text{ULN}$ ) with Grade $\geq 2$ muscle pain or weakness	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to Grade <math>\leq 1</math> (<math>\leq 2,5 \times \text{ULN}</math>) <u>elevation of CPK</u> or to baseline, then resumed at the same dose.</li> <li>• If Grade 3 or 4 elevation of CPK recurs <u>with Grade <math>\geq 2</math> muscle pain or weakness</u>, <b>ALUNBRIG™</b> should be withheld until recovery to Grade <math>\leq 1</math> (<math>\leq 2,5 \times \text{ULN}</math>) <u>elevation of CPK</u> or to baseline, then resumed at the next lower dose level per Table 1.</li> </ul>
Elevation of lipase or amylase	Grade 3 elevation of lipase or amylase ( $> 2 \times \text{ULN}$ )	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to Grade <math>\leq 1</math> (<math>\leq 1,5 \times \text{ULN}</math>) or to baseline, then resumed at same dose.</li> <li>• If Grade 3 elevation of lipase or amylase recurs, <b>ALUNBRIG™</b> should be withheld until recovery to Grade <math>\leq 1</math> (<math>\leq 1,5 \times \text{ULN}</math>) or to baseline, then resumed at the next lower dose level per Table 1.</li> </ul>
	Grade 4 elevation of lipase or amylase ( $> 5 \times \text{ULN}$ )	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to Grade <math>\leq 1</math> (<math>\leq 1,5 \times \text{ULN}</math>), then resumed at the next lower dose level per Table 1.</li> </ul>

Hepatotoxicity	Grade $\geq 3$ elevation ( $> 5 \times$ ULN) of either alanine aminotransferase (ALT) or aspartate aminotransferase (AST) with bilirubin $\leq 2 \times$ ULN	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to baseline or less than or equal to <math>3 \times</math> ULN, then resumed at next lower dose per Table 1.</li> </ul>
	Grade $\geq 2$ elevation ( $> 3 \times$ ULN) of ALT or AST with concurrent total bilirubin elevation $> 2 \times$ ULN in the absence of cholestasis or haemolysis	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be permanently discontinued.</li> </ul>
Hyperglycaemia	For Grade 3 (greater than 250 mg/dL or 13,9 mmol/L) or greater	<ul style="list-style-type: none"> <li>• If adequate hyperglycaemic control cannot be achieved with optimal medical management, <b>ALUNBRIG™</b> should be withheld until adequate hyperglycaemic control is achieved. Upon recovery, <b>ALUNBRIG™</b> may either be resumed at the next lower dose per Table 1 or permanently discontinued.</li> </ul>
Visual Disturbance	Grade 2 or 3	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to Grade 1 or baseline, then resumed at the next lower dose level per Table 1.</li> </ul>
	Grade 4	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be permanently discontinued.</li> </ul>
Other adverse reactions	Grade 3	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at the same dose level.</li> <li>• If the Grade 3 event recurs, <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at the next lower dose level as per Table 1 or permanently discontinued.</li> </ul>

	Grade 4	<ul style="list-style-type: none"> <li>• <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at the next lower dose level as per Table 1.</li> <li>• If the Grade 4 event recurs, <b>ALUNBRIG™</b> should be withheld until recovery to baseline, then resumed at the next lower dose level as per Table 1 or permanently discontinued.</li> </ul>
bpm = beats per minute; CPK = Creatine Phosphokinase; DBP = diastolic blood pressure; SBP = systolic blood pressure; ULN = upper limit of normal		

\*Graded per National Cancer Institute Common Terminology Criteria for Adverse Events. Version 4.0 (NCI CTCAE v4).

### Special populations

#### *Elderly patients*

The limited data on the safety and efficacy of **ALUNBRIG™** in patients aged 65 years and older suggest that a dose adjustment is not required in elderly patients (see section 4.8). There are no available data on patients over 85 years of age.


#### *Hepatic impairment*

No dose adjustment of **ALUNBRIG™** is required for patients with mild hepatic impairment (Child-Pugh class A) or moderate hepatic impairment (Child-Pugh class B). A reduced starting dose of 60 mg once daily for the first 7 days, then 120 mg once daily is recommended for patients with severe hepatic impairment (Child-Pugh class C) (see section 5.2).

#### *Renal impairment*

No dose adjustment of **ALUNBRIG™** is required for patients with mild or moderate renal impairment (estimated glomerular filtration rate (eGFR)  $\geq$  30 mL/min).

A reduced starting dose of 60 mg once daily for the first 7 days, then 90 mg once daily is recommended for patients with severe renal impairment (eGFR < 30 mL/min) (see section 5.2). Patients with severe renal impairment should be closely monitored for new or worsening respiratory symptoms that may indicate ILD/pneumonitis (e.g., dyspnoea, cough, etc.) particularly in the first week (see section 4.4).

DocuSigned by:  
  
 9B2F238576344AB...

### *Paediatric population*

The safety and efficacy of **ALUNBRIG™** in patients less than 18 years of age have not been established. No data are available.

### Method of administration

**ALUNBRIG™** is for oral use. The tablets should be swallowed whole and with water. **ALUNBRIG™** may be taken with or without food.

Grapefruit or grapefruit juice may increase plasma concentrations of **ALUNBRIG™** and should be avoided (see section 4.5).

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Pregnancy and lactation (see section 4.6).

### **4.4 Special warnings and precautions for use**

#### Pulmonary adverse reactions

Severe, life-threatening, and fatal pulmonary adverse reactions, including those with features consistent with ILD/pneumonitis, can occur in patients treated with **ALUNBRIG™** (see section 4.8).

Most pulmonary adverse reactions were observed within the first 7 days of treatment. Grade 1-2 pulmonary adverse reactions resolved with interruption of treatment or dose modification. Increased age and shorter interval (less than 7 days) between the last dose of crizotinib and the first dose of **ALUNBRIG™** were independently associated with an increased rate of these pulmonary adverse reactions. These factors should be considered when initiating treatment with **ALUNBRIG™**. Patients with a history of ILD or drug-induced pneumonitis were excluded from the pivotal trial.

Some patients experienced pneumonitis later in treatment with **ALUNBRIG™**.

Patients should be monitored for new or worsening respiratory symptoms (e.g., dyspnoea, cough, etc.), particularly in the first week of treatment. Evidence of pneumonitis in any patient with worsening respiratory symptoms should be promptly investigated. If pneumonitis is suspected, the dose of

**ALUNBRIG™** should be withheld, and the patient evaluated for other causes of symptoms (e.g., pulmonary embolism, tumour progression, and infectious pneumonia). The dose should be modified accordingly (see section 4.2).

#### Hypertension

Hypertension has occurred in patients treated with **ALUNBRIG™** (see section 4.8). Blood pressure should be monitored regularly during treatment with **ALUNBRIG™**. Hypertension should be treated according to standard guidelines to control blood pressure. Heart rate should be monitored more frequently in patients if concomitant use of a medicine known to cause bradycardia cannot be avoided. For severe hypertension ( $\geq$  Grade 3), **ALUNBRIG™** should be withheld until hypertension has recovered to Grade 1 or to baseline. The dose should be modified accordingly (see section 4.2).

#### Bradycardia

Bradycardia has occurred in patients treated with **ALUNBRIG™** (see section 4.8). Caution should be exercised when administering **ALUNBRIG™** in combination with other medicines known to cause bradycardia. Heart rate and blood pressure should be monitored regularly.

If symptomatic bradycardia occurs, treatment with **ALUNBRIG™** should be withheld and concomitant medicines known to cause bradycardia should be evaluated. Upon recovery, the dose should be modified accordingly (see section 4.2). In case of life-threatening bradycardia, if no contributing concomitant medicines is identified or in case of recurrence, treatment with **ALUNBRIG™** should be discontinued (see section 4.2).

#### Visual disturbance

Visual disturbance adverse reactions have occurred in patients treated with **ALUNBRIG™** (see section 4.8). Patients should be advised to report any visual symptoms. For new or worsening severe visual symptoms, an ophthalmologic evaluation and dose reduction should be considered (see section 4.2).

#### Creatine phosphokinase (CPK) elevation

Elevations of CPK have occurred in patients treated with **ALUNBRIG™** (see section 4.8). Patients should be advised to report any unexplained muscle pain, tenderness, or weakness. CPK levels

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

should be monitored regularly during **ALUNBRIG™** treatment. Based on the severity of the CPK elevation, and or if associated with muscle pain or weakness, treatment with **ALUNBRIG™** should be withheld, and the dose modified accordingly (see section 4.2).

#### Elevations of pancreatic enzymes

Elevations of amylase and lipase have occurred in patients treated with **ALUNBRIG™** (see section 4.8). Lipase and amylase should be monitored regularly during treatment with **ALUNBRIG™**. Based on the severity of the laboratory abnormalities, treatment with **ALUNBRIG™** should be withheld, and the dose modified accordingly (see section 4.2).

#### Hepatotoxicity

Elevations of hepatic enzymes (aspartate aminotransferase, alanine aminotransferase) and bilirubin have occurred in patients treated with **ALUNBRIG™** (see section 4.8).

Liver function, including AST, ALT and total bilirubin should be assessed prior to the initiation of **ALUNBRIG™** and then every 2 weeks during the first 3 months of treatment. Thereafter, monitoring should be performed periodically. Based on the severity of the laboratory abnormalities, treatment should be withheld, and the dose modified accordingly (see section 4.2).

#### Hyperglycaemia

Elevations of serum glucose have occurred in patients treated with **ALUNBRIG™**. Fasting serum glucose should be assessed prior to initiation of **ALUNBRIG™** and monitored periodically thereafter. Antihyperglycaemic treatment should be initiated or optimised as needed. If adequate hyperglycaemic control cannot be achieved with optimal medical management, **ALUNBRIG™** should be withheld until adequate hyperglycaemic control is achieved; upon recovery reducing the dose as described in Table 1 may be considered or **ALUNBRIG™** may be permanently discontinued.

#### Interactions

The concomitant use of **ALUNBRIG™** with strong CYP3A inhibitors should be avoided. If concomitant use of strong CYP3A inhibitors cannot be avoided, the dose of **ALUNBRIG™** should be reduced from 180 mg to 90 mg, or from 90 mg to 60 mg. After discontinuation of a strong CYP3A inhibitor, **ALUNBRIG™** should be resumed at the dose that was tolerated prior to the initiation of the

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

strong CYP3A inhibitor.

The concomitant use of **ALUNBRIG™** with strong and moderate CYP3A inducers should be avoided (see section 4.5).

Vaccinations with live attenuated vaccines are not advised for patients taking **ALUNBRIG™**.

### Fertility

Women of childbearing potential should use effective non-hormonal contraception during treatment with **ALUNBRIG™** and for at least 4 months following the final dose. Men with female partners of childbearing potential should use effective contraception during treatment and for at least 3 months after the last dose of **ALUNBRIG™** (see section 4.6).

### Lactose

**ALUNBRIG™** contains lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, e.g., galactosemia total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

## **4.5 Interaction with other medicines and other forms of interaction**

### Medicines that may increase brigatinib plasma concentrations

#### CYP3A inhibitors

*In vitro* studies demonstrated that brigatinib is a substrate of CYP3A4/5. In healthy subjects, coadministration of multiple 200 mg twice daily doses of itraconazole, a strong CYP3A inhibitor, with a single 90 mg brigatinib dose increased brigatinib  $C_{max}$  by 21 %,  $AUC_{0-120}$  by 101 % (2-fold), and  $AUC_{0-120}$  by 82 % (< 2-fold), relative to a 90 mg brigatinib dose administered alone.

The concomitant use of strong CYP3A inhibitors with **ALUNBRIG™**, including but not limited to certain antivirals (e.g., indinavir, nelfinavir, ritonavir, saquinavir), macrolide antibiotics (e.g., clarithromycin, telithromycin, troleandomycin), antifungals (e.g., ketoconazole, voriconazole), and nefazodone should be avoided. If concomitant use of strong CYP3A inhibitors cannot be avoided, the dose of **ALUNBRIG™** should be reduced by approximately 50 % (i.e., from 180 mg to 90 mg, or from 90 mg to 60 mg). After discontinuation of a strong CYP3A inhibitor, **ALUNBRIG™** should be resumed

at the dose that was tolerated prior to the initiation of the strong CYP3A inhibitor.

Moderate CYP3A inhibitors (e.g., diltiazem and verapamil) may increase the AUC of brigatinib by approximately 40 % based on simulations from a physiologically-based pharmacokinetic model. No dose adjustment is required for **ALUNBRIG™** in combination with moderate CYP3A inhibitors. Patients should be closely monitored when **ALUNBRIG™** is coadministered with moderate CYP3A inhibitors.

Grapefruit or grapefruit juice may also increase plasma concentrations of brigatinib and should be avoided (see section 4.2).

#### CYP2C8 inhibitors

*In vitro* studies demonstrated that brigatinib is a substrate of CYP2C8. In healthy subjects, coadministration of multiple 600 mg twice daily doses of gemfibrozil, a strong CYP2C8 inhibitor, with a single 90 mg brigatinib dose reduced brigatinib  $C_{max}$  by 41 %,  $AUC_{0-INF}$  by 12 %, and  $AUC_{0-120}$  by 15 %, relative to a 90 mg brigatinib dose administered alone. The effect of gemfibrozil on the pharmacokinetics of brigatinib is not clinically meaningful and the underlying mechanism for the decreased exposure of brigatinib is unknown. No dose adjustment is required during coadministration with strong CYP2C8 inhibitors.

#### P-gp and BCRP inhibitors

Brigatinib is a substrate of P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP) *in vitro*. Given that brigatinib exhibits high solubility and high permeability, inhibition of P-gp and BCRP is not expected to result in a clinically meaningful change in the systemic exposure of brigatinib. No dose adjustment is required for **ALUNBRIG™** during coadministration with P-gp and BCRP inhibitors.

#### Medicines that may decrease brigatinib plasma concentrations

##### CYP3A inducers

In healthy subjects, coadministration of multiple 600 mg daily doses of rifampicin, a strong CYP3A inducer, with a single 180 mg brigatinib dose decreased brigatinib  $C_{max}$  by 60%,  $AUC_{0-INF}$  by 80 % (5-fold), and  $AUC_{0-120}$  by 80 % (5-fold), relative to a 180 mg brigatinib dose administered alone. The concomitant use of strong CYP3A inducers with **ALUNBRIG™**, including but not limited to rifampicin,

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd  
carbamazepine, phenytoin, rifabutin, phenobarbitone, and St. John's wort should be avoided.

Moderate CYP3A inducers may decrease the AUC of brigatinib by approximately 50 % based on simulations from a physiologically-based pharmacokinetic model. The concomitant use of moderate CYP3A inducers with **ALUNBRIG™**, including but not limited to efavirenz, modafinil, bosentan, etravirine, and nafcillin should be avoided.

#### Medicines that may have their plasma concentrations altered by brigatinib

##### CYP3A substrates

*In vitro* studies in hepatocytes have shown that brigatinib is an inducer of CYP3A4. Clinical interaction studies with sensitive CYP3A substrates have not been conducted. Brigatinib may reduce plasma levels of coadministered medicines that are predominantly metabolised by CYP3A. Therefore, coadministration of **ALUNBRIG™** with CYP3A substrates with a narrow therapeutic index (e.g., alfentanil, fentanyl, quinidine, cyclosporin, sirolimus, tacrolimus) should be avoided as their effectiveness may be reduced.

**ALUNBRIG™** may also induce other enzymes and transporters (e.g., CYP2C, P-gp) via the same mechanisms responsible for induction of CYP3A (e.g., pregnane X receptor activation).

##### Transporter substrates

Coadministration of brigatinib with substrates of P-gp, (e.g., digoxin, dabigatran, colchicine, pravastatin), BCRP (e.g., methotrexate, rosuvastatin, sulfasalazine), organic cation transporter 1 (OCT1), multidrug and toxin extrusion protein 1 (MATE1), and 2K (MATE2K) may increase their plasma concentrations. Patients should be closely monitored when **ALUNBRIG™** is coadministered with substrates of these transporters with a narrow therapeutic index (e.g., digoxin, dabigatran, methotrexate).

#### **4.6 Fertility, pregnancy and lactation**

**ALUNBRIG™** is contraindicated during pregnancy.(see section 4.3).

##### Women of childbearing potential/Contraception in males and females

Women of childbearing age being treated with **ALUNBRIG™** should be advised not to become pregnant and men being treated with **ALUNBRIG™** should be advised not to father a child during

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

treatment. Women of reproductive potential should use effective non-hormonal contraception during treatment with **ALUNBRIG™** and for at least 4 months following the final dose. Men with female partners of reproductive potential should use effective contraception during treatment and for at least 3 months after the last dose of **ALUNBRIG™**.

#### Pregnancy

**ALUNBRIG™** may cause foetal harm when administered to a pregnant woman. Studies in animals have shown reproductive toxicity (see section 5.3). There are no clinical data on the use of **ALUNBRIG™** in pregnant women. **ALUNBRIG™** should not be used during pregnancy. If **ALUNBRIG™** is used during pregnancy, or if the patient becomes pregnant while taking this medicine, the patient should be apprised of the potential hazard to a foetus.

#### Breast-feeding

Breast-feeding should be stopped during treatment with **ALUNBRIG™**.

#### Fertility

No human data on the effect of **ALUNBRIG™** on fertility are available. Based on repeat-dose toxicity studies in male animals, **ALUNBRIG™** may cause reduced fertility in males (see section 5.3).

### **4.7 Effects on ability to drive and use machines**


Caution should be exercised when driving or operating machines as patients may experience visual disturbance, dizziness, or fatigue while taking **ALUNBRIG™**.

### **4.8 Undesirable effects**

#### Summary of the safety profile

The adverse reactions described in this section were identified three clinical trials with limited patient numbers, 222 ( ALTA trial ), 275 (ALTA 1L) and 25 (Study 101).

The most common adverse reactions ( $\geq 25\%$ ) reported in patients treated with Alunbrig at the recommended dosing regimen were increased AST, increased CPK, hyperglycaemia, increased lipase, hyperinsulinaemia, diarrhoea, increased ALT, increased amylase, anaemia, nausea,

DocuSigned by:  
  
9B2F238576344AB...

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

fatigue, hypophosphataemia, decreased lymphocyte count, cough, increased alkaline phosphatase, rash, increased APTT, myalgia, headache, hypertension, decreased white blood cell count, dyspnoea, and vomiting.

The most common serious adverse reactions ( $\geq 2\%$ ) reported in patients treated with Alunbrig at the recommended dosing regimen other than events related to neoplasm progression were pneumonia, pneumonitis, dyspnoea and pyrexia.

The data described below reflect exposure to Alunbrig at the recommended dosing regimen in three clinical trials: a Phase 3 trial (ALTA 1L) in patients with advanced ALK positive NSCLC previously not treated with an ALK inhibitor (N = 136), a Phase 2 trial (ALTA) in patients treated with Alunbrig with ALK positive NSCLC who previously progressed on crizotinib (N = 110), and a phase 1/2 dose escalation/expansion trial in patients with advanced malignancies (N = 28). Across these studies, the median duration of exposure in patients receiving Alunbrig at the recommended dosing regimen was 21,8 months

Adverse reactions reported are presented in Table 3 and are listed by system organ class, preferred term and frequency. Frequency categories are very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ) and uncommon ( $\geq 1/1,000$  to  $< 1/100$ ). Within each frequency grouping, undesirable effects are presented in order of frequency.

**Table 3: Adverse reactions reported in patients treated with ALUNBRIG™ (per Common Terminology Criteria for Adverse Events (CTCAE) version 4.03) at the 180 mg regimen (N = 274)**

<b>System organ class</b>	<b>Frequency category</b>	<b>Adverse reactions† all grades</b>	<b>Adverse reactions Grade 3-4</b>
Infections and infestations	Very common	Pneumonia <sup>a,b</sup> Upper respiratory tract infection	
	Common		Pneumonia <sup>a</sup>

Blood and lymphatic system disorders	Very common	Anaemia Lymphocyte count decreased APTT increased White blood cell count decreased Neutrophil count decreased	Lymphocyte count decreased
	Common	Decreased platelet count	APTT increased Anaemia
	Uncommon		Neutrophil count decreased
Metabolism and nutrition disorders	Very common	Hyperglycaemia Hyperinsulinaemia <sup>c</sup> Hypophosphataemia Hypomagnesaemia Hypercalcaemia Hyponatraemia Hypokalaemia Decreased appetite	
	Common		Hypophosphataemia Hyperglycaemia Hyponatraemia Hypokalaemia Decreased appetite
Psychiatric disorders	Common	Insomnia	
Nervous system disorders	Very common	Headache <sup>d</sup> Peripheral neuropathy <sup>e</sup> Dizziness	
	Common	Memory impairment Dysgeusia	Headache <sup>d</sup> Peripheral neuropathy <sup>e</sup>
	Uncommon		Dizziness
Eye disorders	Very common	Visual disturbance <sup>f</sup>	
	Common		Visual disturbance <sup>f</sup>
Cardiac disorders	Common	Bradycardia <sup>g</sup> Electrocardiogram QT prolonged Tachycardia <sup>h</sup> Palpitations	Electrocardiogram QT prolonged
	Uncommon		Bradycardia <sup>g</sup>
Vascular disorders	Very common	Hypertension <sup>i</sup>	Hypertension <sup>i</sup>
Respiratory, thoracic and mediastinal disorders	Very common	Cough Dyspnoea <sup>l</sup>	
	Common	Pneumonitis <sup>k</sup>	Pneumonitis <sup>k</sup> Dyspnoea <sup>l</sup>
Gastrointestinal disorders	Very common	Lipase increased Diarrhoea <sup>t</sup> Amylase increased Nausea Vomiting Abdominal pain <sup>l</sup> Constipation Stomatitis <sup>m</sup>	Lipase increased
	Common	Dry mouth Dyspepsia Flatulence	Amylase increased Nausea Abdominal pain <sup>l</sup> Diarrhoea <sup>t</sup>

	Uncommon	Pancreatitis	Vomitin <sup>g</sup> Stomatitis <sup>m</sup> Dyspepsia Pancreatitis
Hepatobiliary disorders	Very common	AST increased ALT increased Alkaline phosphatase increased	
	Common	Blood lactate dehydrogenase increased Hyperbilirubinaemia	ALT increased AST increased Alkaline phosphatase increased
	Uncommon		Hyperbilirubinaemia
Skin and subcutaneous tissue disorders	Very common	Rash <sup>n</sup> Pruritus <sup>o</sup>	
	Common	Dry skin Photosensitivity reaction	Rash <sup>n</sup> Photosensitivity reaction
	Uncommon		Dry skin Pruritus <sup>o</sup>
Musculoskeletal and connective tissue disorders	Very common	Blood CPK increased Myalgia <sup>p</sup> Arthralgia	Blood CPK increased
	Common	Musculoskeletal chest pain Pain in extremity Musculoskeletal stiffness	
	Uncommon		Pain in extremity Musculoskeletal chest pain Myalgia <sup>p</sup>
Renal and urinary disorders	Very common	Blood creatinine increased	
General disorders and administration site conditions	Very common	Fatigue <sup>q</sup> Oedema <sup>r</sup> Pyrexia	
	Common	Non-cardiac chest pain Chest discomfort Pain	Fatigue <sup>q</sup>
	Uncommon		Pyrexia Oedema <sup>r</sup> Non-cardiac chest pain
Investigations	Common	Blood cholesterol increased <sup>s</sup> Weight decreased	
	Uncommon		Weight decreased
<p>† The frequencies for ADR terms associated with chemistry and haematology laboratory changes were determined based on the frequency of abnormal laboratory shifts from baseline.</p> <p><sup>a</sup> Includes atypical pneumonia, pneumonia, pneumonia aspiration, pneumonia cryptococcal, lower respiratory tract infection, lower respiratory tract infection viral, lung infection</p> <p><sup>b</sup> Includes Grade 5 events</p> <p><sup>c</sup> Grade not applicable</p> <p><sup>d</sup> Includes headache, sinus headache, head discomfort, migraine, tension headache</p> <p><sup>e</sup> Includes paraesthesia, peripheral sensory neuropathy, dysaesthesia, hyperaesthesia, hypoaesthesia, neuralgia, neuropathy peripheral, neurotoxicity, peripheral motor neuropathy, polyneuropathy, burning sensation, post herpetic neuralgia</p> <p><sup>f</sup> Includes altered visual depth perception, cataract, colour blindness acquired, diplopia, glaucoma, intraocular pressure increased, macular oedema, photophobia, photopsia, retinal oedema, vision blurred, visual acuity reduced, visual field defect, visual impairment, vitreous detachment, vitreous floaters, amaurosis fugax</p> <p><sup>g</sup> Includes bradycardia, sinus bradycardia</p> <p><sup>h</sup> Includes sinus tachycardia, tachycardia, atrial tachycardia, heart rate increased</p>			

<sup>i</sup> Includes blood pressure increased, diastolic hypertension, hypertension, systolic hypertension  
<sup>j</sup> Includes dyspnoea, dyspnoea exertional  
<sup>k</sup> Includes interstitial lung disease, pneumonitis  
<sup>l</sup> Includes abdominal discomfort, abdominal distension, abdominal pain, abdominal pain lower, abdominal pain upper, epigastric discomfort.  
<sup>m</sup> Includes aphthous stomatitis, stomatitis, aphthous ulcer, mouth ulceration, oral mucosal blistering  
<sup>n</sup> Includes dermatitis acneiform, erythema, exfoliative rash, rash, rash erythematous, rash macular, rash maculo-papular, rash papular, rash pruritic, rash pustular, dermatitis, dermatitis allergic, dermatitis contact, generalised erythema, rash follicular, urticaria, drug eruption, toxic skin eruption  
<sup>o</sup> Includes pruritus, pruritus allergic, pruritus generalised, pruritus genital, vulvovaginal pruritus  
<sup>p</sup> Includes musculoskeletal pain, myalgia, muscle spasms, muscle tightness, muscle twitching, musculoskeletal discomfort  
<sup>q</sup> Includes asthenia, fatigue  
<sup>r</sup> Includes eyelid oedema, face oedema, oedema peripheral, periorbital oedema, swelling face, generalised oedema, peripheral swelling, angioedema, lip swelling, periorbital swelling, skin swelling, swelling of eyelid  
<sup>s</sup> Includes blood cholesterol increased, hypercholesterolemia  
<sup>t</sup> diarrhoea

### Description of selected adverse reactions

#### *Pulmonary adverse reactions*

*In ALTA 1L, 2,9 % of patients experienced any Grade ILD/pneumonitis early in treatment (within 8 days), with Grade 3 4 ILD/pneumonitis in 2,2 % of patients. There were no fatal ILD/pneumonitis. Additionally, 3,7 % of patients experienced pneumonitis later in treatment.*

In ALTA, 6,4 % of patients experienced pulmonary adverse reactions of any grade, including ILD/pneumonitis, pneumonia and dyspnoea, early in treatment (within 9 days, median onset: 2 days); 2,7 % of patients had Grade 3-4 pulmonary adverse reactions and 1 patient (0,5 %) had fatal pneumonia.

Following Grade 1-2 pulmonary adverse reactions, treatment with **ALUNBRIG™** was either interrupted and then restarted or the dose was reduced. Early pulmonary adverse reactions also occurred in a dose escalation study in patients (N = 137) (Study 101) including three fatal cases (hypoxia, acute respiratory distress syndrome and pneumonia).

Additionally, 2,3 % of patients in ALTA experienced pneumonitis later in treatment, with 2 patients having Grade 3 pneumonitis (see sections 4.2 and 4.4).

#### Elderly

Early pulmonary adverse reaction was reported in 10,1 % of patients ≥ 65 years of age compared with 3,1 % of patients < 65 years of age.

### Hypertension

Hypertension was reported in 30 % of patients treated with ALUNBRIG™ at the 180 mg regimen with 11 % having Grade 3 hypertension. Dose reduction for hypertension occurred in 1,5 % at the 180 mg regimen. Mean systolic and diastolic blood pressure, in all patients, increased over time (see sections 4.2 and 4.4).

### Bradycardia

Bradycardia was reported in 8,4 % of patients treated with ALUNBRIG™ at the 180 mg regimen.

Heart rates of less than 50 beats per minute (bpm) were reported in -8,4 % of patients at the 180 mg regimen. (see sections 4.2 and 4.4).

### Visual disturbance

Visual disturbance adverse reactions were reported in 14 % of patients treated with ALUNBRIG™ at the 180 mg regimen. Of these, three Grade 3 adverse reactions (1,1 %) including macular oedema and cataract were reported.

Dose reduction for visual disturbance occurred in two patients (0,7 %) at the 180 mg regimen (see sections 4.2 and 4.4).

### Peripheral neuropathy

Peripheral neuropathy adverse reactions were reported in 20 % of patients treated at the 180 mg regimen. Thirty-threepercent of patients had resolution of all peripheral neuropathy adverse reactions. The median duration of peripheral neuropathy adverse reactions was 6,6 months, with a maximum duration of 28,9 months.

### Creatine phosphokinase (CPK) elevation

In ALTA 1L and ALTA, elevations of CPK were reported in 64 % of patients treated with ALUNBRIG™ at the 180 mg regimen. The incidence of Grade 3-4 elevations of CPK was 18 %. The median time to onset for CPK elevations was 28 days.

Dose reduction for CPK elevation occurred in 10 % patients at the 180 mg regimen (see

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd  
sections 4.2 and 4.4).

#### Elevations of pancreatic enzymes

In ALTA, elevations of amylase and lipase were reported in 47 % and 54 % of patients treated with **ALUNBRIG™**, respectively at the 180 mg regimen. For elevations to Grade 3 and 4, the incidences for amylase and lipase were 7,7 % and 15 %, respectively. The median time to onset for amylase elevations and lipase elevations was 17 days and 29 days, respectively.

Dose reduction for elevation of lipase and amylase occurred in 4,7 % and 2,9 % of patients, respectively at the 180 mg regimen (see sections 4.2 and 4.4).

#### Elevation of hepatic enzymes

Elevations of ALT and AST were reported in 49 % and 68 % of patients treated with **ALUNBRIG™**, respectively at the 180 mg regimen.

For elevations to Grade 3 and 4, the incidences for ALT and AST were 4,7 % and 3,6 %, respectively.

Dose reduction for elevation of ALT and AST occurred in 0,7% and 1,1% of patients, respectively at the 180 mg regimen (see sections 4.2 and 4.4).

#### Hyperglycaemia

Sixty one percent of patients experienced hyperglycaemia. Grade 3 hyperglycemia occurred in 6,6 % of patients.


No patients had dose reductions due to hyperglycaemia.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important.

It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers are asked to report any suspected adverse reactions to SAHPRA via “**6.04 Adverse Drug Reactions**

**Reporting Form**” found online under SAHPRA’s publications:

DocuSigned by:  
  
9B2F238576344AB...

#### 4.9 Overdose

There is no specific antidote for overdose with **ALUNBRIG™**. In the event of an overdose, monitor the patient for adverse reactions (see section 4.8) and provide appropriate supportive care.

### 5. PHARMACOLOGICAL PROPERTIES

#### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antineoplastic agent, protein kinase inhibitors, ATC code: L01XE43

##### Mechanism of action

Brigatinib is a tyrosine kinase inhibitor that targets ALK, c-ros oncogene 1 (ROS1), and insulin-like growth factor 1 receptor (IGF-1R). Brigatinib inhibited autophosphorylation of ALK and ALK-mediated phosphorylation of the downstream signalling protein STAT3 in *in vitro* and *in vivo* assays.

Brigatinib inhibited the *in vitro* proliferation of cell lines expressing EML4-ALK and NPM-ALK fusion proteins and demonstrated dose-dependent inhibition of EML4-ALK-positive NSCLC xenograft growth in mice. Brigatinib inhibited the *in vitro* and *in vivo* viability of cells expressing mutant forms of EML4-ALK associated with resistance to ALK inhibitors, including G1202R and L1196M.

##### *Cardiac electrophysiology*

*In Study 101, the QT interval prolongation potential of brigatinib was assessed in 123 patients with advanced malignancies following once daily brigatinib doses of 30 mg to 240 mg. The maximum mean QTcF (corrected QT by the Fridericia method) change from baseline was less than 10 msec.*

*An exposure QT analysis suggested no concentration dependent QTc interval prolongation.*

*In 274 patients treated with brigatinib 180 mg once a day QT prolongation was reported as common adverse event. (see Table 3).*

##### *Clinical efficacy and safety*

###### ALTA 1L

*The safety and efficacy of Alunbrig was evaluated in a randomised (1:1), open label, multicentre trial (ALTA 1L) in 275 adult patients with advanced ALK positive NSCLC who had not previously received*

an ALK targeted therapy. Patients were allowed to have up to 1 prior regimen of chemotherapy in the locally advanced or metastatic setting.

Patients were randomised in a 1:1 ratio to receive Alunbrig 180 mg once daily with a 7 day lead in at 90 mg once daily (N = 137) or crizotinib 250 mg orally twice daily (N = 138).

The primary endpoint measure was progression free survival (PFS) according to Response Evaluation Criteria in Solid Tumours (RECIST v1.1) as evaluated by a Blinded Independent Review Committee (BIRC). Additional outcome measures included confirmed objective response rate (ORR), duration of response (DOR), time to response, disease control rate (DCR), intracranial ORR, intracranial PFS, and intracranial DOR.

At the primary analysis performed at a median follow-up duration of 11 months in the Alunbrig arm, the ALTA 1L study met its primary endpoint demonstrating a statistically significant improvement in PFS by BIRC. The second interim analysis performed at a median follow up duration of 24,9 months in the Alunbrig arm formed the basis for the results from this study (Table 4).

**Table 4: Efficacy Results in ALTA IL (second interim analysis in ITT Population)**

<b><u>Efficacy Parameters</u></b>	<b><u>Alunbrig N = 137</u></b>	<b><u>Crizotinib N = 138</u></b>
<b><u>Median duration of follow-up (months)</u></b>	<u>24,9</u> (range: 0–34,1)	<u>15,2</u> (range: 0,1–36)
<b><u>Primary efficacy parameters</u></b>		
<b><u>PFS (BIRC)</u></b>		
<u>Number of Patients with Events, n (%)</u>	<u>63 (46 %)</u>	<u>87 (63 %)</u>
<u>Progressive Disease, n (%)</u>	<u>56 (40,9 %)<sup>a</sup></u>	<u>82 (59,4 %)<sup>b</sup></u>
<u>Death, n (%)</u>	<u>7 (5,1 %)</u>	<u>5 (3,6 %)</u>
<u>Median (in months) (95 % CI)</u>	<u>24 (18,5, NE)</u>	<u>11 (9,2; 12,9)</u>
<u>Hazard ratio (95 % CI)</u>	<u>0.49 (0.35; 0.68)</u>	
<u>Log-rank p-value<sup>c</sup></u>	<u>&lt;0.0001</u>	
<b><u>Secondary efficacy parameters</u></b>		
<b><u>Confirmed Objective Response Rate (BIRC)</u></b>		
<u>Responders, n (%) (95 % CI)</u>	<u>101 (73,7 %)</u> (65,5; 80,9)	<u>85 (61,6 %)</u> (52,9; 69,7)
<u>p-value<sup>c,d</sup></u>	<u>0,0342</u>	
<u>Complete Response, %</u>	<u>14,6 %</u>	<u>8,7 %</u>
<u>Partial Response, %</u>	<u>59,1 %</u>	<u>52,9 %</u>

<b>Duration of Confirmed Response (BIRC)</b>		
Median (months) (95 % CI)	NE (19,4; NE)	13,8 (9,3; 20,8)
<b>Overall Survival</b>		
Number of Events, n (%)	33 (24,1)	37 (26,8)
Median (in months) (95 % CI)	NE (NE; NE)	NE (NE; NE)
Hazard ratio (95 % CI)	0.92 (0,57; 1,47)	
Log-rank p-value <sup>d</sup>	0,7710	

BIRC = Blinded Independent Review Committee; NE = Not Estimable; CI = Confidence Interval

a includes 2 patients with palliative radiotherapy to the brain

b includes 8 patients with palliative radiotherapy to the brain

c Stratified by presence of iCNS metastases at baseline and prior chemotherapy for locally advanced or metastatic disease for log-rank test and Cochran Mantel-Haenszel test, respectively

d From a Cochran Mantel-Haenszel test

BIRC assessment of intracranial efficacy according to RECIST v1.1 in patients with any brain metastases and patients with measurable brain metastases ( $\geq 10$  mm in longest diameter) at baseline are summarised in Table 5.

**Table 5: Intracranial Efficacy in Patients in ALTA 1L**

Efficacy Parameters	Patients with Measurable Brain Metastases at Baseline	
	Alunbrig N = 18	Crizotinib N = 23
<b>Confirmed Intracranial Objective Response Rate</b>		
Responders, n (%) (95 % CI)	14 (77,8 %) (52,4; 93,6)	6 (26,1 %) (10,2; 48,4)
p-value <sup>a,b</sup>	0,0014	
Complete Response %	27,8 %	0
Partial Response %	50 %	26,1 %
<b>Duration of Confirmed Intracranial Response<sup>c</sup></b>		
Median (months) (95 % CI)	NE (5,7; NE)	9,2 (3,9; 9,2)
	Patients with Any Brain Metastases at Baseline	
	Alunbrig N = 47	Crizotinib N = 49
<b>Confirmed Intracranial Objective Response Rate</b>		
Responders, n (%) (95 % CI)	31 (66 %) (50,7; 79,1)	8 (16,3 %) (7,32; 29,7)
p-value <sup>a,b</sup>	< 0,0001	
Complete Response (%)	44,7 %	4,1 %
Partial Response (%)	21,3 %	12,2 %
<b>Duration of Confirmed Intracranial Response<sup>c</sup></b>		
Median (months) (95 % CI)	24 (16,9; NE)	9,2 (3,9; NE)
<b>Intracranial PFS<sup>d</sup></b>		
Number of Patients with Events, n (%)	21 (44,7 %)	32 (65,3 %)
Progressive Disease, n (%)	21 (44,7 %) <sup>e</sup>	29 (59,2 %) <sup>f</sup>
Death, n (%)	0	3 (6,1 %)

Median (in months) (95 % CI)	24 (13, NE)	5,6 (3,7; 7,5)
Hazard ratio (95 % CI)	0,31 (0,17; 0,56)	
Log-rank p-value	< 0,0001	

BIRC = Blinded Independent Review Committee CI = Confidence Interval;  
NE = Not Estimable

<sup>a</sup> Stratified by presence prior chemotherapy for locally advanced or metastatic disease for log-rank test and Cochran Mantel-Haenszel test, respectively

<sup>b</sup> From a Cochran Mantel-Haenszel test

<sup>c</sup> measured from date of first confirmed intracranial response until date of intracranial disease progression (new intracranial lesions, intracranial target lesion diameter growth  $\geq$  20% from nadir, or unequivocal progression of intracranial nontarget lesions) or death or censoring

<sup>d</sup> measured from date of randomisation until date of intracranial disease progression (new intracranial lesions, intracranial target lesion diameter growth  $\geq$  20% from nadir, or unequivocal progression of intracranial nontarget lesions) or death or censoring.

<sup>e</sup> includes 1 patient with palliative radiotherapy to the brain

<sup>f</sup> includes 2 patients with palliative radiotherapy to the brain

## ALTA

The safety and efficacy of Alunbrig was evaluated in a randomised (1:1), open label, multicenter trial (ALTA) in 222 adult patients with locally advanced or metastatic ALK positive NSCLC who had progressed on crizotinib.

Patients were randomised in a 1:1 ratio to receive Alunbrig either 90 mg once daily (90 mg regimen, N = 112) or 180 mg once daily with 7 day lead in at 90 mg once daily (180 mg regimen, N = 110).

The major outcome measure was confirmed objective response rate (ORR) according to Response Evaluation Criteria in Solid Tumours (RECIST v1.1) as evaluated by investigator. Additional outcome measures included confirmed ORR as evaluated by an Independent Review Committee (IRC); time to response; progression free survival (PFS); duration of response (DOR); overall survival; and intracranial ORR and intracranial DOR as evaluated by an IRC.

Efficacy results from ALTA analysis are summarised in Table 6.

Table 6: Efficacy results in ALTA (ITT population)

Efficacy parameter	Investigator assessment		IRC assessment	
	90 mg regimen* N = 112	180 mg regimen† N = 110	90 mg regimen* N = 112	180 mg regimen† N = 110
<b>Objective response rate</b>				
(%)	46 %	56 %	51 %	56 %
CI‡	(35; 57)	(45; 67)	(41; 61)	(47; 66)

<b>Time to response</b>				
Median (months)	1,8	1,9	1,8	1,9
<b>Duration of response</b>				
Median (months)	12,0	13,8	16,4	15,7
95 % CI	(9,2;17,7)	(10,2;19,3)	(7,4; 24,9)	(12,8; 21,8)
<b>Progression-free survival</b>				
Median (months)	9,2	15,6	9,2	16,7
95 % CI	(7,4; 11,1)	(11,1; 21)	(7,4; 12,8)	(11,6; 21,4)
<b>Overall survival</b>				
Median (months)	29,5	34,1	NA	NA
95 % CI	(18,2; NE)	(27,7; NE)	NA	NA
12-month survival probability (%)	70,3 %	80,1 %	NA	NA

CI = Confidence Interval; NE = Not Estimable; NA = Not Applicable

\*90 mg once daily regimen

†180 mg once daily with 7 day lead in at 90 mg once daily

‡Confidence Interval for investigator assessed ORR is 97,5 % and for IRC assessed ORR is 95 %

In patients with any brain metastases at baseline, intracranial disease control rate was 77,8 % (95 % CI 67,2-86,3) in the 90 mg arm (N = 81) and 85,1 % (95 % CI 75-92,3) in the 180 mg arm (N = 74).

#### Paediatric population


(see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

### Absorption

Following administration of a single oral dose of brigatinib (30-240 mg) in patients with NSCLC, the median time to peak concentration ( $T_{max}$ ) was 1-4 hours postdose. After a single dose and at steady state, systemic exposure was dose proportional over the dose range of 60-240 mg once daily. Modest accumulation was observed upon repeated dosing (geometric mean accumulation ratio: 1,9 to 2,4). The geometric mean steady state  $C_{max}$  of brigatinib at doses of 90 mg and 180 mg once daily was 552 and 1,452 ng/mL, respectively, and the corresponding  $AUC_{0-t}$  was 8,165 and 20,276 h·ng/mL, respectively. Brigatinib is a substrate of the transporter proteins P-gp and BCRP.

In healthy subjects, compared to overnight fasting, a high fat meal reduced brigatinib  $C_{max}$  by 13 % with no effect on AUC. Brigatinib can be administered with or without food.

DocuSigned by:  
  
 9B2F238576344AB...

### Distribution

Brigatinib was moderately bound (91 %) to human plasma proteins and binding was not concentration-dependent. The blood-to-plasma concentration ratio is 0,69. In patients given brigatinib 180 mg once daily, the geometric mean apparent volume of distribution ( $V_z/F$ ) of brigatinib at steady state was 307453 L, indicating moderate distribution into tissues.

### Biotransformation

*In vitro* studies demonstrated that brigatinib is primarily metabolised by CYP2C8 and CYP3A4, and to a much lesser extent by CYP3A5.

Following oral administration of a single 180 mg dose of [<sup>14</sup>C]brigatinib to healthy subjects, N-demethylation and cysteine conjugation were the two major metabolic clearance pathways. In urine and faeces combined, 48 %, 27 %, and 9,1 % of the radioactive dose was excreted as unchanged brigatinib, N-desmethyl brigatinib (AP26123), and brigatinib cysteine conjugate, respectively. Unchanged brigatinib was the major circulating radioactive component (92 %) along with AP26123 (3,5 %), the primary metabolite also observed *in vitro*. In patients, at steady state, the plasma AUC of AP26123 was < 10 % of brigatinib exposure. In *in vitro* kinase and cellular assays, the metabolite, AP26123, inhibited ALK with approximately 3-fold lower potency than brigatinib.

### Elimination

In patients given brigatinib 180 mg once daily, the geometric mean apparent oral clearance (CL/F) of brigatinib at steady state was 8,9 L/h and the median plasma elimination half-life was 24 h.

The primary route of excretion of brigatinib is in faeces. In six healthy male subjects given a single 180 mg oral dose of [<sup>14</sup>C]brigatinib, 65 % of the administered dose was recovered in faeces and 25 % of the administered dose was recovered in urine. Unchanged brigatinib represented 41 % and 86 % of the total radioactivity in faeces and urine, respectively, the remainder being metabolites.

### Specific populations

#### Hepatic impairment

The pharmacokinetics of brigatinib was characterised in healthy subjects with normal hepatic function

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

(N = 9), and patients with mild hepatic impairment (Child-Pugh class A, N = 6), moderate hepatic impairment (Child-Pugh class B, N = 6), or severe hepatic impairment (Child-Pugh class C, N = 6). The pharmacokinetics of brigatinib was similar between healthy subjects with normal hepatic function and patients with mild (Child-Pugh class A) or moderate (Child-Pugh class B) hepatic impairment. Unbound AUC<sub>0-INF</sub> was 37 % higher in patients with severe hepatic impairment (Child-Pugh class C) as compared to healthy subjects with normal hepatic function (see section 4.2).

### Renal impairment

The pharmacokinetics of brigatinib is similar in patients with normal renal function and in patients with mild or moderate renal impairment (eGFR  $\geq$  30 mL/min) based on the results of population pharmacokinetic analyses. In a pharmacokinetic study, unbound AUC<sub>0-INF</sub> was 94 % higher in patients with severe renal impairment (eGFR < 30 mL/min, N = 6) as compared to patients with normal renal function (eGFR  $\geq$  90 mL/min, N = 8) (see section 4.2).

### **5.3 Preclinical safety data**

Safety pharmacology studies with brigatinib identified potential for pulmonary effects (altered respiration rate; 1-2 times the human C<sub>max</sub>), cardiovascular effects (altered heart rate and blood pressure; at 0.5 times the human C<sub>max</sub>), and renal effects (reduced renal function; at 1-2,5 times the human C<sub>max</sub>), but did not indicate any potential for QT prolongation or neurofunctional effects.

Adverse reactions seen in animals at exposure levels similar to clinical exposure levels with possible relevance to clinical use were as follows: gastrointestinal system, bone marrow, eyes, testes, liver, kidney, bone, and heart. These effects were generally reversible during the non-dosing recovery period; however, effects in the eyes and testes were notable exceptions due to lack of recovery.

Carcinogenicity studies have not been performed with brigatinib.

Brigatinib may impair male fertility. Testicular toxicity was observed in repeat-dose animal studies. In rats, findings included lower weight of testes, seminal vesicles and prostate gland, and testicular tubular degeneration; these effects were not reversible during the recovery period. In monkeys, findings included reduced size of testes along with microscopic evidence of hypospermatogenesis; these effects were reversible during the recovery period. Overall, these effects on the male

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

reproductive organs in rats and monkeys occurred at exposures  $\geq 0,2$ -times the AUC observed in patients at the 180 mg once daily dose. No apparent adverse effects on female reproductive organs were observed in general toxicology studies in rats and monkeys.

In an embryo-foetal development study in which pregnant rats were administered daily doses of brigatinib during organogenesis; dose-related skeletal anomalies were observed at doses as low as approximately 0,7-times the human exposure by AUC at the 180 mg once daily dose. Findings included embryo-lethality, reduced foetal growth, and skeletal variations.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Lactose monohydrate

Microcrystalline cellulose

Sodium starch glycolate (type A)

Silica colloidal hydrophobic

Magnesium stearate

#### Tablet coating

Talc

Macrogol

Polyvinyl alcohol

Titanium dioxide


### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

#### **ALUNBRIG™ 30 mg film-coated tablets**

3 years

DocuSigned by:  
  
9B2F238576344AB...

**ALUNBRIG™ 90 mg film-coated tablets**

3 years

**ALUNBRIG™ 180 mg film-coated tablets**

3 years

**6.4 Special precautions for storage**

Store at or below 30 °C. KEEP OUT OF REACH OF CHILDREN.

HDPE bottle: Keep bottle tightly closed.

Blisters: Keep blisters enclosed in carton until required for use.

**6.5 Nature and contents of container**

**ALUNBRIG™ 30 mg film-coated tablets**

White round wide mouth high density polyethylene (HDPE) bottles with white two-piece polypropylene child resistant screw cap closures with foil induction seal liner, containing either 60 or 120 film-coated tablets, together with one HDPE canister containing a molecular sieve desiccant.

Clear thermoformable Aclar® blister with heat sealable paper-laminated foil lidding in a carton, containing either 28, 56 or 112 film-coated tablets.


**ALUNBRIG™ 90 mg film-coated tablets**

White round wide mouth high density polyethylene (HDPE) bottles with white two-piece polypropylene child resistant screw cap with foil induction seal liner closures, containing either 7 or 30 film-coated tablets, together with one HDPE canister containing a molecular sieve desiccant.

Clear thermoformable Aclar® blister with heat sealable paper-laminated foil lidding in a carton, containing either 7 or 28 film-coated tablets.

**ALUNBRIG™ 180 mg film-coated tablets**

White round wide mouth high density polyethylene (HDPE) bottles with white two-piece polypropylene child resistant screw cap with foil induction seal liner closures, containing

DocuSigned by:  
  
9B2F238576344AB...

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd  
30 film-coated tablets, together with one HDPE canister containing a molecular sieve desiccant.

Clear thermoformable Aclar® blister with heat sealable paper-laminated foil lidding in a carton, containing 28 film-coated tablets.

Treatment initiation pack **ALUNBRIG™** 90 mg and 180 mg film-coated tablets

Each pack consists of an outer carton with two inner cartons containing:

- **ALUNBRIG™** 90 mg film-coated tablets  
1 clear thermoformable Aclar® blister with heat sealable paper-laminated foil lidding in a carton, containing 7 film-coated tablets.
- **ALUNBRIG™** 180 mg film-coated tablets  
3 clear thermoformable Aclar® blisters with heat sealable paper-laminated foil lidding in a carton, containing 21 film-coated tablets.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

Patients should be advised to keep the desiccant canister in the bottle and not to swallow it.

Any unused medicine or waste material should be disposed of in accordance with local requirements

## **7. HOLDER OF CERTIFICATE OF REGISTRATION**


**TAKEDA (Pty) Ltd**

Building A, Monte Circle  
64 Montecasino Boulevard  
Fourways, 2191  
South Africa

## **8. REGISTRATION NUMBER(S)**

**ALUNBRIG™** 30 mg film-coated tablets

52/26/0353

DocuSigned by:  
  
9B2F238576344AB...

**ALUNBRIG™** 30 mg, **ALUNBRIG™** 90 mg and **ALUNBRIG™** 180 mg (brigatinib film-coated Tablets) Takeda (Pty) Ltd

**ALUNBRIG™ 90 mg film-coated tablets**

52/26/0354

**ALUNBRIG™ 180 mg film-coated tablets**

52/26/0355

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

24 July 2020.

**10. DATE OF REVISION OF THE TEXT**

17 February 2022