

1.3.2 PATIENT INFORMATION LEAFLET

Patient Information leaflet

SCHEDULING STATUS

S4

DESCOVY 200 mg/10 mg film-coated tablets

DESCOVY 200 mg/25 mg film-coated tablets

emtricitabine/tenofovir alafenamide

Sugar Free

WARNING:

TOO MUCH LACTIC ACID IN YOUR BLOOD (LACTIC ACIDOSIS) IS A SERIOUS BUT RARE MEDICAL EMERGENCY THAT CAN LEAD TO DEATH. TELL YOUR HEALTHCARE PROVIDER RIGHT AWAY IF YOU GET THESE SYMPTOMS: WEAKNESS OR BEING TIRED MORE THAN USUAL, UNUSUAL MUSCLE PAIN, BEING SHORT OF BREATH OR FAST BREATHING, STOMACH PAIN WITH NAUSEA AND VOMITING, COLD OR BLUE HANDS AND FEET, FEEL DIZZY OR LIGHTHEADED, OR A FAST OR ABNORMAL HEARTBEAT.

IF YOU HAVE HEPATITIS B INFECTION, YOUR DOCTOR WILL CAREFULLY CONSIDER THE BEST TREATMENT REGIMEN FOR YOU. IF YOU HAVE HEPATITIS B INFECTION, LIVER PROBLEMS MAY BECOME WORSE AFTER YOU STOP TAKING DESCOVY. IT IS IMPORTANT NOT TO STOP TAKING DESCOVY WITHOUT TALKING TO YOUR DOCTOR. YOU MUST REMAIN UNDER THE CARE OF YOUR DOCTOR WHILE TAKING DESCOVY.

Read all of this leaflet carefully before you start taking DESCOVY because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist, nurse or healthcare provider.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours

What is in this leaflet

1. What DESCOVY is and what it is used for
2. What you need to know before you take DESCOVY
3. How to take DESCOVY
4. Possible side effects
5. How to store DESCOVY
6. Contents of the pack and other information

1. What DESCOVY is and what it is used for

DESCOVY contains two active substances:

- **emtricitabine**, an antiretroviral medicine of a type known as a nucleoside reverse transcriptase inhibitor (NRTI)
- **tenofovir alafenamide**, an antiretroviral medicine of a type known as a nucleotide reverse transcriptase inhibitor (NtRTI)

DESCOVY blocks the action of the reverse transcriptase enzyme, which is essential for the virus to multiply. DESCOVY therefore reduces the amount of HIV in your body.

DESCOVY in combination with other medicines is for the **treatment of human immunodeficiency virus 1 (HIV-1) infection** in adults and adolescents 12 years of age and older, who weigh at least 35 kg.

2. What you need to know before you take DESCOVY

Do not take DESCOVY:

If you are allergic to emtricitabine, tenofovir alafenamide or any of the other ingredients of this medicine (listed in section 6 of this leaflet).

Warnings and precautions

Too much lactic acid in your blood (lactic acidosis) is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you get these symptoms: weakness or being more tired than usual, unusual muscle pain, being short of breath or fast breathing, stomach pain with nausea and vomiting, cold or blue hands and feet, feel dizzy or lightheaded, or a fast or abnormal heartbeat.

If you have **hepatitis B infection**, your doctor will carefully consider the best treatment regimen for you. If you have hepatitis B infection, liver problems may become worse after you stop taking DESCOVY. It is important not to stop taking DESCOVY without talking to your doctor.

You must remain under the care of your doctor while taking DESCOVY.

You can still pass on HIV when taking DESCOVY. Discuss with your doctor the precautions needed to avoid infecting other people. DESCOVY is not a cure for HIV infection. While taking DESCOVY you may still develop infections or other illnesses associated with HIV infection.

During HIV therapy such as with DESCOVY there may be **an increase in weight and in levels of blood lipids and glucose**. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Autoimmune disorders (the immune system attacks healthy body tissue), may also occur after you start taking medicines for HIV infection such as DESCOVY. Autoimmune disorders may occur many months after the start of treatment. Look out for any symptoms of infection or other symptoms such as:

- muscle weakness
- weakness beginning in the hands and feet and moving up towards the trunk of the body
- palpitations, tremor or hyperactivity

Talk to your doctor before taking DESCOVY:

If you have liver problems or have suffered liver disease, including hepatitis.

- Patients with liver disease including chronic hepatitis B or C, who are treated with antiretrovirals, have a higher risk of severe and potentially fatal liver complications. If you have hepatitis B infection, your doctor will carefully consider the best treatment regimen for you.
- If you have hepatitis B liver problems it may become worse after you stop taking DESCOVY. Do not stop taking DESCOVY without talking to your doctor: see section 3, *Do not stop taking DESCOVY*.
- Your doctor may not prescribe DESCOVY to you if the virus has a K65R mutation.

While you are taking DESCOVY

Once you start taking DESCOVY, look out for:

- **Signs of inflammation or infection**
- Any signs of inflammation or infection. In some patients with advanced HIV infection (AIDS) and who have had opportunistic infections in the past (infections that occur in people with a weak

immune system), signs and symptoms of inflammation from previous infections may occur soon after antiretroviral treatment, such as with DESCOVY, is started. It is thought that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.

- **Joint pain, stiffness or bone problems**

- Bone problems. Some patients taking combination antiretroviral medicines such as Descovy may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). Taking this type of medicine for a long time, taking corticosteroids, drinking alcohol, having a very weak immune system, and being overweight, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are:
 - o joint stiffness
 - o joint aches and pains (especially of the hip, knee and shoulder)
 - o difficulty with movement

→ **If you notice any of these symptoms, tell your doctor immediately.** For more information see section 4, *Possible side effects*.

Although kidney problems have not been observed with DESCOVY, there is a possibility that you may experience kidney problems when taking DESCOVY over a long period of time.

Children and adolescents

Do not give this medicine to children under 12 years of age, or weighing less than 35 kg. The use of DESCOVY in children under 12 years of age has not been studied.

Other medicines and DESCOVY

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. DESCOVY may interact with other medicines. As a result, the amounts of DESCOVY or other medicines in your blood may change. This may stop your medicines from working properly, or may make any side effects worse. In some cases, your doctor may need to adjust your dose or check your blood levels.

Medicines used in treating hepatitis B infection:

You should not take DESCOVY with medicines containing:

- **tenofovir alafenamide**
- **tenofovir disoproxil**
- **lamivudine**

- **adefovir dipivoxil**
- **Tell your doctor** if you are taking any of these medicines.

Other types of medicine:

Talk to your doctor if you are taking:

antibiotics, used to treat bacterial infections including tuberculosis, containing:

- rifabutin, rifampicin, and rifapentine

antiviral medicines used to treat HIV:

- emtricitabine and tipranavir

anticonvulsants, used to treat epilepsy, such as:

- carbamazepine, oxcarbazepine, phenobarbitone and phenytoin

herbal remedies used to treat depression and anxiety containing:

- St. John's wort (*Hypericum perforatum*)

→ **Tell your doctor if you are taking these or any other medicines.** Do not stop your treatment without contacting your doctor.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking DESCOVY.

Use effective contraception while taking DESCOVY.

Ask your doctor or pharmacist for advice before taking any medicine when pregnant.

If you have taken DESCOVY during your pregnancy, your doctor may request regular blood tests and other diagnostic tests to monitor the development of your child. In children whose mothers took NRTIs during pregnancy, the benefit from the protection against HIV outweighed the risk of side effects.

Do not breast-feed your baby during treatment with DESCOVY.

One of the active substances in DESCOVY (emtricitabine) passes into breast milk and possible harm to your baby cannot be excluded.

It is also recommended that you do not breast-feed your baby to avoid passing the HIV virus to the baby through your breast milk.

Driving and using machines

Descovy may affect your ability to drive and use machines. Do not drive and use machines until you know how treatment with DESCOVY affects you.

DESCOVY can cause dizziness and fatigue/tiredness. If you feel tired and/or dizzy when taking DESCOVY, do not drive and do not use any tools or machines.

3. How to take DESCOVY

Do not share medicines prescribed for you with any other person.

Always take DESCOVY exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is:

Adults: one tablet each day, with or without food

Adolescents 12 years of age and older, who weigh at least 35 kg: one tablet each day with or without food

Do not chew, crush or split the tablet.

Always take the dose recommended by your doctor. This is to make sure that your medicine is fully effective, and to reduce the risk of developing resistance to the treatment. Do not change the dose unless your doctor tells you to.

If you are on dialysis, take your daily dose of DESCOVY following completion of dialysis.

If you take more DESCOVY than you should

If you take more than the recommended dose of DESCOVY you may be at higher risk of side effects of DESCOVY (see section 4, *Possible side effects*).

Contact your doctor or nearest emergency department immediately for advice. Keep the tablet bottle with you so that you can show what you have taken.

If you forget to take DESCOVY

It is important not to miss a dose of DESCOVY.

If you do miss a dose:

If you notice within 18 hours of the time you usually take DESCOVY, you must take the tablet as soon as possible. Then take the next dose as usual.

If you notice 18 hours or more after the time you usually take DESCOVY, then do not take the missed dose. Wait and take the next dose at your usual time.

If you vomit less than 1 hour after taking DESCOVY, take another tablet.

Do not stop taking DESCOVY

Do not stop taking DESCOVY without talking to your doctor. Stopping DESCOVY can seriously affect how well future treatment works. If DESCOVY is stopped for any reason, speak to your doctor before you restart taking DESCOVY tablets.

When your supply of DESCOVY starts to run low, get more from your doctor or pharmacist. This is very important because the amount of virus may start to increase if DESCOVY is stopped for even a short time. The disease may then become harder to treat.

If you have both HIV infection and hepatitis B, it is very important not to stop taking DESCOVY without talking to your doctor first. You may require blood tests for several months after stopping treatment. In some patients with advanced liver disease or cirrhosis, stopping treatment may lead to worsening of hepatitis, which may be life-threatening.

→ **Tell your doctor immediately** about new or unusual symptoms after you stop treatment, particularly symptoms you associate with hepatitis B infection.

If you have any further questions on the use of DESCOVY, ask your doctor or pharmacist.

4. Possible side effects

DESCOVY can have side effects.

Not all side effects reported for DESCOVY are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking DESCOVY, please consult your doctor, pharmacist or other healthcare professional for advice.

Possible serious side effects: tell a doctor immediately

Any signs of inflammation or infection. In some patients with advanced HIV infection (AIDS) and who have had opportunistic infections in the past (infections that occur in people with a weak immune system), signs and symptoms of inflammation from previous infections may occur soon after antiretroviral treatment such as with DESCOVY is started. It is thought that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.

- **Autoimmune disorders** (the immune system attacks healthy body tissue), may also occur after you start taking medicines for HIV infection such as DESCOVY. Autoimmune disorders may occur many months after the start of treatment. Look out for any symptoms of infection or other symptoms such as:
 - muscle weakness
 - weakness beginning in the hands and feet and moving up towards the trunk of the body

- palpitations, tremor or hyperactivity

→ If you notice the side effects described above, tell your doctor immediately.

Frequent side effects

feeling sick (*nausea*)

abnormal dreams

headache

dizziness

diarrhoea

vomiting

stomach pain

wind (*flatulence*)

rash

tiredness (*fatigue*)

Less frequent side effects

swelling of the face, lips or throat with difficult breathing (*angioedema*)

These are all very serious side effects. If you have them, you may have had a serious reaction to DESCOVY. You may need urgent medical attention or hospitalisation.

itching (*pruritus*),

hives (*urticaria*)

low red blood cell count (*anaemia*)

problems with digestion resulting in discomfort after meals (*dyspepsia*)

joint pain (*arthralgia*)

→ If any of the side effects get serious tell your doctor.

Other effects that may be seen during HIV treatment

The frequency of the following side effects is not known (frequency cannot be estimated from the available data).

Bone problems. Some patients taking combination antiretroviral medicines such as DESCOVY may develop a bone disease called *osteonecrosis* (death of bone tissue caused by loss of blood supply to the bone). Taking this type of medicine for a long time, taking corticosteroids, drinking alcohol, having a very weak immune system, and being overweight, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are:

- o joint stiffness

- joint aches and pains (especially of the hip, knee and shoulder)
- difficulty with movement

→ **If you notice any of these symptoms tell your doctor.**

During HIV therapy such as with DESCOVY there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects to SAHPRA via the “6.04 Adverse Drug Reaction Reporting Form”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>.

By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store DESCOVY

Store all medicines out of reach of children.

Do not use this medicine after the expiry date which is stated on the carton, label, bottle. The expiry date refers to the last day of that month.

Store in the original package in order to protect from moisture. Keep the bottle tightly closed.

Store at or below 30 °C.

Return all unused medicine to your pharmacist Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What DESCOVY contains

The active substances are emtricitabine and tenofovir alafenamide.

Each DESCOVY film-coated tablet contains either:

200 mg of emtricitabine and tenofovir alafenamide fumarate equivalent to 10 mg of tenofovir alafenamide.

OR

200 mg of emtricitabine and tenofovir alafenamide fumarate equivalent to 25 mg of tenofovir alafenamide

The other ingredients are

Tablet core:

Microcrystalline cellulose, croscarmellose sodium, magnesium stearate.

DESCOVY 200mg/10mg Film-coated Tablets
DESCOVY 200mg/25mg Film-coated Tablets
Gilead Sciences South Africa (Pty) Ltd
3 November 2020

Film-coating:

DESCOVY 200mg/10mg: Polyvinyl alcohol, titanium dioxide, macrogol 3350, talc, iron oxide black (E172).

DESCOVY 200mg/25mg: Polyvinyl alcohol, titanium dioxide, macrogol 3350, talc, indigo carmine aluminium lake (E132).

What DESCOVY looks like and contents of the pack

DESCOVY film-coated tablets are grey, rectangular-shaped tablets, debossed on one side with “GSI” and the number “210” on the other side of the tablet.

DESCOVY film-coated tablets are blue, rectangular-shaped tablets, debossed on one side with “GSI” and the number “225” on the other side of the tablet.

DESCOVY comes in bottles of 30 tablets (with a silica gel desiccant that must be kept in the bottle to help protect your tablets). The silica gel desiccant is contained in a separate sachet or canister and should not be swallowed.

The following pack sizes are available: outer cartons containing 1 bottle of 30 film-coated tablets.

Holder of Certificate of Registration

Gilead Sciences South Africa (Pty) Ltd,
Ground Floor,
West Wing,
No.6 Kikuyu Road,
Sunninghill Extension 56,
Johannesburg, 2191

This leaflet was last revised in

3 November 2020

Registration Number

DESCOVY 200/10: 51/20.2.8/0506

DESCOVY 200/25: 51/20.2.8/0507