

## PATIENT INFORMATION LEAFLET

**SCHEDULING STATUS:** **S4**

### **CLAR HEXAL 500 (tablets)**

Clarithromycin

Contains sugar (10, 8 mg lactose monohydrate per tablet).

#### **Read all of this leaflet carefully before you start taking CLAR HEXAL 500**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor, pharmacist, nurse or other health care provider.
- CLAR HEXAL 500 has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

#### **What is in this leaflet**

1. What CLAR HEXAL 500 is and what it is used for
2. What you need to know before you take CLAR HEXAL 500
3. How to take CLAR HEXAL 500
4. Possible side effects
5. How to store CLAR HEXAL 500
6. Contents of the pack and other information

#### **1. What CLAR HEXAL 500 is and what it is used for**

CLAR HEXAL 500 belongs to a group of medicines called macrolide antibiotics. Antibiotics stop the growth of bacteria (bugs) that cause infections. CLAR HEXAL 500 contains the active

ingredient, clarithromycin. Clarithromycin is used to treat bacterial infections in many different parts of the body.

Your doctor will prescribe CLARIHEXAL 500 to treat the following infections such as:

- Lower respiratory tract such as bronchitis and pneumonia (chest infections).
- Upper respiratory tract infections such as pharyngitis (throat) and sinusitis (sinus) infections.
- Mild to moderately severe acute otitis media (ear infections, particularly inflammation of the middle ear).
- Skin and soft tissue infections such as folliculitis, cellulitis or erysipelas.
- CLARIHEXAL 500 may also be used to treat gastrointestinal ulcers caused by an organism called *Helicobacter pylori*. Together with CLARIHEXAL 500, your doctor will also prescribe other medication that you must use at the same time, to prevent the ulcer from coming back.

## **2. What you need to know before you take CLARIHEXAL 500**

### **Do not take CLARIHEXAL 500:**

- If you are hypersensitive (allergic) to clarithromycin, any macrolide antibiotics (erythromycin or azithromycin) or any of the other ingredients of CLARIHEXAL 500 (listed in section 6).
- If you are taking medicines called ergotamine or dihydroergotamine tablets or use ergotamine inhalers for migraine.
- If you are taking oral midazolam (a sedative).
- If you are taking lomitapide, lovastatin or simvastatin (HMG-CoA reductase inhibitors, commonly known as statins, used to lower levels of cholesterol (a type of fat) in the blood).
- If you are taking medicines called terfenadine or astemizole (widely taken for hay fever or allergies), domperidone or cisapride (for stomach problems) or pimozone (for mental health problems), as combining these medicines can sometimes cause serious disturbances in heart rhythm.
- Have a history of QT prolongation (an abnormality of the electrocardiogram (ECG)) or an irregular heartbeat (ventricular cardiac dysrhythmia).

V5.0 (05/08/2021)

- If you are taking medicines called ticagrelor or ranolazine (for heart attack, chest pain or angina).
- If you are taking colchicine (usually taken for gout).
- If you have hypokalaemia (a condition characterized by low levels of potassium in the blood).
- If you have severe liver disease with kidney disease.
- If you suffer from porphyria.

### **Warnings and precautions**

#### **Take special care with CLARIHEXAL 500:**

- If you have kidney and severe liver disease. You may have an increased chance of side effects.
- If you experience rhabdomyolysis, which is a condition in which skeletal muscle tissue dies, releasing substances into the blood that cause kidney failure.
- If you develop severe or prolonged diarrhoea (pseudomembranous colitis), which may have blood or mucus in it, during or after taking CLARIHEXAL 500, contact your doctor immediately.
- If you have, or have had, heart problems (e.g. heart disease, heart failure, an unusually slow heart rate) or abnormally low levels of magnesium in the blood (hypomagnesaemia).
- If you have pneumonia (serious lung infection with fever, chills, shortness of breath, cough, phlegm or coughing blood).
- In the event of severe acute hypersensitivity reactions, such as anaphylaxis, Stevens-Johnson Syndrome, and toxic epidermal necrolysis, CLARIHEXAL 500 therapy should be discontinued immediately and appropriate treatment should be urgently initiated.
- If you suffer from blood clots or currently using blood thinning medicine.
- If you have a weak immune system (immunocompromised) you may experience nausea, vomiting, irregular taste, stomach pain, runny stomach, rash, flatulence (gas), headache, hearing disturbance, irregular liver test results and irregular low white blood cell and platelet counts or difficulty breathing, restlessness and dry mouth.

### **Other medicines and CLARIHEXAL 500**

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor, nurse or pharmacist if you are taking any of the following medicines:

- Terfenadine or astemizole (used to treat hay fever or allergies) cisapride or domperidone (used to treat stomach disorders) or pimozide (used to treat certain mental disorders) as combining these medicines can sometimes cause serious disturbances in heart rhythm.
- Ergot alkaloids, e.g. ergotamine or dihydroergotamine used to treat migraines.
- Alprazolam, midazolam, temazepam, nitrazepam, lorazepam or triazolam used to treat anxiety disorders or insomnia (difficulty sleeping).
- Atorvastatin, rosuvastatin, lovastatin or simvastatin commonly known as statins, and used to lower levels of cholesterol (a type of fat) in the blood). Statins can cause rhabdomyolysis (a condition which causes the breakdown of muscle tissue which can result in kidney damage) and signs of myopathy (muscle pain or muscle weakness) should be monitored.
- Carbamazepine, valproate, phenobarbital or phenytoin used to treat for epilepsy (fits).
- Rifabutin, rifampicin, rifapentine, fluconazole or itraconazole used to treat bacterial infections.
- Etravirine, efavirenz, nevirapine, ritonavir, zidovudine, atazanavir or saquinavir used to treat HIV.
- Warfarin, rivaroxaban or apixaban also known as blood thinners used to treat blood clots.
- Quetiapine used to treat certain mental/mood disorders.
- Sildenafil, tadalafil or vardenafil used to treat erectile dysfunction.
- Tacrolimus used to treat eczema.
- Sirolimus used to prevent rejection of kidney transplant.
- Vinblastine used to treat certain types of cancer.
- Theophylline used to treat asthma.

V5.0 (05/08/2021)

- Nateglinide or repaglinide used to treat Type 2 diabetes.
- Omeprazole used to treat heart burn or stomach ulcers.
- Tolterodine used to treat overactive bladder.
- Colchicine used to treatment of gout.
- Digoxin, quinidine, disopyramide, calcium channel blockers (verapamil, amlodipine, diltiazem) used to treat irregular heartbeat.
- Itraconazole used to treat fungal infections.
- Contraceptives used to prevent pregnancy.

### **Pregnancy, breastfeeding and fertility**

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking CLARIHEXAL 500.

### **Driving and using machines**

It is not always possible to predict to what extent CLARIHEXAL 500 may interfere with the daily activities of a patient. Patients should ensure that they do not engage in the above activities until they are aware of the measure to which CLARIHEXAL 500 affects them. CLARIHEXAL 500 may make you dizzy, drowsy, confused or disorientated.

### **CLARIHEXAL 500 contains sugar**

CLARIHEXAL 500 contains lactose, which is a type of sugar. Patients with the rare hereditary condition of galactose intolerance e.g. galactosaemia, Lapp lactase deficiency, glucose-galactose malabsorption or fructose intolerance, should not take CLARIHEXAL 500.

CLARIHEXAL 500 contains lactose monohydrate which may have an effect on the control of your blood sugar if you have diabetes mellitus.

### **3. How to take CLARIHEXAL 500**

Do not share medicines prescribed for you with any other person.

V5.0 (05/08/2021)

Always take CLAR HEXAL 500 exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

The usual duration of treatment is 5 to 10 days and will depend on the type of infection and its seriousness.

Children who weigh less than 8 kg will usually be given a dose of approximately 7,5 mg/kg twice a day.

The usual dose in adults is 250 mg twice a day, which could be increased to 500 mg twice a day for more severe infections.

If you suffer from severe renal impairment, your doctor will reduce the dosage of CLAR HEXAL 500 by half.

#### **If you take more CLAR HEXAL 500 than you should**

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

#### **If you forget to take CLAR HEXAL 500**

Do not take a double dose to make up for forgotten individual doses

#### **4. Possible side effects**

CLAR HEXAL 500 can have side effects.

Not all side effects reported for CLAR HEXAL 500 are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking CLAR HEXAL 500, please consult your health care provider for advice.

**If any of the following happens, stop taking CLAR HEXAL 500 and tell your doctor immediately or go to the casualty department at your nearest hospital:**

- Swelling of the hands, feet, ankles, face, lips and mouth or throat, which may cause difficulty in swallowing or breathing.

V5.0 (05/08/2021)

- Severe skin reactions such as painful blistering of the skin, mouth, lips, eyes and genitals (symptoms of a rare allergic reaction called Stevens-Johnson syndrome/toxic epidermal necrolysis).
- A red, scaly rash with bumps under the skin and blisters (symptoms of exanthematous pustulosis).
- A rare allergic skin reactions which cause severe illness with ulceration of the mouth, lips and skin which causes severe illness with rash, fever and inflammation of internal organs (DRESS).

These are all very serious side effects. If you have them, you may have had a serious reaction to CLARIHEXAL 500. You may need urgent medical attention or hospitalisation.

**Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:**

- Changes in the way your heart beats for example if you notice it beating faster.
- Muscle pain or weakness known as rhabdomyolysis (a condition which causes the breakdown of muscle tissue which can result in kidney damage).
- Fits.
- Yellowing of the skin and eyes, dark urine, and tiredness which may be symptoms of liver problems.
- Watery and severe diarrhoea, which may be bloody.

These are all serious side effects. You may need urgent medical attention.

**Tell your doctor if you notice any of the following:**

***Frequent side effects:***

- Difficulty sleeping
- Headache or irregular taste
- Nausea
- Vomiting

V5.0 (05/08/2021)

- Stomach pain
- Diarrhoea
- Indigestion
- Irregular liver function test
- Rash
- Excessive sweating

***Less frequent side effects:***

- Thrush
- Mild to severe nausea, vomiting, cramps and diarrhoea, usually caused by a virus
- Infection
- Irregular blood cells that may cause fever, severe chills, sore throat or mouth ulcers, bleeding or bruising more easily than normal
- Weight loss
- Lack of appetite
- Anxiety
- Nervousness
- Dizziness
- Sleepiness
- Shaking
- Ringing in the ear or hearing loss
- Vertigo
- Nosebleed
- Inflammation of the lining of the oesophagus and lining of the stomach
- Anal pain
- Bloating, constipation, wind, burping
- Dry mouth

V5.0 (05/08/2021)

- Situation where the bile (fluid made by the liver and stored in the gallbladder) cannot flow from the gallbladder to the duodenum (cholestasis)
- Itching or itchy rash
- Muscle spasms or muscle pain
- Aching muscles, muscle tenderness or weakness, not caused by exercise

***Frequency unknown:***

- Bacterial infection of the skin
- Confusion
- Depression
- Seeing things (hallucinations)
- Abnormal dreams or nightmares
- Mood of excitement, over activity and uninhibited behaviour
- Loss of taste or smell
- Tingling feeling in hands and feet also known as “pins and needles”
- Deafness
- Bleeding
- Teeth or mouth discolouration
- Acne
- Generally feeling unwell
- Fever
- Chest pain
- Chills
- Fatigue

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

**Reporting of side effects**

V5.0 (05/08/2021)

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of CLARIHEXAL 500.

Suspected side effects can also be reported directly to the HCR via [Patientsafety.sacg@novartis.com](mailto:Patientsafety.sacg@novartis.com).

## **5. How to store CLARIHEXAL 500**

- Store at or below 25 °C.
- **STORE ALL MEDICINES OUT OF REACH OF CHILDREN.**

## **6. Contents of the pack and other information**

### **What CLARIHEXAL 500 contains**

Active ingredient: Clarithromycin.

Inactive ingredients: Cellulose microcrystalline, cellulose powdered, croscarmellose sodium, magnesium stearate, HPMC-based white coating premix [consists of 36 % lactose monohydrate (EP), 26 % titanium dioxide (EP), 28 % hydroxypropyl methylcellulose (EP) and 10 % polyethylene glycol (EP)], silica colloidal anhydrous.

### **What CLARIHEXAL 500 looks like and contents of the pack**

Tablets are available in Alu/Alu blister packs packed into the carton boxes.

1 blister strip of 10 tablets packed into a carton i.e. 10 tablets per carton.

Not all pack sizes may be marketed.

### **Holder of Certificate of Registration**

Sandoz SA (Pty) Ltd<sup>1</sup>

Waterfall 5-Ir

Magwa Crescent West

Waterfall City

Jukskei View

V5.0 (05/08/2021)

2090

**This leaflet was last revised in**

05 August 2021

**Registration number**

38/20.1.1/0006

**Access to the corresponding Professional Information**

Not applicable.

<sup>1</sup>Company Reg. No.: 1990/001979/07