

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: **S4**

DIFLUCAN® CAPSULES 50 mg

DIFLUCAN® CAPSULES 150 mg

DIFLUCAN® CAPSULES 200 mg

Fluconazole

Contains sugar

Each DIFLUCAN 50 mg capsule contains 49,708 mg lactose monohydrate.

Each DIFLUCAN 150 mg capsule contains 149,123 mg lactose monohydrate.

Each DIFLUCAN 200 mg capsule contains 198,830 mg lactose monohydrate.

Read all of this leaflet carefully before you start taking DIFLUCAN

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other health care provider.
- DIFLUCAN has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What DIFLUCAN is and what it is used for
2. What you need to know before you take DIFLUCAN
3. How to take DIFLUCAN
4. Possible side effects
5. How to store DIFLUCAN
6. Contents of the pack and other information

1. What DIFLUCAN is and what it is used for

DIFLUCAN is one of a group of medicines called “antifungals”. The active ingredient is fluconazole.

DIFLUCAN is used to treat infections caused by fungi including yeasts and may also be used to stop you from getting a fungal infection.

2. What you need to know before you take DIFLUCAN

Do not take DIFLUCAN

- if you are hypersensitive (allergic) to fluconazole, other medicines you have taken to treat a fungal infection, or to any of the other ingredients of DIFLUCAN (listed in section 6). The symptoms of an allergic reaction may include itching, reddening of the skin or difficulty in breathing.
- if you are taking astemizole or terfenadine (an antihistamine for allergies)
- if you are taking cisapride (used for stomach upsets)
- if you are taking erythromycin (for treating infections (antibiotic))
- if you are taking pimozide (for treating schizophrenia)
- if you are taking quinidine (used for irregular heartbeats)
- if you are pregnant or breastfeeding

Warnings and precautions

Take special care with DIFLUCAN:

- if you have liver or kidney problems
- if you suffer from heart disease, including heart rhythm problems
- if laboratory investigations have shown abnormal levels of potassium, calcium or magnesium in your blood
- if you develop severe skin reactions (itching, reddening of the skin or difficulty in breathing)
- if you develop signs of ‘adrenal insufficiency’ where the adrenal glands do not produce adequate amounts of certain steroid hormones such as cortisol (chronic, or long-lasting fatigue, muscle weakness, loss of appetite,

weight loss, abdominal pain)

- if you have ever developed a severe skin rash or skin peeling, blistering and/or mouth sores after taking DIFLUCAN

Serious skin reactions including drug reaction with eosinophilia and systemic symptoms (DRESS) have been reported in association with DIFLUCAN treatment. Stop taking DIFLUCAN and seek medical attention immediately if you notice any of the symptoms related to these serious skin reactions described in section 4.

Talk to your health care provider if the fungal infection does not improve, as alternative antifungal therapy may be needed.

Other medicines and DIFLUCAN

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines).

Tell your doctor immediately if you are taking astemizole or terfenadine (an antihistamine for treating allergies) or cisapride (used for stomach upsets) as these should not be taken with DIFLUCAN.

There are some medicines that may interact with DIFLUCAN. Make sure your doctor knows if you are taking any of the following medicines as a dose adjustment or monitoring may be required to check that the medicines are still having the desired effect:

- halofantrine (to treat malaria)
- amiodarone (used for treating uneven heartbeats 'arrhythmias')
- water tablets, such as hydrochlorothiazide, used to treat fluid retention and high blood pressure
- medicines for treating infections (antibiotics) such as azithromycin, rifampicin or rifabutin
- alfentanil, fentanyl or methadone (for treating pain)
- amitriptyline, nortriptyline (used to treat depression)

- amphotericin B, voriconazole (used to treat serious fungal infections)
- warfarin (or similar medicines) that thin the blood to prevent blood clots
- benzodiazepines such as midazolam, triazolam used to help you sleep or for anxiety
- carbamazepine or phenytoin (used to control epilepsy)
- calcium channel blockers or losartan for lowering blood pressure
- celecoxib (used to treat some types of arthritis)
- ciclosporin, sirolimus or tacrolimus (to prevent transplant rejection)
- cyclophosphamide, prednisone, vinca alkaloids or ibrutinib (for treating some forms of blood cancer e.g. leukaemia)
- atorvastatin, simvastatin or fluvastatin (for lowering cholesterol)
- ivacaftor, alone or in combination with other medicines (used for treating cystic fibrosis)
- lurasidone (used to treat schizophrenia)
- non-steroidal anti-inflammatory drugs (such as aspirin and ibuprofen) that are used to treat pain and fever
- olaparib (used for treating ovarian cancer)
- oral contraceptives (the pill) (to prevent pregnancy)
- saquinavir or zidovudine, also known as AZT (used in HIV-infected patients)
- medicines for diabetes such as chlorpropamide, glibenclamide, glipizide or tolbutamide
- theophylline (used to control asthma)
- tofacitinib (used for treating rheumatoid arthritis)
- tolvaptan (used to treat low levels of sodium in blood or to slow kidney function decline)
- vitamin A (as a trans-retinoid acid used to treat acne)

Pregnancy and breastfeeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking this medicine.

You should not take DIFLUCAN while you are pregnant or if you are breastfeeding.

Effective contraceptive measures must be considered in women of childbearing potential throughout treatment with DIFLUCAN and for approximately 1 week after the final dose.

Driving and using machines

Treatment with DIFLUCAN may lead to dizziness or seizures to occur and could affect your ability to drive or use machinery.

It is not always possible to predict to what extent DIFLUCAN may interfere with the daily activities of a patient. Patients should ensure that they do not engage in the above activities until they are aware of the measure to which DIFLUCAN affects them.

DIFLUCAN capsules contain lactose (sugar)

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take DIFLUCAN

Do not share medicines prescribed for you with any other person.

Always take DIFLUCAN exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

You can take DIFLUCAN capsules with or without a meal at any time of the day.

Elderly

The usual adult dose should be given unless you have kidney problems.

Patients with kidney problems

Your doctor may modify your dose, depending on your kidney function.

If you are not sure, ask your doctor or pharmacist.

It is best to take your medicine at the same time each day. You may take it with or without meals.

Your doctor will tell you how long your treatment with DIFLUCAN will last. If you have the impression that the effect of DIFLUCAN is too strong or too weak, tell your doctor or pharmacist.

If you take more DIFLUCAN than you should

Taking too much DIFLUCAN may make you unwell. In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to take DIFLUCAN

Do not take a double dose to make up for forgotten individual doses. If you forget to take a dose, take it as soon as you remember. If it is almost time for your next dose, do not take the dose that you missed.

4. Possible side effects

DIFLUCAN can have side effects.

Not all side effects reported for DIFLUCAN are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking DIFLUCAN, please consult your health care provider for advice.

You may develop allergic reactions. If any of the following happens, tell your doctor immediately.

- Sudden wheezing, difficulty in breathing or tightness in chest

- Swelling of eyelids, face or lips
- Itching all over the body, reddening of the skin or itchy red spots
- Skin rash
- Severe skin reactions such as a rash that causes blistering (this can affect the mouth and tongue)
- Hypersensitivity reaction with skin rash, fever, swollen glands, increase in a type of white blood cell (eosinophilia) and inflammation of internal organs (liver, lungs, heart, kidneys and large intestine); stop taking DIFLUCAN and seek medical attention immediately if you notice any of the following symptoms: widespread rash, high body temperature and enlarged lymph nodes (swollen glands) (DRESS syndrome or drug hypersensitivity syndrome)

These are all serious side effects. You may need urgent medical attention.

If you are an AIDS patient, you are more likely to get severe skin reactions to medicines including DIFLUCAN.

Tell your doctor if you notice any of the following:

Frequent side effects

- headache
- stomach discomfort
- diarrhoea
- feeling sick (nausea)
- vomiting
- increases in blood tests of liver function
- rash

Less frequent side effects

- insomnia (difficulty in sleeping)
- somnolence (feeling sleepy)

- seizures (fits)
- dizziness
- tingling or pricking (pins and needles)
- altered sense of taste
- sensation of spinning (vertigo)
- indigestion
- wind
- dry mouth
- decrease in bile flow
- yellowing of the skin and eyes (jaundice)
- higher than normal levels of bilirubin in your blood (seen in blood tests)
- itching
- increased sweating
- myalgia (muscle pain)
- fatigue
- tiredness
- general feeling of being unwell
- fever
- lower than normal white blood cells that help defend against infections and blood cells that help to stop bleeding
- high blood levels of cholesterol, fats or salt
- low blood potassium levels
- tremor
- change in heart rate or rhythm
- liver damage or failure
- hair loss

If any of the side effects gets serious, or if you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reactions Reporting Form**”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of DIFLUCAN.

5. How to store DIFLUCAN

- Store all medicines out of reach of children.
- Store at or below 30 °C in a dry place.
- Do not use DIFLUCAN after the expiry date which is stated on the pack. The expiry date refers to the last day of the month.
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What DIFLUCAN contains

- The active substance is fluconazole.
- Each DIFLUCAN Capsule 50 mg contains 50 mg fluconazole.
- Each DIFLUCAN Capsule 150 mg contains 150 mg fluconazole.
- Each DIFLUCAN Capsule 200 mg contains 200 mg fluconazole.
- The other ingredients are lactose, maize starch, colloidal silicone dioxide, magnesium stearate and sodium lauryl sulphate in a hard gelatin capsule containing titanium dioxide with or without patent blue as colourants erythrosine (200 mg only) and indigotine (200 mg only). The printing ink contains black iron oxide, shellac, industrial methylated spirit, N-Butyl alcohol, soya lecithin and antifoam DC 1510.

What DIFLUCAN looks like and contents of the pack

DIFLUCAN Capsules 50 mg have a white body and a turquoise blue cap, containing a white powder. They have FLU-50 and the Pfizer logo printed on them.

DIFLUCAN Capsules 150 mg are turquoise blue (cap and body), containing a white powder. They have FLU-150 and the Pfizer logo printed on them.

DIFLUCAN Capsules 200 mg have a white body and a purple cap, containing a white powder. They have FLU-200 and the Pfizer logo printed on them.

DIFLUCAN Capsules 50 mg come in blister packs of 14 capsules.

DIFLUCAN Capsules 150 mg come in blister packs of 1 or 4 capsules.

DIFLUCAN Capsules 200 mg come in blister packs of 28 and 30 capsules.

Not all pack sizes may be marketed.

Holder of Certificate of Registration

Pfizer Laboratories (Pty) Ltd

85 Bute Lane

Sandton 2196

South Africa

Tel: +27(0)11 320 6000 / 0860 734 937 (Toll-free South Africa)

This leaflet was last revised in

01 September 2022

Registration number

DIFLUCAN Capsules 50 mg: V/20.2.2/339

DIFLUCAN Capsules 150 mg: V/20.2.2/340

DIFLUCAN Capsules 200 mg: 28/20.2.2/151