

SCHEDULING STATUS: S3**Novabrex 100**, hard capsule**Novabrex 200**, hard capsule

Celecoxib

Contains sugar (Novabrex 100 contains 21,4 mg and Novabrex 200 contains 42,8 mg lactose monohydrate per capsule)**Read all of this leaflet carefully before you start taking Novabrex**

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other healthcare provider.
- Novabrex has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What Novabrex is and what it is used for
2. What you need to know before you take Novabrex
3. How to take Novabrex
4. Possible side effects
5. How to store Novabrex
6. Contents of the pack and other information

1. What Novabrex is and what it is used for

Novabrex belongs to a group of medicines called nonsteroidal anti-inflammatory drugs (NSAID), and specifically a sub-group known as cyclooxygenase-2 (COX-2) inhibitors.

Your body makes substances called prostaglandins. Some prostaglandins may cause pain and inflammation. In conditions such as rheumatoid arthritis and osteoarthritis your body makes more of these. Novabrex acts by reducing the production of prostaglandins, thereby reducing the pain and inflammation.

Novabrex is used in adults for the relief of signs and symptoms of rheumatoid arthritis, osteoarthritis and ankylosing spondylitis (progressive stiffening of the spine).

Novabrex is also used in the treatment of pain following an operation, including dental surgery, musculoskeletal pain, primary dysmenorrhoea (painful menstrual periods).

2. What you need to know before you take Novabrex

Do not take Novabrex:

- if you are hypersensitive (allergic) to celecoxib or any of the ingredients of Novabrex (see section 6 “What this product contains”);
- if you had an allergic reaction to a group of medicines called “sulphonamides” (medicines that prevent the growth of bacteria);
- if you have severe liver disease;
- if you have severe kidney disease;
- if you have had allergic reactions such as asthma, nose polyps, severe nasal congestion, or an itchy skin rash, swelling of the face, lips, tongue or throat, breathing difficulties or wheezing from aspirin or other NSAIDs (anti-inflammatory and pain-relieving medicines, such as diclofenac or ibuprofen);
- Established ischaemic heart disease, or cerebrovascular disease, e.g. you have been diagnosed with a heart attack, stroke, or blockages of blood vessels to the heart or brain;
- if you are about to have coronary artery bypass graft surgery;
- if you currently have an ulcer in your stomach or intestines, or if you are bleeding in your stomach or intestines (you may notice that you have black sticky stools or bloody diarrhoea);
- if you have an inflammatory disease of the intestines such as ulcerative colitis or Crohn’s disease;
- if you are pregnant or it is possible that you could become pregnant;
- if you are breastfeeding.

Warnings and precautions

Talk to your doctor or pharmacist before taking Novabrex

Take special care with Novabrex:

- if you smoke, have diabetes or high blood cholesterol;
- if you have had a serious allergic reaction or a serious skin reaction to any medicines;
- if you have previously had an ulcer or bleeding in your stomach or intestines. (**Do not take Novabrex** if you currently have an ulcer or bleeding in your stomach or intestine);
- if you are taking corticosteroid medicines (for example prednisone);
- if you are taking aspirin (even at a low dose for heart protective purposes);
- if you use medicines to reduce blood clotting (e.g. warfarin/warfarin like anticoagulants or novel oral anti-clotting medicines, e.g. apixaban);
- if you are taking Novabrex at the same time as other non-aspirin NSAIDs such as ibuprofen or diclofenac. The use of these medicines together should be avoided;
- if you have fluid retention (such as swollen ankles and feet);
- if you have heart failure or other heart problems;
- if you have high blood pressure;
- if your liver or kidneys are not working well, as your doctor may want to keep a regular check on you;
- if you are dehydrated, for instance due to sickness, diarrhoea or the use of diuretics (medicines used to get rid of excess fluid in the body);
- if you feel ill due to an infection, or think you have an infection, as Novabrex may mask a fever or other signs of infection and inflammation;
- if you are above 65 years of age your doctor will want to monitor you regularly;
- as the consumption of alcohol and NSAIDs may increase the risk of gastrointestinal problems;
- Novabrex may lead to an increase in blood pressure. Your doctor may want to monitor your blood pressure on a regular basis;
- some cases of severe liver reactions, including severe liver inflammation, liver damage, liver failure (some with fatal outcomes or requiring liver transplant), have been reported with celecoxib. Of the cases that reported time to onset, most severe liver reactions occurred within one month of start of treatment.

Other medicines and Novabrex

Always tell your healthcare provider if you are taking any other medicine. (This includes complementary or traditional medicines.)

Before you take Novabrex, make sure your doctor knows if you are taking the following medicines:

- Warfarin (used to prevent blood from clotting);
- ACE inhibitors, angiotensin II receptor antagonists, beta blockers (used for high blood pressure and heart failure);
- Diuretics (used to treat fluid retention);
- Ciclosporin and tacrolimus (used for immune system suppression e.g. after transplants);
- Aspirin (used for pain and fever). Novabrex can be taken with low dose aspirin (75 mg or less daily).

Ask your doctor for advice before taking both medicines together.

Novabrex has no antiplatelet effect and is therefore not a substitute for aspirin for prevention of thrombosis;

- Neuroleptics (used to treat some mental disorders);
- Medicines for anxiety (diazepam) or to treat depression (citalopram, imipramine);
- Medicines to treat an irregular heartbeat (amiodarone, quinidine);
- Dextromethorphan (used to treat coughs);
- Methotrexate (used to treat rheumatoid arthritis, psoriasis and leukaemia);
- Lithium (used to treat some types of depression);
- Oral contraceptives;
- Glibenclamide, tolbutamide (medicines used in the management of type 2 diabetes);
- Phenytoin, a medicine used in epilepsy (seizure disorder);
- Fluconazole (used to treat fungal infection);
- Rifampicin (used to treat bacterial infections);
- Carbamazepine (used to treat epilepsy/seizures and some forms of pain or depression);
- Barbiturates (used to treat epilepsy/seizures and some sleep disorders);

- Ketoconazole (an antifungal medicine);
- Antacids (used for heartburn).

Pregnancy, breastfeeding and fertility

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking Novabrex

Pregnancy

Novabrex must not be used by women who are pregnant or can become pregnant (i.e. women of childbearing potential who are not using adequate contraception) during ongoing treatment. If you become pregnant during treatment with Novabrex you should discontinue the treatment and contact your doctor for alternative treatment.

Breastfeeding

Novabrex must not be used during breastfeeding.

Fertility

NSAIDs, including Novabrex, may make it more difficult to become pregnant. You should tell your doctor if you are planning to become pregnant or if you have problems becoming pregnant.

Driving and using machines

You should be aware of how you react to Novabrex before you drive or operate machinery. If you feel dizzy or drowsy after taking Novabrex, do not drive or operate machinery until these effects wear off.

Novabrex contains lactose

Novabrex contains lactose (a type of sugar). If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking Novabrex.

3. How to take Novabrex

Do not share medicines prescribed for you with any other person.

Always take Novabrex exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

If you think or feel that the effect of Novabrex is too strong or too weak, talk to your doctor or pharmacist.

Use the lowest effective dose for the shortest possible time.

Your doctor will tell you what dose you should take.

The recommended dose for osteoarthritis is 200 mg taken as a single dose, or as two divided doses.

The recommended dose for rheumatoid arthritis is 100 mg or 200 mg twice daily.

The recommended dose for pain following dental surgery is 100 mg or 200 mg twice daily.

The recommended dose for pain following an operation is 200 mg once daily.

The recommended dose for musculoskeletal pain is 200 mg twice daily.

The recommended dose for primary dysmenorrhoea is 400 mg initially, followed by an additional 200 mg on the first day if needed. On subsequent days, the recommended dose is 200 mg twice daily.

The recommended dose for ankylosing spondylitis is 200 mg daily, taken as a single dose or as 100 mg twice daily.

If you have liver problems, or if you weigh less than 50 kg, your doctor will recommend the lowest recommended dose.

Children:

No information is available on the use of Novabrex in children. It is therefore not recommended for use in children.

Method of administration:

Novabrex is for oral use. Swallow the capsules whole with a glass of water at any time of the day, with or without food. However, try to take each dose of Novabrex at the same time each day. Do not chew or crush the granules.

Contact your doctor within two weeks of starting treatment if you do not experience any benefit.

If you take more Novabrex than you should

You should not take more capsules than your doctor tells you to.

In the event of overdose, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to take Novabrex

If you forget to take a capsule, take it as soon as you remember.

Do not take a double dose to make up for forgotten individual doses.

If you stop taking Novabrex

Suddenly stopping your treatment with Novabrex may lead to your symptoms getting worse. Do not stop taking Novabrex unless your doctor tells you to. Your doctor may tell you to reduce the dose over a few days before stopping completely.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Novabrex can have side effects.

Not all side effects reported for Novabrex are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking Novabrex, please consult your doctor, pharmacist or other healthcare provider for advice,

If any of the following happens, stop taking Novabrex and tell your doctor immediately or go to the casualty department at your nearest hospital:

- swelling of the hands, feet, ankles, face, lips and mouth or throat, which may cause difficulty in swallowing or breathing.
- a serious skin reaction with rash, blistering or peeling of the skin.

These are all very serious side effects. If you have them, you may have had a serious reaction to Novabrex. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

- weakness or clumsiness of a hand, arm or leg, difficulties with speech and swallowing, vision problems and headache. These may be signs of a stroke or bleeding in the brain, which may be fatal;
- shortness of breath or difficulty breathing, sudden chest pain, cramping or tingling sensation up the left arm (heart attack);
- problems breathing, shortness of breath, chest pain. You may have a blood clot in the lung (pulmonary embolism) and will need urgent medical attention;
- throbbing or cramping pain in a leg, swelling in a leg, warm skin around painful area (deep vein thrombosis);
- severe stomach pain or any sign of bleeding in the stomach or intestines, such as passing black or bloodstained stools, or vomiting blood;
- liver failure [symptoms may include nausea (feeling sick), diarrhoea, jaundice (your skin or the whites of your eyes look yellow)];
- bladder or kidney infections. Some signs may be fever, chills, blood in the urine or frequent urination;
- acute renal failure/nephrotic syndrome (decreased urine output, swelling of legs, fatigue, foamy urine, loss of appetite);
- painful blisters, sensitive to touch (shingles);
- red, swollen, painful skin with fever (a bacterial skin infection, known as erysipelas);
- ache or a burning pain in your abdomen (Helicobacter infection).

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Frequent side effects:

- infections of the airways such as sinusitis (swelling and redness of the nasal sinuses) or bronchitis (inflammation of the mucous membranes of the airways);
- worsening of existing allergies;
- weight gain;

- trouble sleeping;
- dizziness, abnormal muscle tension;
- headache;
- high blood pressure, worsening of existing high blood pressure. You may not know that you have high blood pressure unless you have your blood pressure tested;
- runny nose, sore throat, shortness of breath;
- nausea (feeling sick), vomiting (being sick);
- indigestion, windiness, pain in the stomach;
- problem swallowing;
- sore teeth;
- skin rash, itching skin;
- accidental injury;
- flu-like illness;
- keeping fluid back in the body tissues, with swollen ankles, legs and/or hands.

Less frequent side effects:

- inflammation or infection of the inner ear (dizziness or feeling that everything is spinning around you);
- infection of the gums;
- blood disorders, such as anaemia (too few red blood cells or their haemoglobin content is too low), leukopenia (a lowered white blood cell count) or thrombocytopenia (deficiency of platelets in the blood)*
*You will only know that you have these disorders if your doctor orders blood tests. See your doctor if you are pale and tired, get more infections than usual or have problems to stop bleeding, or if you have unexpected bruising;
- increased sodium in the blood, increased potassium in the blood (determined by blood tests);
- feeling tired, confused;
- feeling anxious or depressed;
- seeing or hearing things that do not exist (hallucinations);

- blurred vision, eye inflammation and bloodshot eyes, vitreous floaters (seeing spots or strings that move when you move your eyes), bleeding in an eye, painful eyes;
- ringing in the ears, above normal hearing sense, voice disorder;
- irregular heart rhythm, increased heart rate;
- flushing, inflammation of a blood vessel (you may feel unwell with fever, sweats and tiredness);
- tightness in the chest with coughing and wheezing (bronchospasm);
- shortness of breath, cough, fatigue (inflammation of the lung tissues - pneumonitis);
- abdominal pain, cramping, bloating, constipation;
- mouth ulcers, numbness, a burning sensation in the mouth (stomatitis);
- hair loss;
- muscle spasms (leg cramps), muscle pain;
- lower limb fractures;
- kidney function disorders (shown by laboratory tests);
- frequent urination during the night (nocturia);
- menstrual disorders;
- swollen face;
- chest pain.

Frequency unknown

- sleepiness;
- abnormal skin sensations (tingling or tickling or itching or burning);
- problems with coordination and speech (ataxia);
- no sense of smell or taste, or taste affected;
- worsening of existing epilepsy (seizure disorder);
- female fertility decreased;
- vaginal bleeding;
- tender breasts.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of Novabrex.

5. How to store Novabrex

Store all medicine out of reach of children.

Store at or below 25 °C.

Store in the blisters until required for use.

Do not use the capsules after the expiry date shown on the carton and blister.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What Novabrex contains

The active substance is celecoxib.

1 capsule contains 100 mg or 200 mg celecoxib

The other ingredients are:

Capsule contents are: Crospovidone, hydroxypropyl cellulose, lactose monohydrate, povidone, sodium stearyl fumarate and sodium lauryl sulfate.

Capsule shell: gelatin, titanium dioxide and water.

Printing ink: butyl alcohol, dehydrated alcohol, FD & C Blue # 2 Aluminium Lake (Indigo Carmine) -Novabrex 100, and yellow iron oxide (Novabrex 200), isopropyl alcohol, propylene glycol, shellac and strong ammonia solution.

What Novabrex looks like and contents of the pack

Novabrex 100: White cap/ White body, Size '4' hard gelatin capsule filled with white to off white granular powder, imprinted with 'Y' on cap and '100' on body with blue ink.

Novabrex 200: White cap/ White body, Size '2' hard gelatin capsule filled with white to off white granular powder, imprinted with 'Y' on cap and '200' on body with gold ink.

Novabrex 100 is packaged in clear transparent PVC film/Aluminium foil blister packs.

Pack size: 60's - Each blister contains 10 capsules with 6 blisters packed per outer carton.

Novabrex 200 is packaged in clear transparent PVC/ -Aluminium foil blister packs.

Pack sizes: 10's and 30's - Each blister contains 10 capsules with 1 or 3 blisters packed per outer carton.

Capsules packed in the above blisters will be further packed in pre-printed cartons with package leaflet according to the approved pack size.

Holder of Certificate of Registration

Novagen Pharma (Pty) Ltd

Office 2, 100 Sovereign Drive

Route 21 Corporate Park

Nellmapius Drive, Irene

0157 – Pretoria

Tel: +27 12 345 3175

This leaflet was last revised in

16 November 2021

REGISTRATION NUMBER

Novabrex 100: 50/3.1/0744

Novabrex 200: 50/1.3/0745