

**Approved Professional information for
Adrenaline (Epinephrine) 1 mg/1 ml (1:1 000) Fresenius**

SCHEDULING STATUS

S4

1. NAME OF THE MEDICINE

ADRENALINE (EPINEPHRINE) 1 mg/1 ml (1: 1 000) FRESENIUS solution for injection

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Active ingredient:

Each 1 ml ampoule contains 1 mg epinephrine (adrenaline) as tartrate.

Excipients with known effect:

Antioxidant: sodium metabisulphite 0,1 % *m/v*.

Sugar free.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for injection.

Clear, colourless to slight straw-coloured solution in amber or clear ampoules.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS may be used in the treatment of acute allergy and anaphylactic shock.

4.2 Posology and method of administration

Discard contents if discoloured.

By subcutaneous or intramuscular injection.

Adults: 0,2 to 0,5 ml

Children: 0,1 to 0,3 ml

Adults:

Bronchodilator: Subcutaneous injection: 0,2 to 0,5 mg every 20 minutes to 4 hours if necessary (maximum of 1 mg per dose).

Anaphylactic reaction: Intramuscular or subcutaneous injection: 0,2 to 0,5 mg repeated every 10 to 15 min if necessary (maximum 1 mg per dose).

Children:

Bronchodilator or anaphylactic reactions: Subcutaneous 0,01 mg per kg body weight (maximum of 0,5 mg per dose) every 15 min for 2 doses and then every 4 hours as needed.

The patient must be continuously monitored.

Adults:

Vasopressor (anaphylactic shock):

0,1 mg to 0,25 mg (base) administered slowly. May be repeated every five to fifteen minutes as needed.

Children:

Vasopressor (anaphylactic shock):

0,01 mg (base) per kg of body weight every five to fifteen minutes as needed, if an inadequate response to IM or SC administration.

4.3 Contraindications

- Hypersensitivity to sympathomimetics, including epinephrine (adrenaline), or to any of the other ingredients of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS (see section 6.1).
- ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS should not be used in fingers, toes, ears, nose or genitalia owing to the risk of ischaemic tissue necrosis.
- Adrenaline is frequently used in emergency situations and any contraindications are therefore relative.
- ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS interacts with monoamine oxidase inhibitors and should not be given to patients receiving such treatment or within 14 days of its termination.

4.4 Special warnings and precautions for use

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS should be used with caution in patients with:

- Parkinson's disease, hyperthyroidism, psychoneurosis, phaeochromocytoma (diagnosed or suspected), narrow angle glaucoma (or predisposition to), diabetes mellitus, hypokalaemia or hypercalcaemia;
- severe renal impairment, prostatic hypertrophy or urination difficulty;
- cerebrovascular disease, organic brain damage or arteriosclerosis;
- autonomic dysreflexia (hyperreflexia), particularly in spinal cord injury (e.g., tetraplegics);
- shock (other than anaphylactic shock; cardiogenic, traumatic, or haemorrhagic);
- organic heart disease, cardiovascular disease, or cardiac dilatation (severe angina pectoris, obstructive cardiomyopathy, hypertension) as well as most patients with dysrhythmias.
Anginal pain may be induced when coronary insufficiency is present;
- congestive heart failure, coronary artery disease, degenerative heart disease, ischaemic heart disease, phenothiazine-induced circulatory collapse or hypotension.

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS should be used with caution in older patients.

Epinephrine (adrenaline) should be used with extreme caution in patients with long-standing bronchial asthma and emphysema who have developed degenerative heart disease.

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS should be avoided or used with caution in patients undergoing anaesthesia with cyclopropane, halothane, or other halogenated anaesthetics, as they may induce ventricular fibrillation (see section 4.5).

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS should not be used during the second stage of labour (see section 4.6).

Accidental intravascular (IV) injection may result in cerebral haemorrhage due to the sudden rise in blood pressure.

The intramuscular (IM) route is generally preferred in the initial treatment of anaphylaxis, the IV route is generally more appropriate in the Intensive Care Unit or Emergency Department setting. Epinephrine (adrenaline) 1 mg/ml (1:1 000) solution for injection is not suitable for IV use. If the adrenaline 0,1 mg/ml (1:10 000) injection is not available, adrenaline 1 mg/ml (1:1 000) solution must be diluted to 0,1 mg/ml (1:10 000) before IV use.

The IV route for injection of adrenaline must be used with extreme caution and is best reserved for specialists familiar with IV use of adrenaline.

Monitor the patient as soon as possible (pulse, blood pressure, electrocardiogram (ECG), pulse oximetry) in order to assess the response to epinephrine (adrenaline).

The best site for IM injection is the anterolateral aspect of the middle third of the thigh. The needle used for injection needs to be sufficiently long to ensure that the epinephrine (adrenaline) is injected into muscle. Intramuscular injections of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS into the buttocks should be avoided because of the risk of tissue necrosis.

Prolonged use of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS can result in severe metabolic acidosis (because of elevated blood concentrations of lactic acid), renal necrosis and tachyphylaxis.

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS contains sodium metabisulphite, which can rarely cause severe hypersensitivity reactions and bronchospasm.

The presence of sodium metabisulphite in parenteral ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESINIUS and the possibility of allergic-type reactions should not deter use of the medicine when indicated for the treatment of serious allergic reactions or for other emergency situations.

4.5 Interaction with other medicines and other forms of interaction

Sympathomimetic medicines:

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESINIUS should not be administered concomitantly with other sympathomimetic medicines because of the possibility of additive effects and increased toxicity.

Alpha-adrenergic medicines:

The vasoconstrictor and pressor effects of epinephrine (adrenaline), mediated by its alpha-adrenergic action, may be enhanced by concomitant administration of medicines with similar effects, such as ergot alkaloids or oxytocin.

Alpha-adrenergic blocking medicines:

Alpha-blockers, such as phentolamine, antagonise the vasoconstriction and hypertension effects of epinephrine (adrenaline). This effect may be beneficial in epinephrine (adrenaline) overdose (see section 4.9). ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESINIUS specifically reverses the antihypertensive effects of adrenergic neurone blockers, such as guanethidine with the risk of severe hypertension.

Beta-adrenergic blocking medicines:

Severe hypertension and reflex bradycardia may occur with non-cardioselective beta-blocking medicines, such as propranolol, due to alpha-mediated vasoconstriction.

Beta-blockers, especially non-cardioselective medicines, also antagonise the cardiac and bronchodilator effects of epinephrine (adrenaline). Patients with severe anaphylaxis who are taking non-cardioselective beta-blockers may not respond to ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS treatment.

General anaesthetics:

Administration of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS in patients receiving halogenated hydrocarbon general anaesthetics that increase cardiac irritability and seem to sensitise the myocardium to epinephrine (adrenaline) may result in dysrhythmias including ventricular premature contractions, tachycardia or fibrillation (see section 4.4).

Antihypertensive medicines:

Epinephrine (adrenaline) specifically reverses the antihypertensive effects of adrenergic neurone blockers, such as guanethidine, with the risk of severe hypertension. ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS increases blood pressure and may antagonise the effects of antihypertensive medicines.

Antidepressant medicines:

Tricyclic antidepressants, such as imipramine, inhibit reuptake of directly acting sympathomimetic medicines, and may potentiate the effect of adrenaline, increasing the risk of development of hypertension and cardiac dysrhythmias.

Concurrent use or use within 2 weeks of a monoamine oxidase inhibitor increases the risk of adverse events (see section 4.3).

Phenothiazines:

Phenothiazines block alpha-adrenergic receptors (see above).

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS should not be used to counteract circulatory collapse or hypotension caused by phenothiazines; a reversal of the pressor effects of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS may result in further lowering of blood pressure.

Other medicines:

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS should not be used in patients receiving high dosage of other medicines (e.g. cardiac glycosides) that can sensitise the heart to dysrhythmias. Some antihistamines (e.g. diphenhydramine) and thyroid hormones may potentiate the effects of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS, especially on heart rhythm and rate. ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS increases the risk of cardiac adverse effects of levodopa. Use of entacapone may potentiate the chronotropic and dysrhythmogenic effects of epinephrine (adrenaline).

Hypokalaemia:

The hypokalaemic effect of epinephrine (adrenaline) may be potentiated by other medicines that cause potassium loss, including corticosteroids, potassium-depleting diuretics, aminophylline and theophylline.

Hyperglycaemia:

Epinephrine (adrenaline)-induced hyperglycaemia may lead to loss of blood-sugar control in diabetic patients treated with insulin or oral hypoglycaemic medicines.

4.6 Fertility, pregnancy and lactation

Pregnancy:

Epinephrine (adrenaline) crosses the placenta. There is some evidence of a slightly increased evidence of congenital abnormalities.

Injection of epinephrine (adrenaline) may cause anoxia to the fetus, fetal tachycardia, cardiac irregularities, extrasystoles and louder heart sounds.

Epinephrine (adrenaline) usually inhibits spontaneous or oxytocin induced contractions of the pregnant human uterus and may delay the second stage of labour. In dosage sufficient to reduce uterine contractions, the medicine may cause a prolonged period of uterine atony with haemorrhage. For this reason parenteral epinephrine (adrenaline) should not be used during the second stage of labour (see section 4.4).

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS should only be used during pregnancy if the potential benefits justify the possible risks to the fetus.

Lactation:

Epinephrine (adrenaline) is distributed into breast milk. Breastfeeding should be avoided in mothers receiving ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS.

4.7 Effects on ability to drive and use machines

The ability of a patient to drive and use machines may be affected by the anaphylactic reaction, as well as by possible adverse reactions to epinephrine (adrenaline) (see section 4.8).

4.8 Undesirable effects

The adverse events of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS mainly relate to the stimulation of both alpha- and beta-adrenergic receptors. The occurrence of undesirable effects depends on the sensitivity of the individual patient and the dose involved.

Immune system disorders:

Anaphylaxis, possibly with severe bronchospasm (see section 4.4).

Metabolism and nutrition disorders:

Hypokalaemia, metabolic acidosis (see section 4.4).

Disturbances of glucose metabolism, inhibition of insulin secretion and hyperglycaemia even with low doses, gluconeogenesis, glycolysis, lipolysis and ketogenesis.

Psychiatric disorders:

Psychotic states, anxiety, fear, confusion, irritability, insomnia, restlessness.

Nervous system disorders:

Headache, dizziness, tremors.

In patients with Parkinsonian syndrome, ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS increases rigidity and tremor.

Subarachnoid haemorrhage and hemiplegia have resulted from hypertension, even following subcutaneous administration of usual doses of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS.

Cardiac disorders:

Disturbances of cardiac rhythm and rate may result in palpitation and tachycardia. There may also be a reflex bradycardia, but stimulation of β_1 -adrenergic receptors of the heart may produce cardiac dysrhythmias, anginal pain and cardiac arrest.

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS can cause potentially fatal ventricular dysrhythmias including fibrillation, especially in patients with organic heart disease or those receiving other medicines that sensitise the heart to dysrhythmias. Myocardial ischaemia and myocardial infarction have been reported.

Epinephrine (adrenaline) causes electrocardiogram (ECG) changes including a decrease in T-wave amplitude in all leads in normal subjects.

In rare cases stress cardiomyopathy has been seen in patients treated with epinephrine (adrenaline).

Vascular disorders:

Stimulation of α -adrenergic receptors produces vasoconstriction with resultant hypertension. This vasoconstriction is sometimes sufficiently severe to produce gangrene when ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS is infiltrated into the digits. The rise in blood pressure may produce cerebral haemorrhage and pulmonary oedema.

Coldness of extremities may occur even with small doses of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS.

Bowel necrosis, hypotension with dizziness and fainting, and flushing may occur.

Respiratory, thoracic and mediastinal disorders:

Dyspnoea. Pulmonary oedema may occur after excessive doses or in extreme sensitivity.

Gastrointestinal disorders:

Dry mouth, reduced appetite, nausea, vomiting, hypersalivation.

Renal and urinary disorders:

Difficulty in micturition, urinary retention.

General disorders and administration site conditions:

Sweating, weakness, pallor.

Extravasation of parenterally-administered catecholamines may result in tissue necrosis and sloughing. Repeated injections of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS can cause necrosis as a result of vascular constriction at the injection site. Tissue necrosis may also occur in the extremities, kidneys and liver.

Reporting of suspected adverse reactions:

Reporting suspected adverse reactions after authorisation of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS is important. It allows continued monitoring of the benefit/risk balance of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESSENIUS. Health care providers are asked to report any suspected adverse reactions to SAHPRA via the “**6.04 Adverse Drug Reactions**”

Reporting Form", found online under SAHPRA's publications:

<https://www.sahpra.org.za/Publications/Index/8>.

Health care providers are asked to report any suspected adverse drug reactions to the Holder of the Certificate of Registration at the following email address: safety.fksa@fresenius-kabi.com and to the relevant medicine's regulatory authority in the country where the product is marketed.

4.9 Overdose

Symptoms:

See section 4.8.

After overdosage or inadvertent intravenous administration of usual intramuscular subcutaneous doses of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS, systolic and diastolic blood pressure rise sharply; venous pressure also rises. Cerebrovascular or other haemorrhages and hemiplegia may result, especially in elderly patients. Pulmonary oedema may occur.

ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS overdosage causes transient bradycardia followed by tachycardia and may cause other potentially fatal cardiac dysrhythmias. Kidney failure, metabolic acidosis and cold white skin may also occur.

Treatment:

Treatment of overdosage is symptomatic and supportive.

Because of the short duration of the adverse effects of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS, due to inactivation in the body, treatment of severe toxic reactions in hypersensitive patients or after overdose is primarily supportive. Prompt injection of a rapidly acting alpha-adrenoceptor blocking medicine, such as phentolamine, followed by a beta blocker, such as

propranolol, has been tried to counteract the pressor and dysrhythmogenic effects of ADRENALINE (EPINEPHRINE) 1 mg/1 ml FRESENIUS; rapidly-acting vasodilators, such as glyceryl trinitrate have also been used.

5. PHARMACOLOGICAL PROPERTIES

Category and class: A. 5.1 Adrenomimetics (sympathomimetics)

Pharmacotherapeutic group: adrenergic and dopaminergic agents, adrenaline.

ATC code: C01 CA 24

Epinephrine (adrenaline) is an adrenomimetic hormone.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium chloride,

sodium hydroxide (for pH-adjustment) (E524),

sodium metabisulphite (antioxidant) (E223),

tartaric acid (for pH-adjustment) (E334),

water for injection.

6.2 Incompatibilities

In the absence of compatibility studies, this medicine must not be mixed with other medicines.

6.3 Shelf life

12 months.

6.4 Special precautions for storage

Store at or below 25 °C. Protect from light.

6.5 Nature and contents of container

1 ml amber or clear ampoule in containers of 10.

6.6 Special precautions for disposal and other handling

None.

7. HOLDER OF CERTIFICATE OF REGISTRATION

Fresenius Kabi Manufacturing SA (Pty) Ltd

6 Gibaud Road

Korsten 6020

Gqeberha

South Africa

8. APPLICATION NUMBER

C923 (Act 101 of 1965)

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Not applicable.

10. DATE OF REVISION OF THE TEXT

04 October 2022