

SOTAURIC[®] (midostaurin)

25 mg soft capsules

Professional Information

Document status: Final

Approval date: January 2020

SCHEDULING STATUS: S4

1. NAME OF THE MEDICINE

Sotauric® 25 mg soft capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each soft gelatine (porcine) capsule contains 25 mg of midostaurin.

Excipients with known effect

Each soft capsule contains approximately 83 mg ethanol anhydrous and 415 mg macrogolglycerol hydroxystearate.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Soft capsule (capsule).

Pale orange, oblong capsule with red imprint "PKC NVR". The dimensions of the capsule are approximately 25.4 x 9.2 mm.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Sotauric is indicated:

- in combination with standard daunorubicin and cytarabine induction and high-dose cytarabine consolidation chemotherapy, and for patients in complete response followed by Sotauric single agent maintenance therapy, for adult patients with newly diagnosed

acute myeloid leukaemia (AML) who are FLT3 mutation-positive (see section 4.2);

- as monotherapy for the treatment of adult patients with aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated haematological neoplasm (SM-AHN), or mast cell leukaemia (MCL).

4.2 Posology and method of administration

Treatment with Sotauric should be initiated by a medical practitioner experienced in the use of anti-cancer therapies.

Before taking midostaurin, AML patients must have confirmation of FLT3 mutation (internal tandem duplication [ITD] or tyrosine kinase domain [TKD]) using a validated test.

Posology

Sotauric should be taken orally twice daily at approximately 12-hour intervals. The capsules should be taken with food (see sections 4.5 and 5.2).

Prophylactic antiemetics should be administered in accordance with local medical practice as per patient tolerance.

AML

The recommended dose of Sotauric is 50 mg orally twice daily.

Sotauric is administered on days 8-21 of induction and consolidation chemotherapy cycles, and then, for patients in complete response, maintenance therapy with Sotauric daily as single medicine, until relapse or for up to 12 cycles of 28 days each (see section 4.1).

In patients receiving a haematopoietic stem cell transplant (SCT), Sotauric should be discontinued 48 hours prior to the conditioning regimen for SCT.

Dose modifications in AML

Recommendations for dose modifications of Sotauric in patients with AML are provided in Table 1.

Table 1 Sotauric dose interruption, reduction and discontinuation recommendations in patients with AML.

Phase	Criteria	Sotauric dosing
Induction, consolidation and maintenance	Grade 3/4 pulmonary infiltrates	Interrupt Sotauric for the remainder of the cycle. Resume Sotauric at the same dose when infiltrate resolves to Grade ≤ 1 .
	Other Grade 3/4 non-haematological toxicities	Interrupt Sotauric until toxicities considered at least possibly related to Sotauric have resolved to Grade ≤ 2 , then resume Sotauric.
	QTc interval > 470 msec and ≤ 500 msec	Decrease Sotauric to 50 mg once daily for the remainder of the cycle. Resume Sotauric at the initial dose in the next cycle provided that QTc interval improves to ≤ 470 msec at the start of that cycle. Otherwise continue Sotauric 50 mg once daily.
	QTc interval > 500 msec	Withhold or interrupt Sotauric for the remainder of the cycle. If QTc improves to ≤ 470 msec just prior to the next cycle, resume Sotauric at the initial dose. If QTc interval is not improved in time to start the next cycle, do not administer Sotauric during that cycle. Sotauric may be held for as many cycles as necessary until QTc improves.
Maintenance only	Grade 4 neutropenia (ANC < $0.5 \times 10^9/l$)	Interrupt Sotauric until ANC $\geq 1.0 \times 10^9/l$, then resume at 50 mg twice daily. If neutropenia (ANC < $1.0 \times 10^9/l$) persists > 2 weeks and is suspected to be related to Sotauric, discontinue Sotauric
	Persistent Grade 1/2 toxicity	Persistent Grade 1 or 2 toxicity that patients deem unacceptable may prompt an interruption for as many as 28 days.
ANC: Absolute Neutrophil Count		

ASM, SM-AHN and MCL

The recommended starting dose of Sotauric is 100 mg orally twice daily. Treatment should be continued as long as clinical benefit is observed or until unacceptable toxicity occurs.

Dose modifications in ASM, SM-AHN and MCL

Recommendations for dose modifications of Sotauric in patients with ASM, SM-AHN and MCL are provided in Table 2.

Table 2 Sotauric dose interruption, reduction and discontinuation recommendations in patients with ASM, SM-AHN or MCL

Criteria	Sotauric dosing
<p>ANC $<1.0 \times 10^9/l$ attributed to Sotauric in patients without MCL, or ANC less than $0.5 \times 10^9/l$ attributed to Sotauric in patients with baseline ANC value of $0.5-1.5 \times 10^9/l$</p>	<p>Interrupt Sotauric until ANC $\geq 1.0 \times 10^9/l$, then resume at 50 mg twice daily and, if tolerated, increase to 100 mg twice daily. Discontinue Sotauric if low ANC persists for > 21 days and is suspected to be related to Sotauric.</p>
<p>Platelet count less than $50 \times 10^9/l$ attributed to Sotauric in patients without MCL, or platelet count less than $25 \times 10^9/l$ attributed to Sotauric in patients with baseline platelet count of $25-75 \times 10^9/l$</p>	<p>Interrupt Sotauric until platelet count greater than or equal to $50 \times 10^9/l$, then resume Sotauric at 50 mg twice daily and, if tolerated, increase to 100 mg twice daily. Discontinue Sotauric if low platelet count persists for > 21 days and is suspected to be related to Sotauric.</p>
<p>Haemoglobin less than 8 g/dl attributed to Sotauric in patients without MCL, or life-threatening anaemia attributed to Sotauric in patients with baseline haemoglobin value of 8-10 g/dl</p>	<p>Interrupt Sotauric until haemoglobin greater than or equal to 8 g/dl, then resume Sotauric at 50 mg twice daily and, if tolerated, increase to 100 mg twice daily. Discontinue Sotauric if low haemoglobin persists for > 21 days and is suspected to be related to Sotauric.</p>

Criteria	Sotauric dosing
Grade 3/4 nausea and/or vomiting despite optimal anti-emetic therapy	Interrupt Sotauric for 3 days (6 doses), then resume at 50 mg twice daily and, if tolerated, gradually increase to 100 mg twice daily.
Other Grade 3/4 non-haematological toxicities	<p>Interrupt Sotauric until event has resolved to Grade \leq 2, then resume Sotauric at 50 mg twice daily and, if tolerated, increase to 100 mg twice daily.</p> <p>Discontinue Sotauric if toxicity is not resolved to Grade \leq 2 within 21 days or severe toxicity recurs at a reduced dose of Sotauric.</p>
<p>ANC: Absolute Neutrophil Count</p> <p>CTCAE severity: Grade 1 = mild symptoms; 2 = moderate symptoms; 3 = severe symptoms; 4 = life-threatening symptoms.</p>	

Missed doses

If a dose is missed, the patient should take the next dose at the scheduled time.

If vomiting occurs, the patient should not take an additional dose of Sotauric, but should take the next scheduled dose.

Special populations

Elderly (\geq 65 years)

No dose adjustment is required in patients aged over 65 years (see section 5.2). **There is limited experience with midostaurin in AML patients aged 60-70 years and no experience in AML patients above 70 years.**

In patients aged \geq 60 years, Sotauric should be used only in patients eligible to receive intensive induction chemotherapy with adequate performance status and without significant

comorbidities.

Renal impairment

No dose adjustment is required for patients with mild or moderate renal impairment. Clinical experience in patients with severe renal impairment is limited and no data are available in patients with end-stage renal disease (see sections 4.4 and 5.2).

Hepatic impairment

No dose adjustment is required in patients with mild or moderate (Child-Pugh A or B) hepatic impairment (see section 5.2). Safety in patients with severe (Child-Pugh C) hepatic impairment has not been established (see section 4.4).

Acute promyelocytic leukaemia

Sotauric has not been studied in patients with acute promyelocytic leukaemia and therefore its use is not recommended in this patient population.

Paediatric population

The safety and efficacy of Sotauric in children and adolescents below 18 years have not been established (see section 5.1). Currently available data are described in section 5.2 but no recommendation on a posology can be made.

Method of administration

Sotauric is for oral use.

Sotauric should be taken with food

The capsules should be swallowed whole with a glass of water. They should not be opened, crushed or chewed to ensure proper dosing and avoid the unpleasant taste of the capsule content.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Concomitant administration of potent CYP3A4 inducers, e.g. rifampicin, St. John's Wort (*Hypericum perforatum*), carbamazepine, enzalutamide, phenytoin (see section 4.5).

Cardiovascular disorders: Congestive cardiac failure NYHA class III or IV; left ventricular ejection fraction (LVEF) < 50%; myocardial infarction within the previous 6 months; heart block of any degree; congenital or acquired prolongation of QT time; poorly controlled hypertension.

Pregnancy and lactation

4.4 Special warnings and precautions for use

Neutropenia and infections

Neutropenia has occurred in patients receiving Sotauric as monotherapy and in combination with chemotherapy (see section 4.8). Severe neutropenia (ANC < 0.5 x 10⁹/l) was generally reversible by withholding Sotauric until recovery and discontinuation in the ASM, SM-AHN and MCL studies. White blood cell counts (WBCs) should be monitored regularly, especially at treatment initiation.

In patients who develop unexplained severe neutropenia, treatment with Sotauric should be interrupted until ANC is ≥ 1.0 x 10⁹/l, as recommended in Tables 1 and 2. Sotauric should be

discontinued in patients who develop recurrent or prolonged severe neutropenia that is suspected to be related to Sotauric (see section 4.2).

Any active serious infection should be under control prior to starting treatment with Sotauric monotherapy. Patients should be monitored for signs and symptoms of infection, including any device-related infections, and if a diagnosis of infection is made appropriate treatment must be instituted promptly, including, as needed, the discontinuation of Sotauric.

Cardiac dysfunction

Patients with symptomatic congestive heart failure were excluded from clinical studies (see section 4.3). In the ASM, SM-AHN and MCL studies cardiac dysfunction such as congestive heart failure (CHF) (including some fatalities) and transient decreases in left ventricular ejection fraction (LVEF) occurred. In the randomised AML study no difference in CHF was observed between the Sotauric + chemotherapy and placebo + chemotherapy arms. In patients at risk, Sotauric should be used with caution and the patient closely monitored by assessing LVEF when clinically indicated (at baseline and during treatment).

An increased frequency of QTc prolongation was noted in midostaurin-treated patients (see section 4.8), however, a mechanistic explanation for this observation was not found (see section 4.3). Caution is warranted in patients at risk of QTc prolongation (e.g. due to concomitant medicines and/or electrolyte disturbances). Interval assessments of QT by ECG should be considered if Sotauric is taken concurrently with medicines that can prolong QT interval (see section 4.3).

Pulmonary toxicity

Interstitial lung disease (ILD) and pneumonitis, in some cases fatal, have occurred in patients treated with Sotauric monotherapy or in combination with chemotherapy. Patients should be

monitored for pulmonary symptoms indicative of ILD or pneumonitis and Sotauric discontinued in patients who experience pulmonary symptoms indicative of ILD or pneumonitis that are \geq Grade 3 (NCI CTCAE).

Embryofoetal toxicity and breast-feeding

Pregnant women should be informed of the potential risk to a foetus; females of reproductive potential should be advised to have a pregnancy test within 7 days prior to starting treatment with Sotauric and to use effective contraception during treatment with Sotauric and for at least 4 months after stopping treatment. Women using hormonal contraceptives should add a barrier method of contraception.

Because of the potential for serious adverse reactions in breast-feeding infants from Sotauric, women should discontinue breast-feeding during treatment with Sotauric and for at least 4 months after stopping treatment (see section 4.6).

Severe hepatic impairment

Safety of Sotauric has not been established in patients with severe hepatic impairment. Caution is warranted when considering the administration of midostaurin in patients with severe hepatic impairment and patients should be carefully monitored for toxicity (see section 5.2).

Severe renal impairment

Caution is warranted when considering the administration of midostaurin in patients with severe renal impairment or end-stage renal disease and patients should be carefully monitored for toxicity (see section 5.2).

Interactions

Caution is advised with the concomitant use of Sotauric with medicines that are strong inhibitors of CYP3A4, such as, but not limited to, antifungals (e.g. ketoconazole), certain antivirals (e.g. ritonavir), macrolide antibiotics (e.g. clarithromycin) and nefazodone because they can increase the plasma concentrations of midostaurin especially when (re-)starting with midostaurin treatment (see section 4.5). Alternative medicines that do not strongly inhibit CYP3A4 activity should be considered. In situations where satisfactory therapeutic alternatives do not exist, patients should be closely monitored for midostaurin-related toxicity.

Excipients

Sotauric contains macrogolglycerol hydroxystearate, which may cause stomach discomfort and diarrhoea.

A 100 mg dose of Sotauric contains approximately 14 vol. % ethanol anhydrous, which corresponds to 333 mg alcohol. This is equivalent to 8.4 ml beer or 3.5 ml wine. Alcohol may be harmful in patients with alcohol-related problems, epilepsy or liver problems or during pregnancy or breast-feeding.

4.5 Interaction with other medicines and other forms of interaction

Midostaurin undergoes extensive hepatic metabolism mainly through CYP3A4 enzymes which are either induced or inhibited by a number of concomitant medicines.

Effect of other medicines on Sotauric

Medicines or substances known to affect the activity of CYP3A4 may affect the plasma

concentrations of midostaurin and therefore the safety and/or efficacy of Sotauric.

Strong CYP3A4 inducers

Concomitant use of Sotauric with strong inducers of CYP3A4 (e.g. carbamazepine, rifampicin, enzalutamide, phenytoin, St. John's Wort [*Hypericum perforatum*]) is contraindicated (see section 4.3). Strong CYP3A4 inducers decrease exposure of midostaurin and its active metabolites (CGP52421 and CGP62221). In a study in healthy subjects, co-administration of the strong CYP3A4 inducer rifampicin (600 mg daily) to steady state with a 50 mg single dose of midostaurin decreased midostaurin C_{max} by 73 % and AUC_{inf} by 96 % on average, respectively. CGP62221 exhibited a similar pattern. The mean AUC_{last} of CGP52421 decreased by 60 %.

Strong CYP3A4 inhibitors

Strong CYP3A4 inhibitors may increase midostaurin blood concentrations. In a study with 36 healthy subjects, co-administration of the strong CYP3A4 inhibitor ketoconazole to steady state with a single dose of 50 mg midostaurin, led to a significant increase in midostaurin exposure (1.8-fold C_{max} increase and 10-fold AUC_{inf} increase) and 3.5-fold increase in AUC_{inf} of CGP62221, while the C_{max} of the active metabolites (CGP62221 and CGP52421) decreased by half (see section 5.2). At steady state of midostaurin (50 mg twice daily for 21 days), with the strong CYP3A4 inhibitor itraconazole at steady state in a subset of patients (N=7), midostaurin steady-state exposure (C_{min}) was increased by 2.09-fold. C_{min} of CGP52421 was increased by 1.3-fold, whereas no significant effect in exposure of CGP62221 was observed (see section 4.4).

Effect of Sotauric on other medicines

Midostaurin is not an inhibitor of CYP3A4 *in vivo*. The pharmacokinetics of midazolam (sensitive CYP3A4 probe) were not affected following three days' dosing of midostaurin in healthy subjects.

Based on *in vitro* data, midostaurin and/or its metabolites have the potential to inhibit CYP1A2, CYP2D6, CYP2C8, CYP2C9, CYP2E1 and CYP3A4/5 enzymes.

Based on *in vitro* data, midostaurin and/or its metabolites have the potential to induce CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19 and CYP3A4/5 enzymes. Midostaurin inhibited OATP1B1, BCRP and P-glycoprotein (P-gp) *in vitro* (see section 5.2). The combination of data on *in vivo* midostaurin auto-induction upon repeated dosing and increase in plasma 4 β -OH cholesterol levels suggest that midostaurin may be at least a moderate CYP3A4 inducer *in vivo*.

In vivo studies have not been conducted for the investigation of induction and inhibition of enzymes and transporters by midostaurin and the active metabolites. Medicines with a narrow therapeutic range that are substrates of CYP1A2 (e.g. tizanidine), CYP2D6 (e.g. codeine), CYP2C8 (e.g. paclitaxel), CYP2C9 (e.g. warfarin), CYP2C19 (e.g. omeprazole), CYP2E1 (e.g. chlorzoxazone), CYP3A4/5 (e.g. tacrolimus), CYP2B6 (e.g. efavirenz), P-gp (e.g. paclitaxel), BCRP (e.g. atorvastatin) or OATP1B1 (e.g. digoxin), should be used with caution when administered concomitantly with midostaurin and may need dose adjustment to maintain optimal exposure (see section 5.2).

It is currently unknown whether midostaurin may reduce the effectiveness of hormonal contraceptives, and therefore women using hormonal contraceptives should add a barrier method of contraception (see section 4.6).

Food interactions

In healthy subjects, midostaurin absorption (AUC) was increased by an average of 22 % when Sotauric was co-administered with a standard meal and by an average of 59 % when co-administered with a high-fat meal. Peak midostaurin concentration (C_{max}) was reduced by 20 % with a standard meal and by 27 % with a high-fat meal versus on an empty stomach (see section 5.2).

Sotauric is recommended to be administered with food.

4.6 Fertility, pregnancy and lactation

Sotauric is contraindicated in pregnancy and lactation

Women of childbearing potential

Women of childbearing potential should be informed that animal studies show midostaurin to be harmful to the developing foetus. Sexually active women of childbearing potential are advised to have a pregnancy test within 7 days prior to starting treatment with Sotauric and that they should use effective contraception (methods that result in less than 1% pregnancy rates) when using Sotauric and for at least 4 months after stopping treatment with Sotauric. It is currently unknown whether midostaurin may reduce the effectiveness of hormonal contraceptives, and therefore women using hormonal contraceptives should add a barrier method of contraception.

Pregnancy

Midostaurin can cause foetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies in pregnant women. Reproductive studies in rats and rabbits demonstrated that midostaurin induced foetotoxicity (see section 5.3). Sotauric should

not be used during pregnancy or in women of childbearing potential not using contraception (see section 4.3). Pregnant women should be advised of the potential risk to the foetus.

Breast-feeding

It is unknown whether midostaurin or its active metabolites are excreted in human milk. Available animal data have shown that midostaurin and its active metabolites pass into the milk of lactating rats. Breast-feeding should be discontinued during treatment with Sotauric and for at least 4 months after stopping treatment.

Fertility

There are no data on the effect of Sotauric on human fertility. Animal studies with midostaurin have shown impaired fertility (see section 5.3).

4.7 Effects on ability to drive and use machines

Sotauric may influence the ability to drive and use machines. Dizziness and vertigo have been reported in patients taking Sotauric and should be considered when assessing a patient's ability to drive or use machines.

4.8 Undesirable effects

Summary of the safety profile

AML

The safety evaluation of Sotauric (50 mg twice daily) in patients with newly diagnosed FLT3-mutated AML is based on a phase III, randomised, double-blind, placebo-controlled study with 717 patients. The overall median duration of exposure was 42 days (range 2 to 576 days) for patients in the Sotauric plus standard chemotherapy arm versus 34 days (range

1 to 465 days) for patients in the placebo plus standard chemotherapy arm.

For the 205 patients (120 in Sotauric arm and 85 in placebo arm) who entered the maintenance phase, the median duration of exposure in maintenance was 11 months for both arms (16 to 520 days for patients in the Sotauric arm and 22 to 381 days in the placebo arm).

The most frequent adverse drug reactions (ADRs) in the Sotauric arm were febrile neutropenia (83.4 %), nausea (83.4 %), exfoliative dermatitis (61.6 %), vomiting (60.7 %), headache (45.9 %), petechiae (35.8 %) and pyrexia (34.5 %). The most frequent Grade 3/4 ADRs were febrile neutropenia (83.5 %), lymphopenia (20.0%), device-related infection (15.7 %), exfoliative dermatitis (13.6 %), hyperglycaemia (7.0 %) and nausea (5.8 %). The most frequent laboratory abnormalities were haemoglobin decreased (97.3%), ANC decreased (86.7 %), ALT increased (84.2 %), AST increased (73.9 %) and hypokalaemia (61.7 %). The most frequent Grade 3/4 laboratory abnormalities were ANC decreased (85.8 %), haemoglobin decreased (78.5 %), ALT increased (19.4 %) and hypokalaemia (13.9 %).

Serious ADRs occurred at similar rates in patients in the Sotauric versus the placebo arm. The most frequent serious ADR in both arms was febrile neutropenia (16 %).

Discontinuation due to any adverse reaction occurred in 3.1 % of patients in the Sotauric arm versus 1.3 % in the placebo arm. The most frequent Grade 3/4 adverse reaction leading to discontinuation in the Sotauric arm was exfoliative dermatitis (1.2 %).

Safety profile during maintenance phase

While Table 3 provides the incidence for ADRs over the total duration of the study, when the maintenance phase (single agent Sotauric or placebo) was assessed separately, a difference in the type and severity of ADRs was observed. The overall incidence of ADRs during the

maintenance phase was generally lower than during the induction and consolidation phase. Incidences of ADRs were, however, higher in the Sotauric arm than in the placebo arm during the maintenance phase. ADRs occurring more often in the midostaurin arm versus placebo during maintenance included: nausea (46.4 % versus 17.9 %), hyperglycaemia (20.2 % versus 12.5 %), vomiting (19 % versus 5.4 %) and QT prolongation (11.9 % versus 5.4 %).

Most of the haematological abnormalities reported, occurred during the induction and consolidation phase when the patients received Sotauric or placebo in combination with chemotherapy. The most frequent Grade 3/4 haematological abnormalities reported in patients during the maintenance phase with Sotauric were ANC decrease (20.8 % versus 18.8 %) and leukopaenia (7.5 % versus 5.9 %).

ADRs reported during the maintenance phase led to discontinuation of 1.2 % of patients in the Sotauric arm and none in the placebo arm.

ASM, SM-AHN and MCL

The safety of Sotauric (100 mg twice daily) as a single agent in patients with ASM, SM-AHN and MCL was evaluated in 142 patients in two single-arm, open-label, multicentre studies. The median duration of exposure to Sotauric was 11.4 months (range: 0 to 81 months).

The most frequent ADRs were nausea (82 %), vomiting (68 %), diarrhoea (51 %), peripheral oedema (35 %) and fatigue (31 %). The most frequent Grade 3/4 ADRs were fatigue (8.5 %), sepsis (7.7 %), pneumonia (7 %), febrile neutropenia (7 %), and diarrhoea (6.3 %). The most frequent non-haematological laboratory abnormalities were hyperglycaemia (93.7 %), total bilirubin increased (40.1 %), lipase increased (39.4 %), aspartate aminotransferase (AST)

increased (33.8%), and alanine aminotransferase (ALT) increased (33.1 %), while the most frequent haematological laboratory abnormalities were absolute lymphocyte count decreased (73.2 %) and ANC decreased (58.5 %). The most frequent Grade 3/4 laboratory abnormalities were absolute lymphocyte count decreased (45.8 %), ANC decreased (26.8 %), hyperglycaemia (19 %), and lipase increased (17.6 %).

Dose modifications (interruption or adjustment) due to ADRs occurred in 31 % of patients. The most frequent ADRs that led to dose modification (incidence ≥ 5 %) were nausea and vomiting. ADRs that led to treatment discontinuation occurred in 9.2 % of patients. The most frequent (incidence ≥ 1 %) were febrile neutropenia, nausea, vomiting and pleural effusion.

Tabulated lists of adverse drug reactions

ADRs are listed according to MedDRA system organ class. Within each system organ class, the ADRs are ranked by frequency, with the most frequent reactions first, using the following convention (CIOMS III): very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

AML

Table 3 presents the frequency category of ADRs reported in the phase III study in patients with newly diagnosed FLT3-mutated AML.

Table 3 Adverse drug reactions observed in the AML clinical study

Adverse drug reaction	All grades	Grades 3/4	Frequency category
	Sotauric + chemo n=229 ¹ %	Sotauric + chemo n=345 ¹ %	
Infections and infestations			
Device-related infection	24	15.7	Very common
Upper respiratory tract infection	5.2	0.6	Common
Neutropenic sepsis	0.9	3.5	Uncommon
Blood and lymphatic system disorders			
Febrile neutropenia; thrombocytopenia	83.4	83.5	Very common
Petechiae	35.8	1.2	Very common
Lymphopenia	16.6	20	Very common
Immune system disorders			
Hypersensitivity	15.7	0.6	Very common
Metabolism and nutrition disorders			
Hyperuricaemia	8.3	0.6	Common
Psychiatric disorders			
Insomnia	12.2	0	Very common
Nervous system disorders			
Headache	45.9	2.6	Very common
Syncope	5.2	4.6	Common
Tremor	3.9	0	Common
Eye disorders			
Eyelid oedema	3.1	0	Common
Cardiac disorders			

Hypotension	14.4	5.5	Very common
Sinus tachycardia; QT prolongation	9.6	1.2	Common
Hypertension	7.9	2.3	Common
Pericardial effusion	3.5	0.6	Common
Respiratory, thoracic and mediastinal disorders			
Epistaxis; interstitial lung disease and pneumonitis	27.5	2.6	Very common
Laryngeal pain	11.8	0.6	Very common
Dyspnoea	10.9	5.5	Very common
Pleural effusion	5.7	0.9	Common
Nasopharyngitis	8.7	0	Common
Acute respiratory distress syndrome	2.2	2.3	Common
Gastrointestinal disorders			
Nausea	83.4	5.8	Very common
Vomiting	60.7	2.9	Very common
Stomatitis	21.8	3.5	Very common
Abdominal pain upper	16.6	0	Very common
Haemorrhoids	15.3	1.4	Very common
Anorectal discomfort	7	0.9	Common
Abdominal discomfort	3.5	0	Common
Skin and subcutaneous tissue disorders			
Dermatitis exfoliative	61.6	13.6	Very common
Hyperhidrosis	14.4	0	Very common
Dry skin	7	0	Common
Keratitis	6.6	0.3	Common
Musculoskeletal and connective tissue disorders			
Back pain	21.8	1.4	Very common
Arthralgia	14	0.3	Very common
Bone pain	9.6	1.4	Common
Pain in extremity	9.6	1.4	Common
Neck pain	7.9	0.6	Common
General disorders and administration site conditions			

Pyrexia	34.5	3.2	Very common
Catheter-related thrombosis	3.5	2	Common
Investigations			
Haemoglobin decreased*	97.3	78.5	Very common
ANC decreased*	86.7	85.8	Very common
ALT increased*	84.2	19.4	Very common
AST increased*	73.9	6.4	Very common
Hypokalaemia*	61.7	13.9	Very common
Hyperglycaemia	20.1	7	Very common
Hypernatraemia*	20	1.2	Very common
Activated partial thromboplastin time prolonged	12.7	2.6	Very common
Hypercalcaemia*	6.7	0.6	Common
Weight increased	6.6	0.6	Common
<p>¹For trial sites in North America, all grades were collected for 13 pre-specified adverse events. For all other adverse events, only grades 3 and 4 were collected. Therefore all grade AEs are summarised only for patients in non-North American trial sites, whereas Grades 3 and 4 are summarised for patients in all trial sites.</p> <p>* Frequency is based on laboratory values.</p>			

ASM, SM-AHN and MCL

Table 4 presents the frequency category of ADRs based on pooled data from two studies in patients with ASM, SM-AHN and MCL.

Table 4 Adverse drug reactions observed in the ASM, SM-AHN and MCL clinical studies

Adverse drug reaction	Sotauric (100 mg twice daily) N=142		Frequency category
	All grades %	Grades 3/4 %	
Infections and infestations			
Urinary tract infection	13	2.8	Very common
Upper respiratory tract infection	11	1.4	Very common
Pneumonia	8.5	7.0	Common
Sepsis	7.7	7.7	Common
Bronchitis	5.6	0	Common
Oral herpes	4.9	0	Common
Cystitis	4.2	0	Common
Sinusitis	4.2	0.7	Common
Erysipelas	3.5	1.4	Common
Herpes zoster	3.5	0.7	Common
Blood and lymphatic system disorders			
Febrile neutropenia	7.7	7.0	Common
Thrombocytopenia			
Immune system disorders			
Hypersensitivity	2.1	0	Common
Anaphylactic shock	0.7	0.7	Uncommon
Nervous system disorders			
Headache	26	1.4	Very common
Dizziness	13	0	Very common
Disturbance in attention	7	0	Common
Tremor	6.3	0	Common
Ear and labyrinth disorders			
Vertigo	4.9	0	Common
Vascular disorders			
Hypotension	9.2	2.1	Common
Haematoma	6.3	0.7	Common
Respiratory, thoracic and mediastinal disorders			
Dyspnoea; interstitial lung	18	5.6	Very common

disease and pneumonitis			
Cough	16	0.7	Very common
Pleural effusion	13	4.2	Very common
Epistaxis	12	2.8	Very common
Oropharyngeal pain	4.2	0	Common
Gastrointestinal disorders			
Nausea	82	5.6	Very common
Vomiting	68	5.6	Very common
Diarrhoea	51	6.3	Very common
Constipation	29	0.7	Very common
Dyspepsia	5.6	0	Common
Gastrointestinal haemorrhage	4.2	3.5	Common
General disorders and administration site conditions			
Oedema peripheral	35	3.5	Very common
Fatigue	31	8.5	Very common
Pyrexia	27	4.2	Very common
Asthenia	4.9	0.7	Common
Chills	4.9	0	Common
Oedema	4.2	0.7	Common
Investigations			
Hyperglycaemia (non-fasting)*	93.7	19.0	Very common
Absolute lymphocyte decreased*	73.2	45.8	Very common
ANC decreased*	58.5	26.8	Very common
Total bilirubin increased*	40.1	4.9	Very common
Lipase increased*	39.4	17.6	Very common
AST increased*	33.8	2.8	Very common
ALT increased*	33.1	3.5	Very common
Amylase increased*	20.4	7.0	Very common
Weight increased	5.6	2.8	Common
Injury, poisoning and procedural complications			
Contusion	6.3	0	Common
Fall	4.2	0.7	Common
* Frequency is based on laboratory values.			

Description of selected adverse drug reactions

Gastrointestinal disorders

Nausea, vomiting and diarrhoea were observed in AML, ASM, SM-AHN and MCL patients. In ASM, SM-AHN and MCL patients these events led to dose adjustment or interruption in 26 % and to discontinuation in 4.2 % of the patients. Most of the events occurred within the first 6 months of treatment and were managed with supportive prophylactic medicines.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of Sotauric is important. It allows continued monitoring of the benefit/risk balance of Sotauric.

4.9 Overdose

Reported experience with overdose in humans is very limited. Single doses of up to 600 mg have been given with acceptable acute tolerability. Adverse reactions observed were diarrhoea, abdominal pain and vomiting.

There is no known specific antidote for midostaurin. In the event of an overdose, patients must be closely monitored for signs or symptoms of adverse reactions, and appropriate symptomatic and supportive treatment initiated.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, protein kinase inhibitors, ATC code: L01XE39

Mechanism of action

Midostaurin inhibits multiple receptor tyrosine kinases, including FLT3 and KIT kinase. Midostaurin inhibits FLT3 receptor signalling and induces cell cycle arrest and apoptosis in leukaemic cells expressing FLT3 ITD or TKD mutant receptors or over-expressing FLT3 wild type receptors. *In vitro* data indicate that midostaurin inhibits D816V mutant KIT receptors at exposure levels achieved in patients (average achieved exposure higher than IC₅₀). *In vitro* data indicate that KIT wild type receptors are inhibited to a much lesser extent at these concentrations (average achieved exposure lower than IC₅₀). Midostaurin interferes with aberrant KIT D816V-mediated signalling and inhibits mast cell proliferation, survival and histamine release.

In addition, midostaurin inhibits several other receptor tyrosine kinases such as PDGFR (platelet-derived growth factor receptor) or VEGFR2 (vascular endothelial growth factor receptor 2), as well as members of the serine/threonine kinase family PKC (protein kinase C). Midostaurin binds to the catalytic domain of these kinases and inhibits the mitogenic signalling of the respective growth factors in cells, resulting in growth arrest.

Midostaurin in combination with chemotherapeutic agents (cytarabine, doxorubicin, idarubicin and daunorubicin) resulted in synergistic growth inhibition in FLT3-ITD expressing AML cell

lines.

Pharmacodynamic properties

Two major metabolites have been identified in murine models and humans, i.e. CGP62221 and CGP52421. In proliferation assays with FLT3-ITD expressing cells, CGP62221 showed similar potency compared to the parent compound, however CGP52421 was approximately 10-fold less potent.

Cardiac electrophysiology

A dedicated QT study in 192 healthy subjects with a dose of 75 mg twice daily did not reveal clinically significant prolongation of QT by midostaurin and CGP62221 but the study duration was not long enough to estimate the QTc prolongation effects of the long-acting metabolite CGP52421. Therefore, the change from baseline in QTcF with the concentration of midostaurin and both metabolites was further explored in a phase II study in 116 patients with ASM, SM-AHN or MCL. At the median peak C_{min} concentrations attained at a dose of 100 mg twice daily, neither midostaurin, CGP62221 nor CGP52421 showed a potential to cause clinically significant QTcF prolongation, since the upper bounds of predicted change at these concentration levels were less than 10 msec (5.8, 2.4, and 4.0 msec, respectively). In the ASM, SM-AHN and MCL population, 25.4 % of patients had at least one ECG measurement with a QTcF greater than 450 ms and 4.7 % greater than 480 ms.

Clinical efficacy

AML

The efficacy and safety of midostaurin in combination with standard chemotherapy versus placebo plus standard chemotherapy and as single agent maintenance therapy was

investigated in 717 patients (18 to 60 years of age) in a randomised, double-blind, phase III study. Patients with newly diagnosed FLT3-mutated AML as determined by a clinical study assay were randomised (1:1) to receive midostaurin 50 mg twice daily (n=360) or placebo (n=357) sequentially in combination with standard daunorubicin (60 mg/m² daily on days 1-3) / cytarabine (200 mg/m² daily on days 1-7) induction and high-dose cytarabine (3 g/m² every 12 hours on days 1, 3, 5) consolidation, followed by continuous midostaurin or placebo treatment according to initial assignment for up to 12 additional cycles (28 days/cycle). While the study included patients with various AML-related cytogenetic abnormalities, patients with acute promyelocytic leukaemia (M3) or therapy-related AML were excluded. Patients were stratified by FLT3 mutation status: TKD, ITD with allelic ratio <0.7, and ITD with allelic ratio ≥ 0.7.

The two treatment groups were generally balanced with respect to the baseline demographics of disease characteristics. The median age of the patients was 47 years (range: 18 to 60 years), a majority of the patients had ECOG performance status of 0 or 1 (88.3%), and most patients had *de novo* AML


(95 %). Of the patients with race information reported, 88.1 % were Caucasian. The majority of patients (77.4 %) had FLT3-ITD mutations, most of them (47.6 %) with a low allelic ratio (< 0.7), and 22.6 % of patients had FLT3-TKD mutations. Forty-eight per cent were male in the midostaurin arm and 41% in the placebo arm.

Patients who proceeded to haematopoietic stem cell transplant (SCT) stopped receiving study treatment prior to the start of the SCT conditioning regimen. The overall rate of SCT was 59.4 % (214/360) of patients in the midostaurin plus standard chemotherapy arm versus 55.2 % (197/357) in the placebo plus standard chemotherapy arm. All patients were followed for

survival.

The primary endpoint of the study was overall survival (OS), measured from the date of randomisation until death by any cause. The primary analysis was conducted after a minimum follow-up of approximately 3.5 years after the randomisation of the last patient. The study demonstrated a statistically significant improvement in OS with a 23 % risk reduction of death for midostaurin plus standard chemotherapy over placebo plus standard chemotherapy (see Table 6 and Figure 1).

Figure 1 Kaplan-Meier curve for overall survival, non-censored for SCT



The key secondary endpoint was event-free survival (EFS; an EFS event is defined as a failure to obtain a complete remission (CR) within 60 days of initiation of protocol therapy, or relapse, or death from any cause). The EFS showed a statistically significant improvement for midostaurin plus standard chemotherapy over placebo plus standard chemotherapy (HR: 0.78 [95 % CI, 0.66 to 0.93] p = 0.0024), and a median EFS of 8.2 months and 3.0 months, respectively; see Table 5.

Table 5 Efficacy of midostaurin in AML


Efficacy Parameter	Midostaurin n=360	Placebo n=357	HR* (95 % CI)	P-value[‡]
Overall Survival (OS)¹				
Median OS in months (95% CI)	74.7 (31.5, NE)	25.6 (18.6, 42.9)	0.77 (0.63, 0.95)	0.0078
Kaplan-Meier estimates at 5 years (95% CI)	0.51 (0.45, 0.56)	0.43 (0.38, 0.49)		
Event Free Survival (EFS)²				
Median EFS in months, considering CRs within 60 days of treatment start (95% CI)	8.2 (5.4, 10.7)	3.0 (1.9, 5.9)	0.78 (0.66, 0.93)	0.0024
Median EFS in months, considering CRs any time during induction (95% CI)	10.2 (8.1, 13.9)	5.6 (2.9, 6.7)	0.73 (0.61, 0.87)	0.0001
Disease Free Survival (DFS)				
Median DFS in months (95% CI)	26.7 (19.4, NE)	15.5 (11.3, 23.5)	0.71 (0.55, 0.92)	0.0051
Complete Remission (CR)				
within 60 days of treatment start (%)	212 (58.9)	191 (53.5)	NE	0.073 [§]
any time during induction (%)	234 (65.0)	207 (58.0)	NE	0.027 [§]
Cumulative incidence of relapse (CIR)				
Median (95% CI)	NE (25.7, NE)	17.6 (12.7, 46.3)	0.68 (0.52, 0.89)	0.0023
¹ primary endpoint; ² key secondary endpoint; NE: Not Estimated				
*Hazard ratio (HR) estimated using Cox regression model stratified according to the randomisation FLT3 mutation factor.				
[‡] 1-sided p-value calculated using log-rank test stratified according to the randomisation FLT3 mutation factor.				
[§] Not significant				

There was a trend favouring midostaurin for CR rate by day 60 for the midostaurin arm (58.9 % versus 53.5 %; $p = 0.073$) that continued when considering all CRs during induction (65.0% versus 58.0%; $p = 0.027$). In addition, in patients who achieved complete remission during induction, the cumulative incidence of relapse at 12 months was 26% in the midostaurin arm versus 41% in the placebo arm.

Sensitivity analyses for both OS and EFS when censored at the time of SCT also supported the clinical benefit with midostaurin plus standard chemotherapy over placebo.

Results for OS by SCT status are shown in Figure 2. For EFS, considering complete remissions within 60 days of study treatment start, the HR was 0.602 (95 % CI: 0.372, 0.974) for patients with SCT and 0.827 (95 % CI: 0.689, 0.993) for patients without SCT, favouring midostaurin.

Figure 2 Kaplan Meier curve for overall survival by SCT status in AML



In a subgroup analysis, no apparent OS benefit was observed in females, however, a treatment benefit

was observed in females in all secondary efficacy endpoints (see Table 6).

Table 6 Overview of OS, EFS, CR, DFS and CIR by gender in AML

Endpoint	Overall 95% CI	Males 95% CI	Females 95% CI
OS (HR)	0.774 (0.629, 0.953)	0.533 (0.392, 0.725)	1.007 (0.757, 1.338)
EFS (CR induction) (HR)	0.728 (0.613, 0.866)	0.660 (0.506, 0.861)	0.825 (0.656, 1.037)
CR induction (OR)	0.743* (0.550, 1.005)	0.675* (0.425, 1.072)	0.824* (0.552, 1.230)
DFS (CR induction) (HR)	0.663 (0.516, 0.853)	0.594 (0.408, 0.865)	0.778 (0.554, 1.093)
CIR (CR induction) (HR)	0.676 (0.515, 0.888)	0.662 (0.436, 1.006)	0.742 (0.516, 1.069)
*Odds ratio calculated as (No complete remission in treatment/Complete remission in treatment) / (No complete remission in placebo/complete remission in placebo) HR= Hazard ratio; OR=odds ratio			

Efficacy and safety in patients 60-70 years old were evaluated in a phase II, single- arm, investigator- initiated study of midostaurin in combination with intensive induction, consolidation including allogenic SCT and single-agent maintenance in patients with FLT3-ITD mutated AML. Based on an interim analysis, the EFS rate at 2 years (primary endpoint) was 27.1 % (95% CI: 16.6, 44.1) and the median OS was 15.5 months in patients older than 60 years of age (46 out of 145 patients).

ASM, SM-AHN and MCL

The efficacy of midostaurin in patients with ASM, SM-AHN and MCL, collectively referred to as advanced systemic mastocytosis (SM), was evaluated in two open-label, single-arm, multicentre studies (142 patients in total).

The pivotal study was a multicentre, single-arm phase II study in 116 patients with advanced SM (Study CPKC412D2201). Midostaurin was administered orally at 100 mg twice daily until disease progression or intolerable toxicity. Of the 116 patients enrolled, 89 were considered eligible for response assessment and constituted the primary efficacy population. Of these, 73 patients had ASM

(57 with an AHN) and 16 patients had MCL (6 with an AHN). The median age in the primary efficacy population was 64 years with approximately half of the patients \geq 65 years. Approximately one third (36 %) received prior anti-neoplastic therapy for ASM, SM-AHN or MCL. At baseline in the primary efficacy population, 65 % of the patients had $>$ 1 measurable C finding (thrombocytopenia, hypoalbuminaemia, anaemia, high total bilirubin, transfusion-dependent anaemia, weight loss, neutropenia, high ALT or high AST). The KIT D816V mutation was detected in 82 % of patients.

The primary endpoint was overall response rate (ORR). Response rates were assessed based on the modified Valent and Cheson criteria and responses were adjudicated by a study steering committee. Secondary endpoints included duration of response, time to response, and overall survival. Responses to midostaurin are shown in Table 7. Activity was observed regardless of number of prior therapies, and presence or absence of an AHN. Confirmed responses were observed in both KIT D816V mutation positive patients (ORR=63 %) and KIT D816V wild type or unknown patients (ORR=43.8 %). However, the median survival for KIT D816V positive patients was longer, i.e. 33.9 months (95 % CI: 20.7, 42), than for KIT D816V wild type or unknown patients, i.e. 10 months (95 % CI: 6.9, 17.4). Forty-six percent of patients had a decrease in bone marrow infiltration that exceeded 50 % and 58 % had a decrease in serum tryptase levels that exceeded 50 %. Spleen volume decreased by \geq 10 % in 68.9 % of patients with at least 1 post-baseline assessment (26.7 % of patients had a reduction of \geq 35 %, which

correlates with a 50 % decrease by palpation).

The median time to response was 0.3 months (range: 0.1 to 3.7 months). The median duration of follow-up was 43 months.

Table 7 Efficacy of midostaurin in ASM, SM-AHN and MCL: primary efficacy population

	All N=89	ASM N=16	SM-AHN N=57	MCL N=16
Primary endpoint				
Overall response, n (%)	53 (59.6)	12 (75.0)	33 (57.9)	8 (50.0)
(95% CI)	(48.6, 69.8)	(47.6, 92.7)	(44.1, 70.9)	(24.7, 75.3)
Major response, n (%)	40 (44.9)	10 (62.5)	23 (40.4)	7 (43.8)
Partial response, n (%)	13 (14.6)	2 (12.5)	10 (17.5)	1 (6.3)
Stable disease, n (%)	11 (12.4)	1 (6.3)	7 (12.3)	3 (18.8)
Progressive disease, n (%)	10 (11.2)	1 (6.3)	6 (10.5)	3 (18.8)
Secondary endpoints				
Median duration of response, months (95% CI)	18.6 (9.9, 34.7)	36.8 (5.5, NE)	10.7 (7.4, 22.8)	NR (3.6, NE)
Median overall survival, months (95% CI)	26.8 (17.6, 34.7)	51.1 (28.7, NE)	20.7 (16.3, 33.9)	9.4 (7.5, NE)
Kaplan-Meier estimates at 5 years (95% CI)	26.1 (14.6, 39.2)	34.8 (1.7, 76.2)	19.9 (8.6, 34.5)	33.7 (12.3, 56.8)

NE: Not Estimated, NR: Not Reached

Patients who received non-study anti-neoplastic therapy were considered as having progressed at the time of the new therapy.

Although the study was designed to be assessed with the modified Valent and Cheson criteria, as a *post-hoc* exploratory analysis, efficacy was also assessed per the 2013 International Working Group - Myeloproliferative Neoplasms Research and Treatment - European Competence Network on Mastocytosis (IWG-MRT-ECNM) consensus criteria. Response to

Sotauric was determined using a computational algorithm applied without any adjudication. Out of 116 patients, 113 had a C-finding as defined by IWG response criteria (excluding ascites as a C-finding). All responses were considered and required a 12-week confirmation (see Table 8).

Table 8 Efficacy of midostaurin in ASM, SM-AHN and MCL per IWG-MRT-ECNM consensus criteria using an algorithmic approach

	All patients evaluated	ASM	SM-AHN	MCL	Subtype unknown
	N=113	N=15	N=72	N=21	N=5
Overall response rate, n (%)	32 (28.3)	9 (60.0)	15 (20.8)	7 (33.3)	1 (20.0)
(95% CI)	(20.2, 37.6)	(32.3, 83.7)	(12.2, 32.0)	(14.6, 57.0)	(0.5, 71.6)
Best overall response, n (%)					
Complete remission	1 (0.9)	0	0	1 (4.8)	0
Partial remission	17 (15.0)	5 (33.3)	8 (11.1)	3 (14.3)	1 (20.0)
Clinical improvement	14 (12.4)	4 (26.7)	7 (9.7)	3 (14.3)	0
Duration of response*					
n/N (%)	11/32 (34.4)	4/9 (44.4)	4/15 (26.7)	3/7 (42.9)	0/1 (0.0)
median (95% CI)	NE (27.0, NE)	36.8 (10.3, 36.8)	NE (17.3, NE)	NE (4.1, NE)	NE
Overall survival					
n/N (%)	65/113 (57.5)	4/15 (26.7)	49/72 (68.1)	12/21 (57.1)	0/5 (0.0)
median (95% CI)	29.9 (20.3, 42.0)	51.1 (34.7, NE)	22.1 (16.8, 32.2)	22.6 (8.3, NE)	NE

*Confirmation period for responses: 12 weeks

Analysis excludes ascites as a C-finding.

Patients who received non-study anti-neoplastic therapy were considered as having progressed at the time of the new therapy.

The supportive study was a single-arm, multicentre, open-label phase II study of 26 patients with ASM, SM-AHN and MCL (CPKC412A2213). Midostaurin was administered orally at 100 mg twice daily in cycles of 28 days. Lack of a major response (MR) or partial response (PR) by the end of the second cycle required discontinuation from the study treatment. Twenty

(76.9 %) patients had ASM (17 [85 %] with AHN) and 6 patients (23.1 %) had MCL (2 [33.3 %] with AHN). The median age was 64.5 years with half of the patients \geq 65 years). At baseline, 88.5 % had >1 C finding and 69.2 % had received at least one prior anti-neoplastic regimen.

The primary endpoint was ORR evaluated by the Valent criteria during the first two cycles of treatment. Nineteen patients (73.1%; 95% CI = [52.2, 88.4]) achieved a response during the first two cycles of treatment (13 MR; 6 PR). The median duration of follow-up was 73 months, and the median duration of response has not been reached. Median overall survival was 40.0 months (patients were only followed up for one year after treatment discontinuation for survival).

5.2 Pharmacokinetic properties

Midostaurin is a compound with good absorption and poor solubility. Two of its metabolites demonstrated pharmacological activities (CGP52421 and CGP62221). Following multiple doses, the pharmacokinetics of midostaurin and CGP62221 were time-dependent, with an initial increase observed in the first week followed by a decline of concentrations until reaching steady state on day 28. CGP52421 concentrations do not appear to decline as significantly as for midostaurin and CGP62221.

Absorption

The absolute bioavailability of midostaurin following oral administration is not known.

In humans, the absorption of midostaurin was rapid after oral administration, with T_{max} of total radioactivity observed at 1-3 hours post dose. The population pharmacokinetic analysis indicated that the absorption in patients was less than dose proportional at doses $>$ 50 mg

twice daily.

In healthy subjects, after administration of a single dose of 50 mg midostaurin with food, AUC of midostaurin was increased to 20800 ng*h/ml and C_{max} was decreased to 963 ng/ml (see section 4.5). Similarly, for CGP52421 and CGP62221 AUC increased to 19000 and 29200 ng*h/ml and C_{max} decreased to 172 and 455 ng/ml, respectively. Time to peak concentration was also delayed in the presence of a high-fat meal. T_{max} was delayed for all entities, midostaurin median T_{max} was 3 h, and for CGP52421 and CGP62221 T_{max} was delayed to 6 and 7 hours respectively.

In clinical studies, the efficacy and safety of Sotauric were investigated following administration with a light meal. After oral administration of a single 100 mg dose of midostaurin under fed conditions in ASM, SM-AHN and MCL patients, AUC_{inf} , C_{max} and T_{max} were 49600 ng*h/ml, 2940 ng/ml and 3 h, respectively, for midostaurin. For CGP52421, AUC_{0-12h} and C_{max} were 2770 ng*h/ml and 299 ng/ml, respectively. AUC_{0-12h} and C_{max} for CGP62221 were 8700 ng*h/ml and 931 ng/ml, respectively. After 100 mg bid multiple oral doses of midostaurin the $C_{min,ss}$ plasma midostaurin in AML and ASM, SM-AHN, MCL patients were 919 and 1060 ng/ml, respectively. The CGP62221 $C_{min,ss}$ in the AML and the ASM, SM-AHN, MCL population were 1610 ng/ml and 2020 ng/ml, respectively. The CGP52421, $C_{min,ss}$ in the AML and the ASM, SM-AHN, MCL population were 8630 ng/ml and 2860 ng/ml, respectively.

Distribution

Midostaurin has a tissue distribution of geometric mean of 95.2 l (Vz/F). Midostaurin and its metabolites are distributed mainly in plasma rather than red blood cells. *In vitro* data showed midostaurin is more than 98 % bound to plasma proteins, such as albumin, α 1-acid glycoprotein (AGP) and lipoprotein.

Biotransformation

Midostaurin is metabolised by CYP3A4 mainly via oxidative pathways. The major plasma components included midostaurin and two major active metabolites, CGP62221 (via O-demethylation) and CGP52421 (via hydroxylation), accounting for $27.7\pm 2.7\%$ and $38.0\pm 6.6\%$, respectively, of the total plasma exposure at 96 hours after a single 50 mg dose of midostaurin.

Elimination

The median terminal half-lives of midostaurin, CGP62221 and CGP52421 in plasma are approximately 20.9, 32.3 and 471 hours. The mean apparent plasma clearance (CL/F) was 2.4-3.1 l/h in healthy subjects. In AML and ASM, SM-AHN and MCL patients, population pharmacokinetic estimates for clearance of midostaurin at steady state were 5.9 l/h and 4.4 l/h, respectively. The Human Mass Balance study results indicated that faecal excretion is the major route of excretion (78 % of the dose), and mostly as metabolites (73 % of the dose), while unchanged midostaurin accounts for 3 % of the dose. Only 4 % of the dose is recovered in urine.

Linearity/non-linearity

In general, midostaurin and its metabolites showed no major deviation from dose-proportionality after a single dose in the range of 25 mg to 100 mg. However, there was a less than dose-proportional increase in exposure after multiple doses within the dose range of 50 mg to 225 mg daily.

Following multiple oral doses, midostaurin displayed time-dependent pharmacokinetics with an initial increase in plasma concentrations during the first week (peak C_{min}) followed by a decline with time to a steady-state after approximately 28 days (2.5-fold decrease). While the exact mechanism for the declining concentration of midostaurin is unclear, it is likely due to the auto-induction properties of midostaurin and its two active metabolites CGP52421 and CGP62221 on CYP3A4. The pharmacokinetics of the CGP62221 metabolite showed a similar trend. However, CGP52421 concentrations increased up to 2.5-fold for ASM, SM-AHN and MCL and up to 9-fold for AML, compared to midostaurin after one month of treatment.

In vitro evaluation of drug-drug interaction potential

Enzyme drug-drug interactions

Cytochrome P450 inhibition

Based on *in vitro* data, midostaurin and its active metabolites, CGP52421 and CGP62221, are considered inhibitors and may potentially cause increases in exposure of co-administered medicines primarily cleared by CYP1A2, CYP2D6, CYP2C8, CYP2C9, CYP2E1 and CYP3A4/5. In addition, a time-dependent inhibition of CYP3A4 by midostaurin, CGP52421 and CGP62221 was also observed *in vitro*.

Cytochrome P450 induction

Based on *in vitro* data, midostaurin and its active metabolites, CGP52421 and CGP62221, are also considered inducers and may cause decreases in exposure of co-administered medicines primarily cleared by CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19 and CYP3A4/5 (see section 4.5).

Transporter drug-drug interactions

In vitro experiments demonstrated that midostaurin, CGP52421 and CPG62221 can potentially inhibit P-gp, BCRP, OATP1B1 and BSEP.

Special populations

Elderly patients

Based on population pharmacokinetic analyses no significant impact of age on the pharmacokinetics of midostaurin and its two active metabolites was identified for patients aged between 65 and 85 years. In adult patients with ASM, SM-AHN and MCL or AML, no midostaurin dose adjustment is required based on age.

Paediatric patients

Sotauric is not recommended to be used in children and adolescents (see section 4.2). The pharmacokinetics of midostaurin in paediatric patients were explored in a phase I dose escalation monotherapy study with 22 patients (12 aged 0-2 years and 10 aged 10-17 years) with AML or MLL-rearranged ALL using a population pharmacokinetic approach. The pharmacokinetics of midostaurin were less than dose proportional with the doses of 30 mg/m² and 60 mg/m² after single and multiple doses. Due to the limited pharmacokinetic data in paediatric patients, no comparison with midostaurin pharmacokinetics in adults can be made.

Gender

Based on population pharmacokinetic model analyses of the effect of gender on clearance of midostaurin and its active metabolites, there was no statistically significant finding and the anticipated changes in exposure (<20 %) were not deemed to be clinically relevant. No

midostaurin dose adjustment is required based on gender.

Hepatic impairment

A dedicated hepatic impairment study assessed the systemic exposure of midostaurin after oral administration of 50 mg twice daily for 6 days in subjects with baseline mild or moderate hepatic impairment (Child-Pugh Class A or B, respectively) and control subjects with normal hepatic function. The maximum concentration was reached between 2 and 3 hours after administration after single or repeated doses for all groups. On day 1, the AUC_{0-12} and C_{max} were 8130 ng*h/ml and 1206 ng/ml, respectively, for healthy subjects. AUC_{0-12} was decreased by 39 % and 36 % in subjects with mild and moderate hepatic impairment, respectively. On day 7, $AUC_{C_{trough}}$ (exposure under the curve of C_{trough} from day 1 to day 7) was 5410 ng*h/ml in healthy subjects and was decreased by 35 % and 20 % in subjects with mild and moderate hepatic impairment, respectively. AUC_{tau} was decreased by 28 % and 20 % on day 7, respectively. Finally, the long-term data from patients were analysed using a population pharmacokinetic approach. No impact of hepatic impairment could be identified in patients with mild or moderate hepatic impairment in the ASM, SM-AHN, MCL and AML populations.

Overall, there was no clinically relevant increase in exposure (AUC) to plasma midostaurin in subjects with mild or moderate hepatic impairment compared to subjects with normal hepatic function. No dosage adjustment is necessary for patients with baseline mild or moderate hepatic impairment. The pharmacokinetics of midostaurin have not been assessed in patients with baseline severe hepatic impairment (Child-Pugh Class C) (see section 4.4).

Renal impairment

Renal elimination is a minor route of elimination for midostaurin. No dedicated renal impairment study was conducted for midostaurin. Population pharmacokinetic analyses were conducted

using data from clinical studies in patients with AML (n=180) and ASM, SM-AHN and MCL (n=141). Out of the 321 patients included, 177 patients showed pre-existing mild (n=113), moderate (n=60) or severe (n=4) renal impairment ($15 \text{ ml/min} \leq \text{creatinine clearance [CrCL]} < 90 \text{ ml/min}$). 144 patients showed normal renal function ($\text{CrCL} > 90 \text{ ml/min}$) at baseline. Based on the population pharmacokinetic analyses, midostaurin clearance was not significantly impacted by renal impairment and therefore no dosage adjustment is necessary for patients with mild or moderate renal impairment.

5.3 Preclinical safety data

Due to dose-limiting toxicity, clinical therapeutic exposure levels could not be reached in animals. All animal findings described below were observed at midostaurin exposure significantly lower than therapeutic levels.

Safety pharmacology and single/repeat dose toxicity

Safety pharmacology studies indicate that midostaurin is unlikely to interfere with vital functions of the central nervous system. *In vitro*, midostaurin did not inhibit hERG channel activity up to the limit of solubility of $12 \mu\text{M}$. The two major human metabolites GGP52421 and CGP62221 (also tested at the limit of solubility) inhibited hERG current with moderate safety margins. In the repeat-dose studies in dogs, a decrease in heart rate, prolongation of the P-Q interval, and sporadically occurring atrioventricular blocks were seen in individual animals.

In the repeat-dose studies, target organs for toxicity were the gastrointestinal tract (emesis in dogs and monkeys, diarrhoea and mucosal alteration), testes (decreased spermatogenesis), bone marrow (hypocellularity) and lymphoid organs (depletion/atrophy). The effect on the bone

marrow and lymphoid organs was accompanied by haematological changes of decreased white blood cells, lymphocytes and erythrocytic parameters. An increase in liver enzymes (ALT and AST) was seen consistently in rats, and in dogs and monkeys in long-term studies of \geq 3 months duration, without histopathological correlates.

Reproductive toxicity

In a fertility study in rats, midostaurin was associated with reduced fertility, testicular degeneration and atrophy, reduced sperm motility, oligo- and aspermia, increased resorptions, decreased pregnancy rate, number of implants and live embryos.

In embryo-foetal development studies in rats and rabbits, increased numbers of late resorptions, reduced foetal weight and reduced skeletal ossification were observed.

In a pre- and post-natal developmental study, maternal dystocia and reduced litter size, lower pup body weights, accelerated complete eye opening and delayed auricular startle ontogeny were noted.

Juvenile animal studies

In a toxicity study in juvenile rats, midostaurin was administered from days 7 to 70 postpartum. A reduction in body weight, haemorrhage and mixed cell infiltration in the lungs, and erythrocytosis/erythrophagocytosis in the mesenteric lymph nodes were seen. There were no effects on physical development, sensory function or behavioural function. Mating index, fertility index and conception rates were reduced at 0, 5 and 15 mg/kg/day, but not at 2 mg/kg/day.

Genotoxicity

In vitro and *in vivo* genotoxicity studies covering relevant genotoxicity endpoints showed no evidence of mutagenic or clastogenic activity. No carcinogenicity studies have been performed.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content

Macrogolglycerol hydroxystearate

Macrogol

Ethanol anhydrous

Maize oil mono-di-triglycerides

All-rac-alpha-tocopherol

Capsule shell

Gelatin

Glycerol

Titanium dioxide (E171)

Iron oxide yellow (E172)

Iron oxide red (E172)

Purified water

Printing ink

Carmine (E120)

Hypromellose

Propylene glycol

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Store at or below 30°C.

Store in the original container in order to protect from moisture.

6.5 Nature and contents of container

Available in packs of 28, 56 and 112 soft capsules per box. Sotauric 25 mg soft capsules are packed in PA/Al/PVC-Al blister packs.

Not all pack sizes and types may be marketed.

6.6 Special precautions for disposal

Any unused medicines or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Novartis South Africa (Pty) Ltd

Magwa Crescent West

Waterfall City, Jukskei View

Johannesburg

2090

South Africa

8. MARKETING AUTHORISATION NUMBER(S)

Sotauric 25 mg soft capsule: 52/32.16/0121

9. DATE OF FIRST AUTHORISATION

January 2020

10. DATE OF REVISION OF THE TEXT

January 2020