

## PROFESSIONAL INFORMATION

### SCHEDULING STATUS

S4

#### 1 NAME OF THE MEDICINE

**Perjeta®** Concentrate solution for infusion

#### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Perjeta contains pertuzumab as the active substance.

Each vial of Perjeta contains 420 mg pertuzumab in 14 mL preservative free concentrate, (equivalent to 30 mg/mL), concentrated solution for infusion.

Contains sugar (sucrose).

For the full list of excipients, see section 6.1.

#### 3 PHARMACEUTICAL FORM

Perjeta is a sterile almost clear to slightly opalescent, colourless to pale yellow liquid which contains no preservatives.

#### 4 CLINICAL PARTICULARS

##### 4.1 Therapeutic Indications

###### **Metastatic Breast Cancer:**

Perjeta is indicated for use in combination with trastuzumab and docetaxel for patients with HER2-positive metastatic or locally recurrent unresectable breast cancer, who have not received previous anti-HER2 therapy or chemotherapy for their metastatic disease.

###### **Early Breast Cancer:**

Perjeta is indicated in combination with trastuzumab and chemotherapy for the:

- neoadjuvant treatment of patients with HER2 positive, locally advanced, inflammatory, or early stage breast cancer (either > 2 cm in diameter or node positive) as part of a complete regimen for early breast cancer.
- adjuvant treatment of patients with HER-2 positive breast cancer at high risk of recurrence.

## 4.2 Posology and method of administration

Perjeta should only be initiated under the supervision of a medical practitioner experienced in the administration of anti-cancer medicines. Perjeta should be administered by a medical practitioner prepared to manage anaphylaxis and in an environment where full resuscitation service is immediately available.

Patients treated with Perjeta must have HER2-positive tumour status, defined as a score of 3+ by immunohistochemistry (IHC) and/or a ratio of  $\geq 2,0$  by *in situ* hybridisation (ISH) assessed by a validated test.

To ensure accurate and reproducible results, the testing must be performed in a specialised laboratory, which can ensure validation of the testing procedures.

Perjeta must be diluted by a healthcare professional and administered as an intravenous infusion. Do not administer as an intravenous push or bolus.

### ***Dosage of Perjeta in combination with trastuzumab and docetaxel***

#### **Metastatic and Early Breast Cancer**

The recommended initial dose of Perjeta is 840 mg administered as a 60 minutes intravenous infusion, followed every 3 weeks thereafter by a maintenance dose of 420 mg administered over a period 30 to 60 minutes.

An observation period of 30 - 60 minutes is recommended after completion of each Perjeta infusion. The observation period should be completed prior to any subsequent dose of trastuzumab or chemotherapy (see section 4.4).

Perjeta and trastuzumab should be administered sequentially and can be given in any order. When administered with Perjeta, the recommendation is to follow a 3-weekly schedule for trastuzumab administered either as:

- an IV infusion with an initial dose of 8 mg/kg followed every 3 weeks thereafter by a dose of 6 mg/kg body weight
- or
- a fixed dose of trastuzumab subcutaneous (SC) injection (600 mg) for the initial dose and every 3 weeks thereafter irrespective of the patient's body weight

In patients receiving a taxane, Perjeta and trastuzumab should be administered prior to the taxane.

When administered with Perjeta, the recommended initial dose of docetaxel is 75 mg/m<sup>2</sup>.

In patients receiving an anthracycline-based regimen, Perjeta and trastuzumab should be administered following completion of the entire anthracycline regimen.

### ***Duration of treatment***

#### **Metastatic Breast Cancer (MBC)**

It is recommended that patients are treated with Perjeta until disease progression or unmanageable toxicity.

#### **Early Breast Cancer (EBC)**

In the neoadjuvant setting (before surgery), it is recommended that patients are treated with Perjeta for 3 to 6 cycles depending on the regimen chosen, in combination with trastuzumab and chemotherapy.

In the adjuvant setting (after surgery), Perjeta should be administered in combination with trastuzumab for a total of one year (maximum 18 cycles or until disease recurrence, or unmanageable toxicity, whichever occurs first), as part of a complete regimen for early breast cancer, including standard anthracycline- and/or taxane-based chemotherapy. Perjeta and trastuzumab should start on Day 1 of the first taxane-containing cycle and should continue even if chemotherapy is discontinued.

Patients who start Perjeta and trastuzumab in the neoadjuvant setting should continue to receive adjuvant Perjeta and trastuzumab to complete 1 year of treatment (maximum 18 cycles).

### ***Delayed or Missed doses***

For recommendations on delayed or missed doses, please refer to Table 1 below.

**Table 1 Recommendations regarding delayed or missed doses**

| Time between two sequential doses | Perjeta                                                                                                                                                                                                       | Trastuzumab                                                                                                                                                                                                                  |                                                                                                                              |
|-----------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------|
|                                   |                                                                                                                                                                                                               | IV                                                                                                                                                                                                                           | SC                                                                                                                           |
| < 6 weeks                         | The 420 mg dose of Perjeta IV should be administered as soon as possible. Do not wait until the next planned dose.                                                                                            | The 6 mg/kg dose of trastuzumab IV should be administered as soon as possible. Do not wait until the next planned dose.                                                                                                      | The fixed dose of 600 mg trastuzumab SC should be administered as soon as possible. Do not wait until the next planned dose. |
| ≥ 6 weeks                         | The loading dose of 840 mg Perjeta IV should be re-administered as a 60 minute infusion, followed by a maintenance dose of 420 mg IV administered over a period of 30 to 60 minutes every 3 weeks thereafter. | The loading dose of 8 mg/kg of trastuzumab IV should be re-administered over approximately 90 minutes, followed by a maintenance dose of 6 mg/kg IV administered over a period of 30 or 90 minutes every 3 weeks thereafter. |                                                                                                                              |

### ***Dose modifications***

Dose reductions are not recommended for Perjeta.

Patients may continue therapy during periods of reversible chemotherapy-induced myelosuppression but they should be monitored carefully for complications of neutropenia during this time. For docetaxel and other chemotherapy dose modifications, see relevant professional informations.

For trastuzumab, dose reductions are not recommended, see trastuzumab professional information. If trastuzumab treatment is discontinued, treatment with Perjeta should be discontinued. If docetaxel is discontinued, treatment with Perjeta and trastuzumab may continue until disease progression or unmanageable toxicity in the metastatic setting.

#### ***Left ventricular dysfunction:***

See section 4.4, Table 2 for information on dose recommendations in the event of left ventricular dysfunction.

#### ***Infusion-related reactions:***

The infusion rate may be slowed or interrupted if the patient develops an infusion-related reaction (see sections 4.4 and 4.8). The infusion may be resumed when symptoms abate. Treatment including oxygen, beta agonists, antihistamines, rapid i.v. fluids and antipyretics may also help alleviate symptoms.

#### ***Hypersensitivity reactions/anaphylaxis:***

The infusion should be discontinued immediately and permanently if the patient experiences a serious hypersensitivity reaction (e.g. anaphylaxis), bronchospasm or acute respiratory distress syndrome (see section 4.4).

### **Special Dosage Instructions**

***Elderly patients:*** No overall differences in efficacy of Perjeta were observed in patients  $\geq 65$  and  $< 65$  years of age. The incidence of the following all grade adverse events was at least 5 % higher in patients aged  $\geq 65$  years of age, compared to patients aged  $< 65$  years of age: decreased appetite,

anaemia, decreased weight, asthenia, dysgeusia, peripheral neuropathy, hypomagnesemia and diarrhoea.

No dose adjustment is necessary in the elderly population  $\geq 65$  years of age.

*Patients with renal impairment:* Dose adjustments of Perjeta are not needed in patients with mild or moderate renal impairment. No dose recommendations can be made for patients with severe renal impairment because of the limited pharmacokinetic data available (see section 5.2 Pharmacokinetic properties).

*Patients with hepatic impairment:* The safety and efficacy of Perjeta have not been studied in patients with hepatic impairment. No specific dose recommendations can be made.

*Paediatric population:* The safety and efficacy of Perjeta in children and adolescents below 18 years of age have not been established. There is no relevant use of Perjeta in the paediatric population in the indication of breast cancer.

### **Method of administration**

Perjeta is administered intravenously by infusion. It should not be administered as an intravenous push or bolus. For instructions on dilution of Perjeta prior to administration, see below. For the initial dose, the recommended infusion period is 60 minutes. If the first infusion is well tolerated, subsequent infusions may be administered over a period of 30 minutes to 60 minutes (see section 4.4).

**Instructions for dilution:** see **Special Instructions for use, Handling and Disposal:** see section 6.6

**Incompatibilities:** see section 6.2

## **4.3 Contraindications**

Perjeta is contraindicated in patients with known hypersensitivity to pertuzumab or any of its excipients.

Pregnancy and Lactation (see section 4.6).

#### 4.4 Special warnings and precautions for use

In order to improve traceability of biological medicines, the trade-name of the administered product should be clearly recorded (or stated) in the patient file.

##### *Infusion-related reactions*

Perjeta has been associated with infusion-related reactions, including events with fatal outcomes (see section 4.8). Close observation of the patient during and for 60 minutes after the first infusion, and during and for 30 minutes following subsequent infusions of Perjeta is recommended. If a significant infusion-related reaction occurs, the infusion should be slowed down or interrupted and appropriate medical therapies should be administered. Patients should be evaluated and carefully monitored until complete resolution of signs and symptoms. Permanent discontinuation should be considered in patients with severe infusion reactions. This clinical assessment should be based on the severity of the preceding reaction and response to administered treatment for the adverse reaction (see section 4.2).

##### *Hypersensitivity reactions/anaphylaxis*

Patients should be observed closely for hypersensitivity reactions. Severe hypersensitivity reactions, including anaphylaxis and events with fatal outcomes, have been observed in patients treated with Perjeta (see section 4.8). Medications to treat such reactions, as well as emergency equipment, should be available for immediate use. Perjeta is contraindicated in patients with known hypersensitivity to pertuzumab or to any of its excipients (see section 4.3).

##### *Left ventricular dysfunction (including congestive heart failure)*

Decreases in LVEF have been reported with medicines that block HER2 activity, including Perjeta. The incidence of symptomatic left ventricular systolic dysfunction (LVD [congestive heart failure]) was higher in patients treated with Perjeta in combination with trastuzumab and chemotherapy compared to trastuzumab and chemotherapy. Patients who have received prior anthracyclines or prior radiotherapy to the chest area may be at higher risk of decreased LVEF. The majority of cases of symptomatic heart failure reported in the adjuvant setting were in patients who received anthracycline-based chemotherapy (see section 4.8).

Perjeta has not been studied in patients with: a pre-treatment LVEF value of  $\leq 50\%$ ; a prior history of congestive heart failure (CHF); decreases in LVEF to  $< 50\%$  during prior trastuzumab adjuvant therapy; or conditions that could impair left ventricular function such as uncontrolled hypertension, recent myocardial infarction, serious cardiac dysrhythmia requiring treatment or a cumulative prior anthracycline exposure to  $> 360 \text{ mg/m}^2$  of doxorubicin or its equivalent.

Assess LVEF prior to initiation of Perjeta and at regular intervals during treatment to ensure that LVEF is within normal limits (see Table 2 below). If the LVEF declines as indicated in Table 2 and has not improved, or has declined further at the subsequent assessment, discontinuation of Perjeta and trastuzumab should be strongly considered.

**Table 2 Dose recommendations for left ventricular dysfunction**

|                                 | <b>Pre-treatment LVEF:</b> | <b>Monitor LVEF every:</b> | <b>Withhold Perjeta and trastuzumab for at least 3 weeks for an LVEF decrease to:</b> |                                                                           | <b>Resume Perjeta and trastuzumab after 3 weeks if LVEF has recovered to:</b> |                                                                       |
|---------------------------------|----------------------------|----------------------------|---------------------------------------------------------------------------------------|---------------------------------------------------------------------------|-------------------------------------------------------------------------------|-----------------------------------------------------------------------|
| <b>Metastatic Breast Cancer</b> | $\geq 50\%$                | ~12 weeks                  | Either                                                                                |                                                                           | Either                                                                        |                                                                       |
|                                 |                            |                            | $<40\%$                                                                               | 40 %-45 %<br>with a fall of $\geq 10\%$ -points below pre-treatment value | $>45\%$                                                                       | 40 %-45 %<br>with a fall of $<10\%$ -points below pre-treatment value |
|                                 | $\geq 55\%^*$              |                            |                                                                                       |                                                                           | Either                                                                        |                                                                       |

|                            |  |                                                      |                                                                   |        |                                                       |
|----------------------------|--|------------------------------------------------------|-------------------------------------------------------------------|--------|-------------------------------------------------------|
| <b>Early Breast Cancer</b> |  | ~12 weeks<br>(once during<br>neoadjuvant<br>therapy) | <50 % with a fall of<br>≥10 %-points below<br>pre-treatment value | ≥ 50 % | < 10 %-<br>points<br>below pre-<br>treatment<br>value |
|----------------------------|--|------------------------------------------------------|-------------------------------------------------------------------|--------|-------------------------------------------------------|

\*for patients receiving anthracycline-based chemotherapy, a LVEF of ≥ 50 % is required after completion of anthracyclines, before starting Perjeta and trastuzumab

**Febrile neutropenia:** Patients treated with Perjeta, trastuzumab and docetaxel are at increased risk of febrile neutropenia especially during the first 3 cycles of treatment. The higher incidence of febrile neutropenia may be associated with the higher incidence of mucositis and diarrhoea in these patients. Symptomatic treatment for mucositis and diarrhoea should be considered.

**Sugars:** Contains sucrose which may have an effect on the glycaemic control of patients with diabetes mellitus. Patients with rare hereditary conditions such as fructose intolerance, glucose-galactose mal-absorption or sucrase-isomaltase insufficiency should not take Perjeta.

#### 4.5 Interaction with other medicines and other forms of interaction

No pharmacokinetic (PK) interactions were observed between Perjeta and trastuzumab, or between Perjeta and docetaxel. In the population PK analysis, no evidence of a medicine interaction has been shown between Perjeta and trastuzumab and between Perjeta and docetaxel. This lack of interaction was confirmed by pharmacokinetic data from the additional neoadjuvant and early breast cancer studies.

Five studies have evaluated the effects of Perjeta on the pharmacokinetics of co-administered cytotoxic agents, docetaxel, paclitaxel, gemcitabine, carboplatin, erlotinib and capecitabine, respectively. There was no evidence of any pharmacokinetics interaction between Perjeta and any of these agents. The pharmacokinetics of Perjeta in these studies were comparable to those observed in single-agent studies.

#### 4.6 Fertility, pregnancy and lactation

### *Pregnancy*

Perjeta should not be used during pregnancy (see section 4.3). Women of child bearing potential and female partners of male patients of child bearing potential should use effective contraception while receiving Perjeta and for 6 months following the last dose of Perjeta.

Combined hormonal and barrier methods are recommended.

In animal studies, Perjeta administered to cynomolgus monkeys during organogenesis led to oligohydramnios, delayed renal development and embryo foetal death.

### *Breastfeeding*

Because human IgG is secreted in human milk, and animal data that indicate Perjeta is foetotoxic, women receiving Perjeta must not breastfeed their infants.

## **4.7 Effects on ability to drive and use machines**

Patients experiencing headache, dizziness or infusion reactions should be advised not to drive and use machines until symptoms abate.

## **4.8 Undesirable effects**

### **a. Summary of the safety profile:**

#### **Clinical Trials**

The safety of Perjeta has been evaluated in more than 6 000 patients in trials conducted in patients with various malignancies and predominantly treated with Perjeta in combination with other anti-neoplastic medicines.

The safety of Perjeta was generally consistent across studies, although the incidence and most common adverse drug reactions (ADRs) varied depending on whether Perjeta was administered as monotherapy or in combination with other anti-neoplastic medicines.

#### **Metastatic and Early Breast Cancer**

Table 3 summarises the ADRs from the pivotal clinical trials, in which Perjeta was given:

- in combination with docetaxel and trastuzumab to patients with metastatic breast cancer (n=453)

- from the neoadjuvant trials, in which Perjeta was given in combination with trastuzumab and chemotherapy to patients with locally advanced, inflammatory or early breast cancer (n=309 and n=218)
- in which adjuvant Perjeta was given in combination with trastuzumab and anthracycline-based or non-anthracycline based, taxane-containing chemotherapy to patients with EBC (n=2 364).

As Perjeta is used with trastuzumab and chemotherapy, it is difficult to ascertain the causal relationship of an adverse reaction to a particular medicine.

#### b. Tabulated list of adverse reactions

The ADRs are listed below by system organ class (SOC) and categories of frequency:

Very common ( $\geq 1/10$ ), Common ( $\geq 1/100$  to  $< 1/10$ ), Uncommon ( $\geq 1/1\ 000$  to  $< 1/100$ ), Rare ( $\geq 1/10\ 000$  to  $< 1/1\ 000$ ), Very rare ( $< 1/10\ 000$ ).

Within each frequency grouping and SOC, adverse reactions are presented in the order of decreasing seriousness.

The most common ADRs ( $\geq 30\%$ ) from this pooled data were diarrhoea, alopecia, nausea, fatigue, neutropenia, and vomiting. The most common NCI-CTCAE Grade 3-4 ADRs ( $\geq 10\%$ ) were neutropenia and febrile neutropenia.

**Table 3: Summary of ADRs in patients treated with Perjeta in the Metastatic and Neoadjuvant setting**

| System<br>Organ Class                      | Very Common                                             | Common     | Uncommon |
|--------------------------------------------|---------------------------------------------------------|------------|----------|
| Infections and infestations                | Upper respiratory tract<br>infection<br>Nasopharyngitis | Paronychia |          |
| Blood and lymphatic<br>system<br>disorders | Febrile neutropenia*<br>Neutropenia<br>Leucopenia       |            |          |

|                                                 |                                                                                                                                     |                                   |                               |
|-------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------|-------------------------------|
|                                                 | Anaemia                                                                                                                             |                                   |                               |
| Immune system disorders                         | Hypersensitivity/<br>anaphylactic reaction°<br><br>Infusion-related<br>reaction, Cytokine<br>release syndrome°°                     |                                   |                               |
| Metabolism and nutrition disorders              | Decreased appetite                                                                                                                  |                                   |                               |
| Psychiatric disorders                           | Insomnia                                                                                                                            |                                   |                               |
| Nervous system disorders                        | Peripheral neuropathy<br><br>Headache<br><br>Dysgeusia<br><br>Peripheral sensory<br>neuropathy<br><br>Dizziness<br><br>Paraesthesia |                                   |                               |
| Eye disorders                                   | Increased lacrimation                                                                                                               |                                   |                               |
| Cardiac disorders                               |                                                                                                                                     | Left ventricular<br>dysfunction** | Congestive heart<br>failure** |
| Vascular disorders                              | Hot flush                                                                                                                           |                                   |                               |
| Respiratory, thoracic and mediastinal disorders | Cough<br><br>Epistaxis<br><br>Dyspnoea                                                                                              | Pleural effusion                  | Interstitial lung<br>disease  |
| Gastrointestinal disorders                      | Diarrhoea<br><br>Vomiting<br><br>Stomatitis<br><br>Nausea                                                                           |                                   |                               |

|                                                      |                                                                                                |                  |  |
|------------------------------------------------------|------------------------------------------------------------------------------------------------|------------------|--|
|                                                      | Constipation<br>Dyspepsia<br>Abdominal pain                                                    |                  |  |
| Skin and subcutaneous tissue disorders               | Alopecia<br>Rash<br>Nail disorder<br>Pruritis<br>Dry skin                                      |                  |  |
| Musculoskeletal and connective tissue disorders      | Myalgia<br>Arthralgia<br>Pain in extremity                                                     |                  |  |
| General disorders and administration site conditions | Mucositis/mucosal inflammation<br>Pain<br>Peripheral oedema<br>Pyrexia<br>Fatigue<br>Asthaenia | Chills<br>Oedema |  |

\* Including adverse reactions with a fatal outcome.

\*\* For the overall treatment period across the 4 studies.

° Hypersensitivity/anaphylactic reaction is based on a group of terms.

°° Infusion related reaction/cytokine release syndrome includes a range of different terms within a time window, see “Description of selected adverse reactions” below.

### c. Description of selected adverse events

***ADRs reported in patients receiving Perjeta and trastuzumab after discontinuation of docetaxel***

ADRs were reported less frequently after discontinuation of docetaxel treatment. After discontinuation of docetaxel, all ADRs in the Perjeta and trastuzumab treated group occurred in < 10 % of patients with the exception of diarrhoea (19,1 %), upper respiratory tract infection (12,8 %), rash (11,7 %), headache (11,4 %) and fatigue (11,1 %).

### ***Further information on selected adverse reactions***

#### ***Left ventricular dysfunction (LVD)***

In metastatic breast cancer pivotal trial, the incidence of LVD during study treatment was higher in the placebo-treated group than in the Perjeta-treated group (8,6 % and 6,6 %, respectively). The incidence of symptomatic LVD was also lower in the Perjeta-treated group (1,8 % in the placebo-treated group vs. 1,5 % in the Perjeta-treated group) (see section 4.4).

In a neoadjuvant trial, in which patients received 4 cycles of Perjeta as neoadjuvant treatment, the incidence of LVD (during the overall treatment period) was 7,5 % in the Perjeta, trastuzumab and docetaxel-treated group.

In a second neoadjuvant trial, the group treated with Perjeta plus trastuzumab and FEC had an incidence of LVD of 8,3 %.

In the neoadjuvant period of the early breast cancer trial, the incidence of asymptomatic LVD was 7 % in the group treated with dose dense AC followed by Perjeta plus trastuzumab and paclitaxel and 3,5 % in the group treated with FEC followed by Perjeta plus trastuzumab and docetaxel.

In the early breast cancer trial, the incidence of symptomatic heart failure (NYHA class III or IV) with a LVEF decline of at least 10 %-points from baseline and to < 50 % was < 1 %.

#### ***Infusion-related reactions***

An infusion-related reaction was defined in pivotal trials as any event reported as hypersensitivity, anaphylactic reaction, acute infusion reaction or cytokine release syndrome occurring during an infusion or on the same day as the infusion. When only Perjeta was administered, the overall frequency of infusion reactions was 13,2 %. The most common infusion reactions (> 1,0 %) were pyrexia, chills, fatigue, headache, asthenia, hypersensitivity and vomiting. During the second cycle

when all medicines were administered on the same day, the most common infusion related reactions (> 1,0 %) were fatigue, dysgeusia, hypersensitivity, myalgia and vomiting.

In the neoadjuvant and adjuvant trials, Perjeta was administered on the same day as the other study treatments. Infusion-related reactions occurred in 18,6 % - 25,0 % of patients on the first day of Perjeta administration (in combination with trastuzumab and chemotherapy). The type and severity of events were consistent with those observed in the MBC trial, with a majority of reactions being mild or moderate.

#### *Hypersensitivity reactions/anaphylaxis*

In the MBC pivotal trial, the overall frequency of hypersensitivity/anaphylaxis events (not including acute infusion reactions/cytokine release syndrome) during the treatment period was 11,3 %, of which 2 % were NCI-CTCAE Grade 3-4, respectively. Overall, 4 patients experienced events described as anaphylaxis (see section 4.4).

In the neoadjuvant and EBC trials, the overall frequency of hypersensitivity/anaphylaxis was highest in the Perjeta group (13,2 %), of which 2,6 % were NCI-CTCAE grade 3-4.

#### *Febrile neutropenia*

The majority of patients experienced at least one leucopenic event, 63,0 % of patients, of which the majority were neutropenic events. Febrile neutropenia occurred in 13,7 %. The proportion of patients experiencing febrile neutropenia was highest in the first cycle of therapy and declined steadily thereafter. An increased incidence of febrile neutropenia was observed for Asian patients compared with patients of other races and from other geographic regions. Among Asian patients, the incidence of febrile neutropenia was 26 %.

#### *Diarrhoea*

In metastatic breast cancer, diarrhoea occurred in 68,8 % of patients. Most events were mild-moderate in severity and occurred in the first few cycles of treatment. The incidence of NCI-CTCAE Grade 3-4 diarrhoea was 9,3 %. The median duration of the longest episode was 18 days.

#### *Rash*

Rash occurred in 45,2 % of patients. Most events of Grade 1 or 2 in severity, occurred in the first two cycles. Rash occurred in 40,2 % of patients treated with neoadjuvant Perjeta, trastuzumab and docetaxel compared with 29,0 % of patients treated with trastuzumab and docetaxel.

#### *Laboratory abnormalities*

In the pivotal trials, the incidence of NCI-CTCAE Grade 3-4 decreases in neutrophil counts were balanced in the Perjeta treated groups.

### **Post Marketing**

The following adverse drug reaction has been identified from post marketing experience with Perjeta:  
Metabolism and nutrition disorders: Tumour lysis syndrome.

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions to SAHPRA via the “6.04 Adverse Drug Reaction Report Form”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/Publications/Index/8>

### **4.9 Overdose**

In case of overdose, patients must be closely monitored for signs or symptoms of adverse reactions and appropriate symptomatic treatment initiated.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antineoplastic agents, monoclonal antibodies, ATC code: L01XC13  
Pertuzumab is a recombinant humanised monoclonal antibody that specifically targets the extracellular dimerisation domain (Subdomain II) of the human epidermal growth factor receptor 2 protein (HER2) and thereby blocks ligand-dependent heterodimerisation of HER2 with other HER family members, including EGFR, HER3 and HER4. As a result, pertuzumab inhibits ligand-initiated

intracellular signalling through two major signal pathways, mitogen-activated protein (MAP) kinase and phosphoinositide 3-kinase (PI3K). Inhibition of these signalling pathways can result in cell growth arrest and apoptosis, respectively. In addition, pertuzumab mediates antibody-dependent cell-mediated cytotoxicity (ADCC).

While pertuzumab alone inhibited the proliferation of human tumour cells, the combination of pertuzumab and trastuzumab significantly augmented anti-tumour activity in HER2-overexpressing xenograft models.

#### *Immunogenicity*

Anti-therapeutic antibodies (ATA) neutralising to pertuzumab were found in 3,3 % of pertuzumab-treated patients tested (13/389 patients) during the metastatic breast cancer trials and 4,1 % (16/392 patients) in the early breast cancer trials. None of these patients experienced anaphylactic/hypersensitivity reactions that were clearly related to ADA.

## **5.2 Pharmacokinetic Properties**

Across multiple clinical trials in various indications there was no change in clearance of pertuzumab at doses of 2 - 25 mg/kg. Based on a population pharmacokinetics (PK) analysis that included 481 patients, the median clearance (CL) of pertuzumab was 0,235 L/day and the median half-life was 18 days.

Pertuzumab displayed linear pharmacokinetics within the recommended dose range.

Baseline albumin and lean body weight were the most significant covariates influencing CL. Clearance decreased in patients with higher baseline albumin concentrations and increased in patients with greater lean body weight. There is no need to adjust the dosage of pertuzumab based on these covariates.

No differences in pertuzumab PK were observed in patients with early breast cancer compared to patients with metastatic breast cancer.

*Absorption:* Pertuzumab is administered as an IV infusion. There have been no studies performed with other routes of administration.

*Distribution:* Across all clinical studies, the volume of distribution of the central (V<sub>c</sub>) and the peripheral (V<sub>p</sub>) compartment in the typical patient was 3,11 L and 2,46 L respectively.

*Metabolism/Biotransformation:* The metabolism of pertuzumab has not been studied. Antibodies are cleared principally by catabolism.

*Elimination:* The median clearance (CL) of pertuzumab is approximately 0,235 L/day and the median half-life was 18 days.

#### *Pharmacokinetics in Special Populations*

*Elderly:* Pertuzumab has not been studied in elderly patients. In a population PK analysis, age was not found to significantly affect PK of pertuzumab. In the population PK analysis, 32,5 % (N=143) patients were ≥ 65 years of age and 9,1 % (N=40) patients were ≥ 75 years of age.

*Patients with renal impairment:* No dedicated renal impairment trial for Perjeta has been conducted. Based on the results of the population PK analysis, Perjeta exposure in patients with mild (creatinine clearance [CL<sub>Cr</sub>] 60 to 90 mL/min, N=200) and moderate renal impairment (CL<sub>Cr</sub> 30 to 60 mL/min, N=71) was similar to that in patients with normal renal function (CL<sub>Cr</sub> greater than 90 mL/min, N=200). No relationship between CL<sub>Cr</sub> and Perjeta exposure was observed over the range of CL<sub>Cr</sub> (27 to 244 mL/min).

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### *Excipients:*

Glacial acetic acid,

l-histidine,

polysorbate 20,

sucrose,

water for injections.

## 6.2 Incompatibilities

No incompatibilities between Perjeta and polyvinylchloride (PVC), polyethylene or non-PVC polyolefin bags have been observed.

Glucose (5 %) solution should not be used to dilute Perjeta since it was chemically and physically unstable in such solutions.

Perjeta must not be mixed with other medicines except those mentioned above.

## 6.3 Shelf life

24 months

## 6.4 Special precautions for storage

Store vials in a refrigerator between 2 °C - 8 °C.

This medicine should not be used after the expiry date (EXP) shown on the pack.

Keep vial in the outer carton in order to protect from light until required for use.

Do not freeze. Do not shake.

Store out of reach of children.

### ***Shelf-life of the solution for infusion containing Perjeta***

Perjeta does not contain any antimicrobial preservative; therefore, care must be taken to ensure the sterility of the prepared solution.

***Diluted solution:*** Chemical and physical in-use stability has been demonstrated for 24 hours at 30 °C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 °C to 8 °C, unless dilution has taken place in controlled and validated aseptic conditions.

## 6.5 Nature and contents of container

Perjeta is presented as a colourless Type I glass vial, sealed with a grey butyl rubber stopper and crimped with a silver aluminium cap fitted with a pink flip-off plastic disc. Pack of 1 vial in a carton.

## 6.6 Special Instructions for Use, Handling and Disposal

Perjeta is for single use only as it does not contain any antimicrobial preservative. Therefore, care must be taken to ensure the sterility of the prepared solution for infusion and should be prepared by a healthcare professional using aseptic technique.

The vial must not be shaken. All the Perjeta concentrate should be withdrawn from the vial (14 mL) using sterile needle and syringe and diluted into a 250 mL PVC or non-PVC polyolefin infusion bag of sodium chloride 9 mg/mL (0,9 %) solution for infusion. Do not withdraw saline out of the infusion bag.

After dilution, the solution should contain a nominal concentration of 3,0 mg/mL of Perjeta for the initial dose where two vials are required and 1,6 mg/mL of Perjeta for the subsequent dose where one vial is required. Glucose (5 %) solution should not be used, see *Incompatibilities* below.

The bag should be gently inverted to mix the solution in order to avoid foaming.

Perjeta should be inspected visually for particulates and discolouration prior to administration. If particulates or discoloration are observed, the solution should not be used. Once the infusion is prepared it should be administered immediately.

## 7. HOLDER OF CERTIFICATE OF REGISTRATION

Roche Products (Pty) Ltd

90 Bekker Road, Hertford Office Park

Building E, Vorna Valley

Midrand, Johannesburg, 1686

South Africa

Roche Ethical Assistance Line (REAL) toll-free: 0800 21 21 25

## **8. REGISTRATION NUMBER**

48/30.1/0370

## **9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of registration: 25 March 2019

## **10. DATE OF REVISION OF THE TEXT**

Last revision: 18 October 2022

Namibia: NS2 19/26/0032

Botswana: NS2 BOT2103721

Zimbabwe: PP 2019/9.7/5926

### **Approved Manufacturer(s):**

Roche Diagnostics GmbH

Sandhoferstrasse 116

68305 Mannheim

Germany