

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S0

BABY'S OWN COUGH SYRUP

45,52 mg ammonium chloride per 5 ml syrup.

Contains 2,49 mg methylparaben and 0,521 mg benzoic acid, 1,87 g glucose and 2,80 g sucrose per 5 ml syrup.

Read all of this leaflet carefully because it contains important information for you

BABY'S OWN COUGH SYRUP is available without a doctor's prescription, for you to treat a mild illness. Nevertheless, you still need to use BABY'S OWN COUGH SYRUP carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share BABY'S OWN COUGH SYRUP with any other person.
- Ask your health care provider or pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or do not improve after 10 days.

What is in this leaflet

1. What BABY'S OWN COUGH SYRUP is and what it is used for
2. What you need to know before you use BABY'S OWN COUGH SYRUP
3. How to use BABY'S OWN COUGH SYRUP
4. Possible side effects
5. How to store BABY'S OWN COUGH SYRUP
6. Contents of the pack and other information

1 What BABY'S OWN COUGH SYRUP is and what it is used for

BABY'S OWN COUGH SYRUP is used for the temporary relief/alleviation of cough.

2 What you need to know before you take BABY'S OWN COUGH SYRUP

Do not take BABY'S OWN COUGH SYRUP:

- if you are hypersensitive (allergic) to ammonium chloride or any of the ingredients of BABY'S OWN COUGH SYRUP (listed in section 6).
- If you have kidney or liver insufficiencies.

Warnings and precautions

Special care should be taken with BABY'S OWN COUGH SYRUP:

- If you have high blood pressure or a heart disease, because BABY'S OWN COUGH SYRUP may lead to high blood pressure, water retention and electrolyte abnormalities.

Persistent coughs should be investigated for possible underlying cause by your doctor.

Children

Do not give BABY'S OWN COUGH SYRUP to children under 2 years of age.

Other medicines and BABY'S OWN COUGH SYRUP

Always tell your healthcare provider if you are taking any other medicine. (This includes complementary or traditional medicines.)

Pregnancy and breastfeeding

The safety of BABY'S OWN COUGH SYRUP in pregnancy and lactation has not been established.

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking this medicine.

Driving and using machines

Large doses of BABY'S OWN COUGH SYRUP may cause drowsiness and interfere with your ability to drive or use machinery.

BABY'S OWN COUGH SYRUP contains benzoic acid, methylparaben, sugar and Carmoisine.

Benzoic acid may increase jaundice (yellowing of the skin and eyes) in newborn babies (up to 4 weeks old). Do not give BABY'S OWN COUGH SYRUP to children under 2 years of age.

BABY'S OWN COUGH SYRUP contains sucrose and glucose, which may have an effect on the blood sugar levels of diabetic patients.

If you suffer from a rare hereditary disease such as fructose-intolerance, glucose-galactose mal-absorption or sucrase-isomaltase insufficiency, you should not take BABY'S OWN COUGH SYRUP.

BABY'S OWN COUGH SYRUP contains methylparaben as preservative and Carmoisine CI 14720 as colourant. These ingredients may cause allergic reactions.

3 How to take BABY'S OWN COUGH SYRUP

Do not share medicines prescribed for you with any other person.

Always take/give BABY'S OWN COUGH SYRUP exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure.

Doses must be taken with an adequate amount of fluid (half to one glass).

Shake the bottle well before use.

The usual dose, to be given orally 2 to 4 hourly is as follows:

Children:

2 to 6 years: 7,5 ml in water

6 to 12 years: 15 ml in water

Adults:

15 ml to 30 ml in water

If you take or give more BABY'S OWN COUGH SYRUP than you should

Large doses of BABY'S OWN COUGH SYRUP may cause:

- Nausea
- Vomiting
- Thirst
- Headache
- abnormally deep and rapid breathing, breathing is difficult and the breath has a characteristic odour, kidney failing to produce urine may occur (anuria)
- a feeling of warmth
- muscle twitching
- progressive drowsiness, delirium (confusion) and dizziness
- recurrent fits

- depression of the central nervous system and coma
- gastric (stomach) irritation and abdominal pain (colic)
- abnormal high body fluid and tissue acidity (acidosis)
- abnormal low doses of potassium in the blood (hypokalaemia)
- the capacity of the liver to convert ammonium ion to urea is exceeded (hepatic encephalopathy).

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre.

Treatment of overdosage is symptomatic and supportive.

If you forget to take or give a dose of BABY'S OWN COUGH SYRUP

Do not take or give a double dose to make up for forgotten individual doses.

4 Possible side effects

BABY'S OWN COUGH SYRUP can have side effects.

Possible side effects include:

- nausea
- vomiting
- abnormal high body fluid and tissue acidity (acidosis)
- abnormal low doses of potassium in the blood (hypokalaemia).

Not all side effects reported for BABY'S OWN COUGH SYRUP are included in this leaflet.

Should your general health worsen or if you experience any untoward effects while taking BABY'S OWN COUGH SYRUP, please consult your health care provider for advice.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of BABY'S OWN COUGH SYRUP.

5 How to store BABY'S OWN COUGH SYRUP

Store all medicines out of reach of children.

Store at or below 25 °C and protect from light.

Do not use after the expiry date stated on the label.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6 Content of the pack and other information

What BABY'S OWN COUGH SYRUP contains

- The active substance is 45,52 mg ammonium chloride per 5 ml syrup.
- The preservatives are methylparaben 0,05 % *m/v* and benzoic acid 0,01 % *m/v*.
- The other ingredients are anise oil, camphor, carmosine, Clark's Caramel, glacial acetic acid, liquorice liquid extract, purified water and strong ammonia solution.
- BABY'S OWN COUGH SYRUP contains 1,87 g glucose and 2,80 g sucrose per 5 ml syrup.

What BABY'S OWN COUGH SYRUP looks like and contents of the pack

Mentholatum SA (Pty) Ltd
Baby's Own Cough Syrup
45,52 mg Ammonium chloride per 5 ml syrup

BABY'S OWN COUGH SYRUP is a red syrupy liquid.

BABY'S OWN COUGH SYRUP is presented in glass bottles of 50 ml, [and] 100 ml and 200 ml.

Holder of Certificate of Registration

Mentholatum South Africa (Pty) Ltd

1st Floor, Silverberg Terrace

Steenberg Office Park

Silverwood Close

Tokai

Cape Town

7945

Tel No: 021 – 702 0620 / 21

This leaflet was last revised in

15 May 2023

Registration number

G600 (Act 101/1965)

Access to the corresponding Professional Information

The PI can be found online under Mentholatum SA's website:

<https://www.mentholatum.co.za/product-information/>.