

Approved Professional Information for Medicines for Human Use:

BIMAGAM 0,03 %

SCHEDULING STATUS

S4

1. NAME OF THE MEDICINE

BIMAGAM 0,03 % eye drops, solution

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each mL of sterile solution contains bimatoprost 0,3 mg.

Contains benzalkonium chloride 0,005 % (*m/v*) as preservative.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Eye drops, solution.

BIMAGAM 0,03 % Eye drops, solution

Colourless solution that is a practically clear and free of particles.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Reduction of elevated intraocular pressure in chronic open-angle glaucoma and ocular hypertension (as monotherapy or as adjunctive therapy to beta-blockers).

4.2 Posology and method of administration

Posology

When used as monotherapy or as adjunctive therapy, the recommended dose is one

Austell Pharmaceuticals (Pty) Ltd, 540318, Bimagam 0,03 %, Eye drops, solution

drop of BIMAGAM 0,03 % eye drops in the affected eye(s) once daily, administered in the evening.

The dose should not exceed once daily as more frequent administration may lessen the intraocular pressure lowering effect.

Special populations

Elderly population

No dosage adjustment in elderly patients is necessary.

Hepatic and renal impairment

BIMAGAM 0,03 % eye drops has not been studied in patients with renal or moderate to severe hepatic impairment and should therefore be used with caution in such patients. In patients with a history of mild liver disease or abnormal ALT, AST and/or bilirubin at baseline, BIMAGAM 0,03 % eye drops had no adverse effect on liver function over 24 months.

Paediatric population

BIMAGAM 0,03 % eye drops has only been studied in adults and therefore its use is not recommended in children or adolescents (under the age of 18).

Method of administration

BIMAGAM 0,03 % is for ocular use only.

To prevent contamination of the dropper tip and solution, care should be taken not to touch the eyelids, surrounding areas or other surfaces with the dropper tip of the bottle.

If more than one topical ophthalmic medication is being used, the medicines should be administered at least 5 minutes apart.

4.3 Contraindications

- Hypersensitivity to bimatoprost, benzalkonium chloride or to any of the other excipients listed in section 6.1.

4.4 Special warnings and precautions for use

Ocular

Before treatment is initiated, patients should be informed of the possibility of prostaglandin analogue periorbitopathy (PAP) eyelash growth, darkening of the eyelid skin and increased iris pigmentation, since these have been observed during treatment with bimatoprost as in BIMAGAM 0,03 %. Some of these changes may be permanent and may lead to differences in appearance between the eyes when only one eye is treated. Increased iris pigmentation is likely to be permanent. The pigmentation change is due to increased melanin content in the melanocytes rather than to an increase in the number of melanocytes. The long-term effects of increased iris pigmentation are not known. Iris colour changes seen with ophthalmic administration of bimatoprost may not be noticeable for several months to years.

Typically, the brown pigmentation around the pupil spreads concentrically towards the periphery of the iris and the entire iris or parts become more brownish. Neither naevi nor freckles of the iris appear to be affected by the treatment. At 12 months, the incidence of iris hyperpigmentation with bimatoprost 0,3 mg/mL was 1,5 % (see section 4.8) and did not increase following 3 years treatment. Periorbital tissue pigmentation has been reported to be reversible in some patients.

Cystoid macular oedema has been less frequently reported following treatment with bimatoprost 0,3 mg/mL eye drops, solution. Therefore, BIMAGAM 0,03 % should be used with caution in patients with known risk factors for macular oedema (e.g. aphakic patients, pseudophakic patients with a torn posterior lens capsule).

There have been less frequent spontaneous reports of reactivation of previous corneal infiltrates or ocular infections with bimatoprost 0,3 mg/mL eye drops, solution. BIMAGAM 0,03 % should be used with caution in patients with a prior history of significant ocular viral infections (e.g. herpes simplex) or uveitis/iritis.

BIMAGAM 0,03 % has not been studied in patients with inflammatory ocular conditions, neovascular, inflammatory, angle-closure glaucoma, congenital glaucoma or narrow-angle glaucoma.

Skin

There is a potential for hair growth to occur in areas where BIMAGAM 0,03 % solution comes repeatedly in contact with the skin surface. Thus, it is important to apply BIMAGAM 0,03 % as instructed and avoid it running onto the cheek or other skin areas.

Respiratory

BIMAGAM 0,03 % has not been studied in patients with compromised respiratory function. While there is limited information available on patients with a history of asthma or COPD, there have been reports of exacerbation of asthma, dyspnoea and COPD, as well as reports of asthma, in post marketing experience. The frequency of

these symptoms is not known. Patients with COPD, asthma or compromised respiratory function due to other conditions should be treated with caution.

Cardiovascular

BIMAGAM 0,03 % has not been studied in patients with heart block more severe than first degree or uncontrolled congestive heart failure. There have been a limited number of spontaneous reports of bradycardia or hypotension with bimatoprost 0,3 mg/mL eye drops, solution. BIMAGAM 0,03 % should be used with caution in patients predisposed to low heart rate or low blood pressure.

Other information

Concomitant use with other prostaglandin analogues

In studies of bimatoprost 0,3 mg/mL in patients with glaucoma or ocular hypertension, it has been shown that the more frequent exposure of the eye to more than one dose of bimatoprost daily may decrease the IOP-lowering effect (see section 4.5). Patients using BIMAGAM 0,03 % with other prostaglandin analogues should be monitored for changes to their intraocular pressure.

Bacterial keratitis

There have been reports of bacterial keratitis associated with the use of multiple dose containers of topical ophthalmic products. These containers had been inadvertently contaminated by patients who, in most cases, had a concurrent ocular disease.

Patients with a disruption of the ocular epithelial surface are at greater risk of developing bacterial keratitis.

Austell Pharmaceuticals (Pty) Ltd, 540318, Bimagam 0,03 %, Eye drops, solution

Patients should be instructed to avoid allowing the tip of the dispensing container to contact the eye or surrounding structures to avoid eye injury and contamination of the solution.

Contact lenses

BIMAGAM 0,03 % contains the preservative benzalkonium chloride, which may be absorbed by and cause discolouration of soft contact lenses. Patients wearing soft (hydrophilic) contact lenses should be instructed to remove contact lenses prior to administration of BIMAGAM 0,03 % and wait at least 15 minutes following administration before reinserting soft contact lenses.

BIMAGAM 0,03 % should not be administered while wearing contact lenses.

Excipient: Benzalkonium chloride

BIMAGAM 0,03 % contains 0,05 mg benzalkonium chloride (a preservative) in each millilitre eye drop solution which is equivalent to 0,005 % (*m/v*).

As the possibility of adverse effects on the corneal permeability, and the danger of disruption of the corneal epithelium with prolonged or repeated usage of benzalkonium chloride preserved ophthalmological preparations, cannot be excluded, regular ophthalmological examination is required. Caution should be exercised in the use of benzalkonium chloride preserved topical medication over an extended period in patients with extensive ocular surface disease.

4.5 Interaction with other medicines and other forms of interaction

No interaction studies have been performed.

No interactions are anticipated in humans, since systemic concentrations of bimatoprost are extremely low (less than 0,2 ng/mL) following ocular dosing with bimatoprost 0,3 mg/mL eye drops, solution. Bimatoprost is biotransformed by any of multiple enzymes and pathways, and no effects on hepatic medicine metabolising enzymes were observed in preclinical studies.

In clinical studies, BIMAGAM 0,03 % eye drops was used concomitantly with a number of different ophthalmic beta-blocking medicines without evidence of interactions.

Concomitant use of BIMAGAM 0,03 % and antiglaucomatous medicines other than topical beta-blockers has not been evaluated during adjunctive glaucoma therapy.

There is a potential for the IOP-lowering effect of prostaglandin analogues (e.g. BIMAGAM 0,03 %) to be reduced in patients with glaucoma or ocular hypertension when used with other prostaglandin analogues (see section 4.4).

4.6 Fertility, pregnancy and lactation

The safety of BIMAGAM 0,03 % during pregnancy and lactation has not been established.

Pregnancy

There are no adequate data from the use of bimatoprost in pregnant women. Animal studies have shown reproductive toxicity at high maternotoxic doses.

BIMAGAM 0,03 % should not be used during pregnancy unless clearly necessary.

Breastfeeding

It is unknown whether bimatoprost is excreted in human breast milk. Animal studies have shown excretion of bimatoprost in breast milk. It is recommended that BIMAGAM 0,03 % not be used in breastfeeding mothers.

Fertility

There are no data on the effects of bimatoprost on human fertility.

4.7 Effects on ability to drive and use machines

BIMAGAM 0,03 % has negligible influence on the ability to drive and use machines. if transient blurred vision or dizziness occurs at instillation, the patient should wait until the vision clears or dizziness subsides before driving or using machines.

4.8 Undesirable effects

a. Summary of the safety profile

In clinical studies, over 1800 patients have been treated with bimatoprost 0,3 mg/mL eye drops. On combining the data from phase III monotherapy and adjunctive bimatoprost 0,3 mg/mL eye drops usage, the most frequently reported adverse reactions were:

- growth of eyelashes in the first year with the incidence of new reports decreasing at 3 years
- conjunctival hyperaemia (mostly trace to mild and thought to be of a non-inflammatory nature) in the first year with the incidence of new reports decreasing at 3 years
- ocular pruritus in the first year with the incidence of new reports decreasing at 3 years.

b. Tabulated list of adverse reactions

The table below shows all adverse drug reactions (ADRs) observed during clinical trials and postmarket spontaneous reports with bimatoprost 0,3 mg/mL eye drops, solution. Most were ocular, mild and none was serious.

System Organ	Frequency		
Class	Frequent	Less Frequent	Not known
Immune system disorders			Hypersensitivity reaction including signs and symptoms of eye allergy and allergic dermatitis
Nervous system disorders	Headache	Dizziness	
Eye disorders	Conjunctival / ocular hyperaemia, eye (ocular) pruritus, prostaglandin analogue periorbitopathy, superficial punctate keratitis, corneal erosion, ocular burning, ocular irritation, allergic conjunctivitis, blepharitis,	Retinal haemorrhage, uveitis, cystoid macular oedema, iritis, blepharospasm, eyelid retraction, periorbital erythema, eyelid oedema	Periorbital and lid changes including deepening of the eyelid sulcus, ocular discomfort

	worsening of visual acuity, asthenopia, conjunctival oedema, foreign body sensation, ocular dryness, eye pain, photophobia, tearing / lacrimation increased, eye discharge, visual disturbance/blurred vision, increased iris pigmentation, eyelash growth, Eyelash darkening, eyelid erythema, eyelid pruritus cataract		
--	--	--	--

Vascular disorders	Hypertension		
Respiratory, thoracic and mediastinal disorders			Asthma, asthma exacerbation, COPD exacerbation and dyspnoea
Gastrointestinal disorders		Nausea	
Skin and subcutaneous tissue disorders	Pigmentation of periocular skin skin hyperpigmentation, blepharal pigmentation	Hirsutism	Skin discoloration (periocular) abnormal hair growth
General disorders and administration site conditions		Asthenia, peripheral oedema	
Investigations	Liver function test abnormal		
Infections and infestations		Infection (primarily colds and upper respiratory tract infections)	

c. Description of selected adverse reactions

Prostaglandin analogue periorbitopathy (PAP)

Prostaglandin analogues including bimatoprost as in BIMAGAM 0,03 % can induce periorbital lipodystrophic changes which can lead to deepening of the eyelid sulcus, ptosis, enophthalmos, eyelid retraction, involution of dermatochalasis and inferior scleral show. Changes are typically mild, can occur as early as one month after initiation of treatment with bimatoprost as in BIMAGAM 0,03 %, and may cause impaired field of vision even in the absence of patient recognition. PAP is also associated with periocular skin hyperpigmentation or discolouration and hypertrichosis. All changes have been noted to be partially or fully reversible upon discontinuation or switch to alternative treatments.

Iris hyperpigmentation

Increased iris pigmentation is likely to be permanent. The pigmentation change is due to increased melanin content in the melanocytes rather than to an increase in the number of melanocytes. The long-term effects of increased iris pigmentation are not known. Iris colour changes seen with ophthalmic administration of bimatoprost as in BIMAGAM 0,03 % may not be noticeable for several months to years. Typically, the brown pigmentation around the pupil spreads concentrically towards the periphery of the iris and the entire iris or parts become more brownish. Neither naevi nor freckles of the iris appear to be affected by the treatment. At 12 months, the incidence of iris hyperpigmentation with bimatoprost as in BIMAGAM 0,03 % eye drops, solution was 0,5 %. At 12 months, the incidence with bimatoprost 0,03 % eye drops, solution was 1,5 % and did not increase following 3 years treatment.

Adverse reactions reported in phosphate containing eye drops

Cases of corneal calcification have been reported less frequently in association with the use of phosphate containing eye drops in some patients with significantly damaged corneas.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare professionals are asked to report any suspected adverse reactions to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>

4.9 Overdose

No case of overdose has been reported and is unlikely to occur after ocular administration.

If overdose occurs, treatment should be symptomatic and supportive.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Category and Class: A. 15.4 Ophthalmological preparations. Others

Pharmacotherapeutic group: Ophthalmologicals, prostaglandin analogues, ATC

Code: S01EE03

Mechanism of action

Bimatoprost is an ocular hypotensive medicine. It is a synthetic prostamide, structurally related to prostaglandin F_{2α} that does not act through any known prostaglandin receptors. Bimatoprost selectively mimics the effects of prostamides. The prostamide receptor, however, has not yet been structurally identified.

Bimatoprost reduces intraocular pressure (IOP) in humans by increasing aqueous humour outflow through the trabecular meshwork and enhancing uveoscleral outflow.

Reduction of the intraocular pressure starts approximately 4 hours after the first administration and maximum effect is reached within approximately 8 to 12 hours.

The duration of effect is maintained for 24 hours.

Austell Pharmaceuticals (Pty) Ltd, 540318, Bimagam, Eye drops, solution and 0,03 %
Limited experience is available with the use of bimatoprost in patients with open-angle glaucoma with pseudoexfoliative and pigmentary glaucoma, and chronic angle-closure glaucoma with patent iridotomy and no recommendation can be made.

No clinically relevant effects on heart rate and blood pressure have been observed in clinical trials.

5.2 Pharmacokinetic properties

Absorption

Bimatoprost penetrates the human cornea and sclera well *in vitro*. After ocular administration, the systemic exposure of bimatoprost is very low with no accumulation over time.

After once daily ocular administration of one drop of 0,03 % bimatoprost to both eyes of 15 healthy subjects for two weeks, blood concentrations peaked within 10 minutes after dosing and declined to below the lower limit of detection (0,025 ng/ mL) in most subjects within 1,5 hours after dosing.

Mean C_{max} and $AUC_{0-24hrs}$ values were similar on days 7 and 14 at approximately 0,08 ng/ mL and 0,09 ng•hr/ mL respectively, indicating that a steady medicine concentration was reached during the first week of ocular dosing.

Distribution

Bimatoprost is moderately distributed into body tissues and the systemic volume of distribution in humans at steady state was 0,67 L/kg.

Austell Pharmaceuticals (Pty) Ltd, 540318, Bimagam, Eye drops, solution and 0,03 %

In human blood, bimatoprost resides mainly in the plasma. The plasma protein binding of bimatoprost is approximately 88 %.

Biotransformation

Bimatoprost is not extensively metabolised in the human eye. Bimatoprost is the major circulating component in the blood once it reaches the systemic circulation following ocular dosing. Bimatoprost then undergoes oxidation, N-deethylation and glucuronidation to form a diverse variety of metabolites.

Elimination

Bimatoprost is eliminated primarily by renal excretion, up to 67 % of an **intravenous** dose administered to healthy volunteers was excreted in the urine, 25 % of the dose was excreted via the faeces. The elimination half-life, determined after **intravenous** administration, was approximately 45 minutes, the total blood clearance was 1,5 L/hr/kg.

Characteristics in elderly patients

After twice daily dosing, the mean AUC_{0-24hr} value of 0,0634 ng•hr/ mL bimatoprost in the elderly (subjects 65 years or older) were significantly higher than 0,0218 ng•hr/ mL in young healthy adults. However, this finding is not clinically relevant as systemic exposure for both elderly and young subjects remained very low from ocular dosing. There was no accumulation of bimatoprost in the blood over time and the safety profile was similar in elderly and young patients.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Benzalkonium chloride

Citric acid monohydrate

Dibasic sodium phosphate heptahydrate

Purified water

Sodium chloride

Sodium hydroxide or hydrochloric acid (for pH-adjustment)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

36 months.

28 days (4 weeks) after first opening.

6.4 Special precautions for storage

Store unopened container at or below 25 °C in the original carton, until required for use.

Opened container must be stored at or below 25 °C. Do not use more than 28 days after opening.

6.5 Nature and contents of container

BIMAGAM 0,03 % eye drops, solution is filled in a white LDPE bottle with white LDPE dropper insert, and closed with a white, tamper-proof HDPE screw cap. Each bottle has a fill volume of 3 mL. Subsequently, the bottle is packed into the respective folding carton together with the leaflet.

The following pack sizes are available: cartons containing 1 or 3 bottles of 3 mL solution. Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements for disposal.

7. HOLDER OF CERTIFICATE OF REGISTRATION

Austell Pharmaceuticals (Pty) Ltd

1 Sherborne Road

Parktown

JOHANNESBURG

2193

South Africa

Tel: 0860287835

8. REGISTRATION NUMBER

BIMAGAM 0,03 %: 54/15.4/0318

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

17 October 2023

10. DATE OF REVISION OF THE TEXT