

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

PROFESSIONAL INFORMATION

SCHEDULING STATUS: **S3**

1. NAME OF THE MEDICINE

Akliel 50 microgram/g cream

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One gram of cream contains 50 micrograms of trifarotene.

Excipients with known effect

One gram of cream contains 300 milligrams of propylene glycol (E1520) and 50 milligrams of ethanol.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Cream.

White and homogenous cream

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Akliel is indicated for the cutaneous treatment of moderate *Acne Vulgaris* of the face and/or the trunk in patients from 12 years of age and older, when many comedones, papules and pustules are present.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Aklief
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

4.2 Posology and method of administration

Posology

Apply a thin layer of Aklief cream to the affected areas of the face and/or trunk once a day, in the evening, on clean and dry skin.

It is recommended that the medical practitioner assesses the continued improvement of the patient after three months of treatment.

Special populations

Renal and Hepatic impairment

Aklief has not been studied in patients with renal and hepatic impairment.

Elderly patients

The safety and efficacy of Aklief in elderly patients aged 65 years and above have not been established.

Paediatric population

The safety and efficacy of Aklief in children below 12 years old have not been established.

Method of administration

Cutaneous use only.

Before using the pump for the first time, prime it by pressing down several times until a small amount of medicine is dispensed (up to 10 times maximum). The pump is now ready to use.

Apply a thin layer of Aklief cream to the affected areas of the face (forehead, nose, chin and right and left cheeks) and all affected areas of the trunk once a day, in the evening, on clean and dry skin:

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

- One pump actuation should be enough to cover the face (i.e. forehead, cheeks, nose, and chin).
- Two pump actuations should be enough to cover the upper trunk (i.e. reachable upper back, shoulders and chest). One additional pump actuation may be used for middle and lower back if acne is present.

Patients should be instructed to avoid contact with the eyes, eyelids, lips and mucous membranes and to wash their hands after applying Akliel.

The use of a moisturiser is recommended as needed from the initiation of treatment, while allowing sufficient time before and after the application of Akliel cream to allow the skin to dry.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Pregnancy (see section 4.6)
- Women planning a pregnancy

4.4 Special warnings and precautions for use

Erythema, scaling, dryness, and stinging/burning may be experienced with use of Akliel cream (see section 4.8). To mitigate the risk of such reactions, patients should be instructed to use a moisturiser from the initiation of treatment, and, if appropriate, reduce the frequency of application of Akliel cream, or suspend use temporarily. Despite mitigation measures, if severe reactions persist the treatment may be discontinued.

Akliel should not be applied to cuts, abrasions, eczematous or sunburned skin.

Use of "waxing" as a depilatory method should be avoided on skin treated with Akliel.

If a reaction suggesting sensitivity to any component of the formula occurs, the use of Akliel should be discontinued. Caution should be exercised if cosmetics or acne medicines with

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

desquamative, irritant or drying effects are concomitantly used with Akliel, as they may produce additive irritant effects.

Akliel should not come into contact with the eyes, eyelids, lips, or mucous membranes. If the product enters the eye, wash immediately and abundantly with lukewarm water.

Excessive exposure to sunlight, including sunlamps or phototherapy should be avoided during the treatment. Use of a broad-spectrum, water-resistant sunscreen with a Sun Protection Factor (SPF) of 30 or higher and protective clothing over treated areas is recommended when exposure cannot be avoided.

Akliel cream contains propylene glycol (E1520) that may cause skin irritation.

Akliel also contains 50 mg alcohol (ethanol) in each gram which is equivalent to 5% w/w. It may cause burning sensation on damaged skin.

4.5 Interaction with other medicines and other forms of interaction

Effect of Akliel cream on other medicines

A clinical drug-drug interaction study has shown that topical application of trifarotene did not affect the circulating concentrations of hormonal contraceptives (ethinyl estradiol and levonorgestrel) administered by oral route.

Effect of other medicines on Akliel cream

No clinical drug-drug interaction studies were performed to assess effects of other medicines on trifarotene systemic levels (see section 5.2).

There is no data on the pharmacodynamic interaction potential of trifarotene.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

Caution should be exercised if cosmetics or acne medicines with desquamative, irritant or drying effects are concomitantly used with Akliel as they may produce additive irritant effects (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

Akliel is contraindicated (see section 4.3) during pregnancy or in women planning a pregnancy. Studies in animals with trifarotene by the oral route have shown reproductive toxicity at high systemic exposure (see section 5.3).

If Akliel is used during pregnancy, or if the patient becomes pregnant while taking Akliel, treatment should be discontinued.

Breastfeeding

It is unknown whether trifarotene or its metabolites are excreted in human milk.

Available pharmacodynamic/toxicological data in animals have shown excretion of trifarotene/metabolites in milk (for details see 5.3).

A risk to the suckling child cannot be excluded.

Use of Akliel during breastfeeding is not recommended.

To avoid the risk of ingestion by, and/or contact exposure of, an infant, nursing women should not apply Akliel cream to the chest or breast area.

Fertility

No human fertility studies were conducted with Akliel.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

No effects of trifarotene on fertility were found in rats in reproductive studies of oral administration. However, after oral administration to dogs, findings of Germ cell degeneration were observed, see section 5.3.

4.7 Effects on ability to drive and use machines

Akliel has no or negligible influence on the ability to drive and use machines.

4.8 Undesirable effects

Summary of the safety profile

Local cutaneous reactions such as erythema, scaling, dryness, and stinging/burning) were collected separately from other adverse events as a measure of local tolerance. These cutaneous reactions are very common and of mild, moderate and severe intensity for up to 39%, 29.7% and 6.2% of patients, respectively on the face. On the trunk, up to 32.9%, 18.9%, 5.2% of patients had mild, moderate and severe reactions respectively. The maximum severity typically occurred at Week 1 for the face, and at Week 2 to 4 for the trunk, and decreased with continued use of the medication (see section 4.4).

The most “commonly” reported adverse reactions as described below in Table 1 are application site irritation, application site pruritus and sunburn, occurring in 1.2% to 6.5% of patients treated with Akliel cream in clinical studies.

Tabulated summary of adverse reactions

Adverse reactions reported in the 12-week vehicle-controlled Phase 3 studies in 1220 patients treated with Akliel cream (and for which the rate for Akliel cream exceeds the rate for vehicle cream) are presented in Table 1.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
 Product name: Akliel
 Dosage form: Cream
 Strength: 50 µg Trifarotene per 1 g of cream
 Registration number: 56/13.12/1007

The adverse reactions are classified by System Organ Class and frequency, using the following convention: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

Table 1 – Adverse reactions

System Organ Class	Frequency	Adverse reactions
<i>General disorders and administration site conditions</i>	Common	Application site irritation Application site pruritus
	Uncommon	Application site pain Application site dryness Application site discolouration Application site erosion Application site rash Application site swelling
	Rare	Application site erythema Application site urticaria Application site vesicles
<i>Injury, poisoning and procedural complications</i>	Common	Sunburn
<i>Skin and subcutaneous tissue disorders</i>	Uncommon	Skin irritation Acne Dermatitis allergic Erythema
	Rare	Eczema asteatotic

Applicant: Galderma Laboratories South Africa (Pty) Ltd
 Product name: Akliel
 Dosage form: Cream
 Strength: 50 µg Trifarotene per 1 g of cream
 Registration number: 56/13.12/1007

		Seborrheic dermatitis Skin burning sensation Skin fissures Skin hyperpigmentation
<i>Eye disorders</i>	Rare	Eyelid exfoliation Eyelid oedema
<i>Gastrointestinal disorders</i>	Rare	Cheilitis
<i>Vascular disorders</i>	Rare	Flushing

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers are asked to report any suspected adverse reactions to SAHPRA via the “6.04 Adverse Drug Reactions Reporting Form”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>

4.9 Overdose

Akliel is for once-daily cutaneous use only.

If the medicine is applied excessively, no more rapid or better results will be obtained and marked redness, scaling, or skin discomfort may occur. In this event, discontinue use and wait until the skin has recovered.

In case of accidental ingestion, appropriate symptomatic measures should be taken. Chronic ingestion of Akliel may lead to the same side effects as those associated with excessive oral intake of vitamin A.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Category and class: A.13.12 Dermatological preparations – Others

ATC code: D10AD06

Mechanism of action

Akliel contains 50 micrograms (mcg/g) (w/w) trifarotene, which is a chemically stable, terphenyl acid derivative with retinoid like activity. It is a potent RAR γ agonist (retinoid acid receptor γ agonist), characterised by its high specificity to this receptor over RAR α & RAR β (50- and 8-fold, respectively, with no Retinoid X Receptor (RXR) activity).

In addition, trifarotene modulates retinoid target genes (differentiation and inflammatory processes) in immortalised keratinocytes and reconstructed epidermis.

Pharmacodynamic effects

Trifarotene has demonstrated, in the Rhino-mouse model, marked comedolytic activity with the reduction in the comedone count and marked increased epidermis thickness. In this model, trifarotene produced the same comedolytic effect as other known retinoids, at about 10 times lower dose.

Trifarotene has also shown anti-inflammatory and depigmenting activities.

Clinical efficacy and safety

Akliel cream applied once daily in the evening was evaluated for 12 weeks in 2 randomised, multi-center, parallel group, double-blind, vehicle-controlled studies of identical design. They were conducted in a total of 2420 patients aged, 9 years and older, with moderate facial and truncal acne vulgaris.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
 Product name: Akliel
 Dosage form: Cream
 Strength: 50 µg Trifarotene per 1 g of cream
 Registration number: 56/13.12/1007

Acne severity was evaluated using the 5-point Investigator’s Global Assessment (IGA) scale for the face and Physician’s Global Assessment (PGA) for the trunk, with moderate acne vulgaris defined as a score of Grade 3-Moderate (see Table 2).

Table 2 Investigator’s Global Assessment and Physician’s Global Assessment Scales

0	Clear	Clear skin with no inflammatory or non-inflammatory lesions.
1	Almost Clear	A few scattered comedones and a few small papules.
2	Mild	Easily recognizable; less than half the surface is involved. Some comedones and some papules and pustules.
3	Moderate	More than half of the surface is involved. Many comedones, papules and pustules. One nodule may be present.
4	Severe	Entire surface is involved. Covered with comedones, numerous papules and pustules. Few nodules may be present.

There were three co-primary efficacy endpoints in both pivotal studies 1) the success rate based on the IGA and PGA outcome (percentage of subjects “clear” and “almost clear” and with at least a 2-grade change from baseline) and absolute and percentage change from baseline in 2) inflammatory and 3) non-inflammatory lesion counts at Week 12.

Overall, 87% of subjects were Caucasian and 55% were female. Thirty four (1.4%) subjects were 9 to 11 years of age, 1128 (47%) subjects were 12 to 17 years and 1258 (52%) subjects were 18 years and older. All patients had moderate acne vulgaris on the face and 99% on the trunk. At baseline subjects had between 7 and 200 (average 36) inflammatory lesions on the

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

face and between 0 and 220 (average 38) on the trunk. Additionally, subjects had 21 to 305 (average 52) non-inflammatory lesions on the face and 0 to 260 (average 46) on the trunk.

The efficacy assessed from total 2420 patients 9 years and older showed Akliel cream is effective in the treatment of moderate acne compared to vehicle. The results from the face showed a significant improvement ($p < 0.001$) of the Success Rate (percentage of subjects “clear” and “almost clear” and with at least a 2-grade change from baseline) and a significant reduction in absolute change ($p < 0.001$) and percentage change ($p < 0.001$) from baseline for both inflammatory lesions and non-inflammatory lesions. The results from the trunk also showed Akliel cream achieved significant improvement in PGA Success Rate (percentage of subjects “clear” and “almost clear” and with at least a 2-grade change from baseline) ($p < 0.001$), and significant reduction in absolute change ($p < 0.001$) and percentage change ($p < 0.001$) from baseline for both inflammatory lesions and non-inflammatory lesions.

Paediatric population

Age group 9 to 11 years: In the phase 3 studies a total of only 34 children of this age group were included – 19 of them in study 18251 and 15 in study 18252. In this age group, patient number was low and efficacy could not be demonstrated.

Age group 12 to 17 years: In the phase 3 studies a total of 1128 children aged 12 to 17 years with moderate acne vulgaris were included: 573 of them in study 18251 and 555 children in study 18252.

In this age group, Akliel cream is effective in the treatment of moderate acne compared to vehicle. The results from the face showed significant improvement in the Success Rate ($p < 0.001$), and significant reduction in absolute change ($p < 0.001$) and percentage change ($p < 0.001$) from baseline for both inflammatory lesions and non-inflammatory lesions were

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

observed; the results from the trunk also showed significant improvement in PGA Success Rate, and significant reduction in absolute change ($p < 0.001$) and percentage change ($p < 0.001$) from baseline for both inflammatory lesions and non-inflammatory lesions.

Long-term efficacy

In Study 3, a one-year open label safety study of 453 patients, 9 years and older, with moderate facial and truncal acne vulgaris, Akliel cream demonstrated a clinically meaningful improvement with IGA and PGA success rates increasing:

- from 26.6% at Week 12 visit to 65.1% at Week 52 visit for the face and
- from 38.6% at Week 12 visit to 66.9% at Week 52 visit for the trunk, respectively.

IGA and PGA success experienced by the same subject increased from 22.0% at Week 12 to 57.9% at Week 52.

5.2 Pharmacokinetic properties

Absorption

The absorption of trifarotene from Akliel cream was evaluated in adult and paediatric (10-17 y.o.) subjects with acne vulgaris. Subjects were treated once daily for 30 days with 2 grams/day of Akliel applied on the face, shoulders, chest, and upper back.

Overall, systemic exposure levels were low and similar between adults and paediatric populations.

After 4 weeks treatment, seven of nineteen (37 %) adult subjects had quantifiable trifarotene plasma levels. C_{max} ranged from below the limit of quantification (LOQ < 5 pg/mL) to 10 pg/mL and AUC 0-24hR ranged from 75 to 104 pg.hr/mL.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

Three of the seventeen (18%) of paediatric subjects presented quantifiable systemic exposure.

C_{max} ranged from below the limit of quantification (LOQ <5 pg/mL) to 9 pg/mL and AUC 0-24hR ranged from 89 to 106 pg.hr/mL

Steady state conditions were achieved in both adult and paediatric subjects following 2 weeks of topical administration. No drug accumulations is expected with long-term use.

Distribution

Trifarotene penetrates into the skin with an exponential distribution from the stratum corneum to the epidermis and dermis.

An in vitro study demonstrated that trifarotene is greater than 99.9% bound to plasma proteins.

No significant binding of trifarotene to erythrocytes was observed.

Biotransformation

In vitro studies using human hepatic microsomes and recombinant CYP450 enzymes have shown that trifarotene is primarily metabolized by CYP2C9, CYP3A4, CYP2C8 and at lesser extent by CYP2B6.

Pharmacokinetic drug interaction potential

In vitro studies show that trifarotene cream at the concentrations achieved systemically after topical administration did not inhibit the CYP450 isoenzymes CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6 and 3A4, and did not induce CYP1A2, 2B6, or 3A4.

In vitro studies have shown that trifarotene cream at the concentrations achieved systemically after topical administration did not inhibit either MATE, OATP, OAT or OCT uptake transporters or BCRP, PgP, BSEP or MPR efflux transporters.

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

5.3 Preclinical safety data

Note: the animal multiples of human systemic exposure calculations were based on Area Under the Curve (AUC) comparisons for a topical human dose of 2 g of trifarotene cream, applied once daily.

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated oral dose toxicity, genotoxicity, or carcinogenic potential.

In dermal repeat dose toxicity studies in minipigs for up to 9 months, systemic exposure to trifarotene was very low, generally below the limit of quantification. There were no systemic effects and the only noteworthy finding consisted of reversible skin irritation at the application sites.

In animal reproduction studies, oral administration of trifarotene in pregnant rats and rabbits during organogenesis was teratogenic and embryotoxic at exposures (AUC) that were 1614 to 18245-times and 800 to 4622-times those observed in humans at the maximum recommended human dose (MRHD) of 2 g.

Trifarotene was not teratogenic in rats and rabbits at systemic exposures corresponding to 534 and 98-times respectively those observed in humans.

Trifarotene had no effects on pre- and post-natal development in rats, up to the highest oral doses tested which corresponded to systemic exposures (AUC) 595 to 1877-times higher than those observed in humans.

Trifarotene showed no adverse effects on fertility in rats administered orally at exposures of approximately 1754 (males) and 1877 (females) times the 2 g dose in humans. However, after oral administration to dogs, Germ cell degeneration with pyknotic/apoptotic germ cells was evident from the lowest dose tested of 0.2 mg/kg/day corresponding to a systemic exposure 1170 times higher than those observed in humans. All animals with this finding also showed hypospermatogenesis and debris in the epididymides. The findings did not completely recover

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

after 8 weeks, suggesting an extended and possibly chronic effect. As these effects were noted also at the lowest dose tested, the relevance of the findings for lower doses is unknown.

Oral study in rats have shown trifarotene and/or related metabolites are excreted into maternal milk.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Allantoin

Simulgel 600 PHA (copolymer of acrylamide and sodium acryloyldimethyltaurate, isohexadecane, polysorbate 80, sorbitan oleate)

Cyclomethicone

Ethanol

Phenoxyethanol

Propylene glycol (E1520)

Triglycerides medium-chain

Purified water

6.2 Incompatibilities

Not applicable

6.3 Shelf life

2 years

After first opening: use within 6 months.

6.4 Special precautions for storage

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

Store at or below 25 °C.

Akliel does not require any special storage conditions.

6.5 Nature and contents of container

5g tube: White Low density polyethylene (LDPE)/Aluminium (Al)/High density polyethylene (HDPE) laminated tubes with a white high density polyethylene (HDPE) head and a white polypropylene (PP) closure.

15g, 30g, 75g Multidose container with airless pump: Polypropylene (PP)/High density polyethylene (HDPE) white airless bottle closed with a white polypropylene (PP) pump and a white polypropylene (PP) overcap.

Pack sizes:

1 tube of 5 g.

1 bottle of 15, 30 or 75 g.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. HOLDER OF CERTIFICATE OF REGISTRATION

Galderma Laboratories South Africa (Pty) Ltd

Nicol Main Office Park, Block C First Floor

FutureSpace,

2 Bruton Road,

Bryanston, 2191

Applicant: Galderma Laboratories South Africa (Pty) Ltd
Product name: Akliel
Dosage form: Cream
Strength: 50 µg Trifarotene per 1 g of cream
Registration number: 56/13.12/1007

South Africa

8. REGISTRATION NUMBER

56/13.12/1007

9. DATE OF FIRST AUTHORISATION

19 September 2023

10. DATE OF REVISION OF THE TEXT