

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S5

PAINOGESIC CO 37.5 mg/325 mg film-coated tablets

Tramadol hydrochloride 37,5 mg and paracetamol 325 mg.

Sugar Free

Contains: Sodium 0.035 mmol

Read all this leaflet carefully before you start taking

PAINOGESIC CO.

- Keep this leaflet. You may need to read it again.
- If you have further questions, please ask your doctor, pharmacist, nurse or other healthcare provider.
- **PAINOGESIC CO** has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet:

1. What **PAINOGESIC CO** is and what it is used for
2. What you need to know before you use **PAINOGESIC CO**
3. How to use **PAINOGESIC CO**
4. Possible side effects
5. How to store **PAINOGESIC CO**
6. Contents of the pack and other information

1. What PAINOGESIC CO is and what it is used for

PAINOGESIC CO contains tramadol and paracetamol.

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

Tramadol is a painkiller belonging to the class of the opioids that acts on the central nervous system. It is used for the management of moderate to moderately severe pain.

Paracetamol, the other active substance, has centrally acting analgesic effects.

2. What you need to know before you take PAINOGESIC CO

Do not take PAINOGESIC CO:

- If you are allergic to tramadol, paracetamol, any of the other ingredients mentioned in section 6.1, or other opioids such as codeine.
- If you are suffering from acute alcohol poisoning.
- If you are taking sleeping pills, pain relievers or medicines that affect mood and emotions.
- If you are also taking medicines such as tranylcypromine and moclobemide or have taken them in the last 14 days before treatment with **PAINOGESIC CO** which are used to treat depression. These medicines are used to treat depression and belong to a group called monoamine oxidase inhibitors (MAOIs).
- If you have a severe liver disorder.
- **PAINOGESIC CO** must not be used for narcotic (an addictive substance) withdrawal treatment.
- You should not take **PAINOGESIC CO** if you suffer from severe breathing problems.
- You should not take **PAINOGESIC CO** if you are suffering from increased intracranial pressure or central nervous system depression, which are characterised by decreased rate of breathing, decreased heart rate, and loss of consciousness.
- You should not take **PAINOGESIC CO** if you have fits that are not controlled

Warnings and precautions

Take special care with PAINOGESIC CO:

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

- If you take other medicines containing paracetamol or tramadol
- If you have liver problems or disease as your eyes and skin may turn yellow, which may suggest jaundice
- If you have kidney problems
- If you have recently suffered from a head injury, shock or severe headaches associated with vomiting (being sick)
- If you are dependent on any other medicines (for example morphine)
- If you have epilepsy or have experienced fits or seizures
- Tramadol exerts its effect by converting (metabolising) into its active component. If you convert (metabolise) Tramadol to this active component more rapidly and completely than other patients, you are known as an ultra-rapid metaboliser. This means you are more likely to have serious side effects, such as breathing difficulties, with slow or shallow breathing. If you experience these types of side effects, stop taking **PAINOGESIC CO** and consult with your doctor immediately.
- If you abruptly stop treatment with **PAINOGESIC CO** you may suffer from panic attacks, severe anxiety (feeling nervous or worried), hallucinations (hearing or seeing things that are not there), paraesthesia (burning or prickling sensation that is usually felt in the hands, arms, legs, or feet) and tinnitus (ringing or buzzing in the ears). Your doctor will gradually decrease your dose of **PAINOGESIC CO** to prevent these side effects from occurring.
- **PAINOGESIC CO** contains paracetamol. If you are allergic to paracetamol, you may break out in a severe skin rash after you take **PAINOGESIC CO**. You should stop using **PAINOGESIC CO** at the first appearance of a skin rash or any other sign of allergy.
- **PAINOGESIC CO** should not be taken with alcohol containing beverages.

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

- While taking **PAINOGESIC CO** you may experience low levels of sodium in your blood (hyponatraemia). Symptoms may include nausea and vomiting, headaches, feeling confused, feeling very tired, feeling restless, feeling irritable, muscle weakness, spasms or cramps and seizures. Elderly patients and patients taking other medicines that lower sodium in the blood are most at risk for this side effect. If you experience any of these symptoms while taking **PAINOGESIC CO**, consult your doctor immediately.
- If you suffer from sleep-related breathing disorders, take special care when taking **PAINOGESIC CO**.
- (SCAR) is a severe skin rash that is life-threatening.
Talk to your doctor if you develop a craving for **PAINOGESIC CO**

Other medicines and PAINOGESIC CO :

Always tell your healthcare professional if you are taking any other medicine. (This includes all complementary or traditional medicines).

PAINOGESIC CO should not be administered with the following:

- A group of antidepressants called Monoamine Oxidase (MAO) Inhibitors (such as moclobemide and tranylcypromine). When taken together with **PAINOGESIC CO** it can cause diarrhoea, tachycardia (irregular heartbeat), hyperhidrosis (excessive sweating), trembling, confusional state, even coma.
- Carbamazepine (used to treat seizures). When taken together with **PAINOGESIC CO** it can cause reduced efficacy and shorter duration due to decreased concentration of tramadol.
- buprenorphine, nalbuphine or pentazocine (opioid-type pain relievers). The effects of these medicines may be decreased.

Caution is advised when the following medicines are given with **PAINOGESIC CO**:

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

- Tramadol can induce convulsions and increase the potential for certain antidepressants such as citalopram, escitalopram, paroxetine, duloxetine, venlafaxine and amitriptyline, antipsychotics and seizure threshold- lowering medicinal products (such as bupropion, mirtazapine, tetrahydrocannabinol) to cause convulsions.
- Concomitant therapeutic use of tramadol and citalopram, escitalopram, paroxetine, duloxetine, venlafaxine and amitriptyline, moclobemide and tranylcypromine and mirtazapine may cause serotonin toxicity. Symptoms include involuntary, rhythmic contractions of muscles, including the muscles that control movement of the eye, agitation, excessive sweating, tremor, exaggeration of reflexes, increased muscle tension, body temperature above 38 °C.
- If you are taking other pain relievers such as morphine and codeine (also as cough medicine), baclofen (a muscle relaxant), medicines used to lower blood pressure, or medicines to treat allergies as well as sedative medicines such as benzodiazepines (alprazolam, diazepam, lorazepam) or related medicines with **PAINOGESIC CO** as there is an increased risk of drowsiness, difficulties in breathing (respiratory depression), coma which may be life-threatening.
- If you are taking warfarin (for blood thinning). The effectiveness of this medicine may be altered, and bleeding may occur.
- If you are taking digoxin used to treat heart problems
- If you are taking diflunisal used for pain
- If you are taking ondansetron used to treat vomiting

PAINOGESIC CO with food and drink

Do not drink alcohol during treatment with **PAINOGESIC CO** as the effects of **PAINOGESIC CO** and alcohol may intensify each other.

Pregnancy, breastfeeding and fertility

PAINOGESIC CO must not be taken during pregnancy.

PAINOGESIC CO must not be taken during breastfeeding since there is no information available regarding safety during breastfeeding.

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare professional for advice before taking this medicine.

Driving and using machines

Do not drive while taking this medicine until you know how it affects you.

PAINOGESIC CO can affect your ability to drive as it may make you sleepy or dizzy.

If you feel drowsy while taking **PAINOGESIC CO**, do not drive, use tools or use machinery.

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3. How to take PAINOGESIC CO

Do not share medicines prescribed for you with any other person.

Always take **PAINOGESIC CO** exactly as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure. Your doctor will tell you how long your treatment with **PAINOGESIC CO** will last. If you have the impression that the effect of **PAINOGESIC CO** is too strong or too weak, talk to your doctor or pharmacist.

- Swallow the tablets whole with sufficient liquid.
- Do not break or chew the tablets.

Adults:

Applicant/PHCR: *Innovata Pharmaceuticals (Pty) Ltd*

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: *Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg*

Application name (number): PAINOGESIC CO

The recommended dosage is 1 or 2 tablets every 4 to 6 hours, unless otherwise prescribed by your doctor. If required, further doses may be taken, as instructed by your doctor.

Do not take more than 8 tablets per day.

If you take more PAINOGESIC CO than you should

In the event of over dosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison center.

Immediate medical advice should be sought in the event of an overdose even if you feel well, because of the risk of delayed, serious liver damage from too much paracetamol. Your kidneys can also become damaged in the absence of liver damage.

If you forget to take PAINOGESIC CO

If you forgot to take a dose, take it as soon as you remember unless it is nearly time for your next dose. Do not take a double dose to make up for forgotten individual doses.

If you stop taking PAINOGESIC CO

Withdrawal symptoms may occur if **PAINOGESIC CO** is discontinued abruptly. You may experience anxiety, nervous excitement, difficulty sleeping, excessive abnormal movements, tremor and gastrointestinal symptoms.

Other symptoms that have very rarely been seen if tramadol hydrochloride is discontinued abruptly include: panic attacks, severe anxiety, hallucinations, tingling or prickling sensation and ringing in the ears.

4. Possible side effects

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

PAINOGESIC CO can have side effects.

Not all side effects reported for **PAINOGESIC CO** are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking **PAINOGESIC CO**, please consult your healthcare provider for advice.

If any of the following happens, stop taking **PAINOGESIC CO** and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Rash or itching all over the body
- Allergic reactions along with breathing problems or closing of the throat
- Increased bleeding time

These are all very serious side effects. If you have them, you may have had a serious reaction to **PAINOGESIC CO**. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following: The following side effects occur frequently:

- Irregular heartbeat that is much rapid and more pronounced than usual
- Elevated liver enzymes, resulting in abdominal pain, dark urine, pale stools , weakness, fatigue

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Frequent side effects:

- confusional state, sleep disorders, mood changes (anxiety, nervousness, feeling of high spirits)
- dizziness, excessive sleepiness
- headache, trembling

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

- nausea, vomiting, constipation, dry mouth, diarrhoea, stomach pain, indigestion, stomach gas

Less frequent side effects:

- depression, seeing, hearing, smelling, tasting or imagining things, nightmares
- involuntary muscular contractions, tingling or prickling sensation, loss of memory
- ringing in the ear
- rapid, strong, or irregular heartbeat, very fast heartbeat, dysrhythmia
- difficulty swallowing, blood in faeces
- chills
- chest pain
- problems with urination
- difficulty breathing
- increase in blood pressure, hot flushes
- low blood sugar
- drug dependence
- mental confusion and emotional disruption
- convulsions, fainting, speech disorders
- slurred speech, stumbling, convulsions, syncope, speech disorders.
- vision blurred, constriction of the pupil (miosis)
- excessive dilation of the pupils (mydriasis)
- severe skin rash

In addition, the following side effects have been reported by people using medicines that contain only tramadol or only paracetamol:

- feeling faint when getting up from a lying or sitting position, slow heart rate, fainting
- changes in appetite
- muscle weakness, slower or weaker breathing

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

- mood changes, changes in activity, changes in perception
- worsening of existing asthma
- Severe skin rash

If you notice any side effects not mentioned in this leaflet, inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects via

<http://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of this medicine.

5. How to store PAINOGESIC CO

PAINOGESIC CO should be stored in the original packaging (keep blisters in the carton until required for use).

Store at or below 30 °C.

Protect from light and moisture.

STORE ALL MEDICINE OUT OF REACH OF CHILDREN

Do not use after the expiry date stated on the label and carton.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g., toilets).

6. Contents of the pack and other information

What **PAINOGESIC CO** contains:

The active substance in **PAINOGESIC CO** is tramadol and paracetamol. Sugar free

Applicant/PHCR: Innovata Pharmaceuticals (Pty) Ltd

Product Proprietary Name: PAINOGESIC CO

Dosage Form & Strength: Film Coated Tablet, Tramadol HCl 37,5 mg & Paracetamol 325 mg

Application name (number): PAINOGESIC CO

The other ingredients are sodium starch glycolate, maize starch, pre-gelatinized starch, colloidal silicon dioxide, magnesium stearate, opary white Y-1-7000.

What PAINOGESIC CO looks like and contents of the pack

PAINOGESIC CO is a white to off-white, round, biconvex film coated tablet plain on both surfaces.

PAINOGESIC CO 37.5/325 mg is available in PVC/PVDC – Aluminium blister strip of 10 tablets per blister strip.

Carton (Duplex Cardboard paper):

20 tablets per carton, 30 tablets per carton, 60 tablets per carton and 100 tablets per carton.

NAME, BUSINESS ADDRESS AND TELEPHONE NUMBER OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION

Innovata Pharmaceuticals (Pty) LTD

Crownwood Office Park

100 Northern Parkway

Ormonde

Johannesburg

2091

South Africa

This leaflet was last revised in

26 June 2023

Registration number

A 49/2.9/0774