

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

S2

RUPASTRAL 10 mg tablets

Rupatadine

Contains sugar: lactose monohydrate 60 mg/tablet

Read all of this leaflet carefully because it contains important information for you

RUPASTRAL is available without a doctor's prescription, for you to treat a mild illness. Nevertheless, you still need to use RUPASTRAL carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share RUPASTRAL with any other person.
- Ask your health care provider or pharmacist if you need more information or advice.

What is in this leaflet

1. What RUPASTRAL is and what it is used for
2. What you need to know before you take RUPASTRAL
3. How to take RUPASTRAL
4. Possible side effects
5. How to store RUPASTRAL
6. Contents of the pack and other information

1. What RUPASTRAL is and what it is used for

RUPASTRAL is an antihistamine.

RUPASTRAL is used to relieve symptoms in adults and adolescents (over 12 years of age) of:

- Allergic rhinitis such as sneezing, runny nose, itching in the eyes and nose.
- Urticaria (an allergic skin rash) such as itching and hives (localised skin redness and swelling).

2. What you need to know before you take RUPASTRAL

Do not take RUPASTRAL:

- If you are hypersensitive (allergic) to RUPASTRAL or any of the other ingredients of RUPASTRAL (listed in section 6).
- If the patient is younger than 12 years of age.
- If you are pregnant or breastfeeding.

Warnings and precautions:

Take special care with RUPASTRAL:

- If you have, or have had liver or kidney disease, ask your doctor for advice.
The use of RUPASTRAL is not recommended in patients with impaired kidney or liver functions
- If you have heart conditions or low potassium blood levels, talk to your doctor or pharmacist, as RUPASTRAL can worsen these conditions
- If you are over the age of 65 years, as you can be more sensitive and can experience side effects.

Children

RUPASTRAL is not recommended for use in children under 12 years as safety is not established.

Other medicines and RUPASTRAL

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor or pharmacist if you are taking any of the following:

- Ketoconazole or erythromycin (medicines used to treat bacterial infections) as these medicines can increase the effect of RUPASTRAL.
- Alcohol.
- CNS depressants (medicines used to treat anxiety, panic attacks and acute stress reactions and sleep disorders).
- Statins (medicines to help lower cholesterol levels in the blood).
- Medicines used to treat bacterial infections, e.g. itraconazole, ketoconazole, voriconazole, posaconazole, erythromycin, fluconazole, clarithromycin.
- Medicines used to treat HIV, e.g. saquinavir, indinavir, ritonavir, nelfinavir, amprenavir, fosamprenavir,
- Nefazodone (medicines used to treat depression).
- Diltiazem (medicine used to treat certain heart conditions).

RUPASTRAL with food and drink

- Do not take RUPASTRAL with grapefruit juice.
- RUPASTRAL can be taken with or without meals.

Pregnancy and breastfeeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking this medicine.

It is not recommended to take RUPASTRAL when you are pregnant or breastfeeding.

Driving and using machines

It is not always possible to predict to what extent RUPASTRAL may interfere with your daily activities. At recommended doses, RUPASTRAL is not expected to influence your ability to drive or use machinery. However, RUPASTRAL may cause somnolence, headache, dizziness and a disturbance in attention; thus, when you first start taking RUPASTRAL you should take care to see how the treatment affects you before driving or using machinery.

RUPASTRAL contains lactose

Patients with the rare hereditary conditions of lactose intolerance should not take RUPASTRAL.

3. How to take RUPASTRAL

Do not share medicines prescribed for you with any other person.

Always take RUPASTRAL exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure.

RUPASTRAL is for adolescents (from 12 years and older) and adults.

The usual dose is one tablet per day.

Take the tablet at the same time of day, with a glass of water.

If you take more RUPASTRAL than you should

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to take RUPASTRAL

Do not take a double dose to make up for forgotten individual doses. If it is almost time for your next dose, skip the missed dose and take your next dose when you are meant to. Otherwise, take it as soon as you remember, and then go back to taking your tablets as you would normally.

If you are not sure whether to skip the dose, talk to your doctor or pharmacist.

4. Possible side effects

RUPASTRAL can have side effects.

Not all side effects reported for RUPASTRAL are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking RUPASTRAL, please consult your health care provider for advice.

If any of the following happens, stop taking RUPASTRAL and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Allergic (itching, hives and swelling of the face, lips, tongue or throat).

These are all very serious side effects. You may need urgent medical attention or hospitalisation.

Tell your doctor if you notice any of the following:

Frequent

- Sleepiness, headache, dizziness
- Dry mouth
- Sensation of weakness and fatigue.

Less frequent

- Sore throat, running nose
- Increased appetite
- Irritability
- Difficulty concentrating
- Rapid or regular heartbeat
- Nosebleed, nasal dryness, sore throat,
- Cough, dry throat, pain in tonsils area
- Nausea, stomach pain, diarrhoea, indigestion, vomiting, constipation
- Rash
- Back pain, joint pain, muscle pain
- Thirst, general discomfort, fever
- Changes in some laboratory blood tests.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. You can also report side effects to SAHPRA via the “6.04 Adverse Drug Reaction Reporting Form”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of RUPASTRAL.

5. How to store RUPASTRAL

Store all medicines out of reach of children.

- Store at or below 25 °C.
- Keep blister in original container to protect from light.
- Do not use after the expiry date stated on the blister or carton.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What RUPASTRAL contains

- The active substance is rupatadine.
- The other ingredients are lactose monohydrate, microcrystalline cellulose, pregelatinised maize starch, colour iron oxide red, colour iron oxide yellow, magnesium stearate.

What RUPASTRAL looks like and contents of the pack

RUPASTRAL tablets are light-salmon coloured, round, biconvex, uncoated tablets, plain on both sides.

It is packed in PVC/PVDC and aluminium foil. Each blister contains 10 tablets.

Blisters of 10 tablets are packed in a carton in pack sizes of 10, 20 or 30 tablets.

Holder of Certificate of Registration

Astral Pharma (Pty) Ltd

125 Meade Street

1st Floor, Beacon Place

George, 6529

This leaflet was last revised in

27 May 2025

Registration number

57/5.7.1/0199

A handwritten signature in black ink, appearing to be a stylized 'SK' or similar initials.