

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS

S4

WEGOVY® 0,25 mg (FlexTouch) solution for injection in pre-filled pen

WEGOVY® 0,5 mg (FlexTouch) solution for injection in pre-filled pen

WEGOVY® 1 mg (FlexTouch) solution for injection in pre-filled pen

WEGOVY® 1,7 mg (FlexTouch) solution for injection in pre-filled pen

WEGOVY® 2,4 mg (FlexTouch) solution for injection in pre-filled pen

Semaglutide

Sugar free.

Read all of this leaflet carefully before you start using WEGOVY®

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist, nurse or other health care provider.
- WEGOVY® has been prescribed for you personally and you should not share your medicine with other people. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What WEGOVY® is and what it is used for
2. What you need to know before you use WEGOVY®
3. How to use WEGOVY®
4. Possible side effects

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

5. How to store WEGOVY®
6. Contents of the pack and other information

1. What WEGOVY® is and what it is used for

What WEGOVY® is

WEGOVY® is a medicine for weight loss and weight maintenance that contains the active substance semaglutide. It is similar to a natural hormone called glucagon-like peptide-1 (GLP-1) that is released from the intestine after a meal. It works by acting on targets (receptors) in the brain that control your appetite, causing you to feel fuller and less hungry and experience less craving for food. This will help you eat less food and reduce your body weight.

What WEGOVY® is used for

WEGOVY® is used together with diet and physical activity for weight loss and to help keep the weight under control. It is used in adults, who have

- a BMI of 30 kg/m² or greater (obesity) or
- a BMI of at least 27 kg/m² but less than 30 kg/m² (overweight) who have weight-related health problems (such as diabetes, high blood pressure, abnormal levels of fats in the blood, breathing problems during sleep called 'obstructive sleep apnoea' or a history of heart attack, stroke or blood vessel problems).

BMI (Body Mass Index) is a measure of your weight in relation to your height.

WEGOVY® is used together with diet and physical activity for weight management in adolescents ages 12 years and above, who have

- obesity and
- body weight > 60 kg.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

As an adolescent patient, you should only continue using WEGOVY® if you have lost at least 5 % of your BMI after 12 weeks on the 2,4 mg dose or maximum tolerated dose (see section 3). Consult your doctor before you continue.

2. What you need to know before you use WEGOVY®

Do not use WEGOVY®

- if you are allergic to semaglutide or any of the other ingredients of this medicine (listed in section 6).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before using WEGOVY®.

The use of WEGOVY® is not recommended if you:

- use other products for weight loss,
- have type 1 diabetes,
- have severely reduced kidney function,
- have severely reduced liver function,
- have severe heart failure,
- have diabetic eye disease (retinopathy).

There is little experience with WEGOVY® in patients:

- of 75 years and older,
- with liver problems,
- with severe stomach or gut problem which results in delayed stomach emptying (called gastroparesis), or if you have an inflammatory bowel disease.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Please consult your doctor if one of the above applies to you.

- **Dehydration**

During treatment with WEGOVY®, you may feel sick (nausea) or be sick (vomiting) or have diarrhoea. These side effects can cause dehydration (loss of fluids). It is important that you drink enough fluids to prevent dehydration. This is especially important if you have kidney problems. Talk to your doctor if you have any questions or concerns.

- **Inflammation of the pancreas**

If you have severe and on-going pain in the stomach area (see section 4) – see a doctor straight away as this could be a sign of inflamed pancreas (acute pancreatitis).

- **People with type 2 diabetes**

WEGOVY® cannot be used as a substitute for insulin. Do not use WEGOVY® in combination with other medicines that contain GLP-1 receptor agonists (such as liraglutide, dulaglutide, exenatide or lixisenatide).

- **Low blood sugar (hypoglycaemia)**

Taking a sulfonylurea or an insulin with WEGOVY® might increase the risk of getting low blood sugar levels (hypoglycaemia). Please see section 4 for the warning signs of low blood sugar levels. Your doctor may ask you to test your blood sugar levels. This will help your doctor decide if the dose of the sulfonylurea or insulin needs to be changed to reduce the risk of low blood sugar.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

- **Diabetic eye disease (retinopathy)**

If you have diabetic eye disease and are using insulin, this medicine may lead to a worsening of your vision, and this may require treatment. Fast improvements in blood sugar control may lead to a temporary worsening of diabetic eye disease. If you have diabetic eye disease and experience eye problems while taking this medicine, talk to your doctor.

Children and adolescents

The safety and efficacy of WEGOVY® in children below 12 years of age have not been studied and are not recommended for use in this population.

Other medicines and WEGOVY®

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

In particular, tell your doctor, pharmacist, or nurse if you are using medicines containing the following:

- Warfarin or other similar medicines taken by mouth to reduce blood clotting (oral anti-coagulants). When you start treatment with e.g., warfarin or similar medicines, frequent blood testing to determine the ability of your blood to clot may be required.

Pregnancy and breast-feeding

This medicine should not be used during pregnancy, as it is not known if it may affect your unborn child. Therefore, it is recommended to use contraception while using this medicine. If you wish to become pregnant, you should stop using this medicine at least two months in advance. If you become or are pregnant, think you may be pregnant or are planning to have

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

a baby when using this medicine, talk to your doctor straight away, as your treatment will need to be stopped.

Do not use this medicine if you are breast-feeding, as it is unknown if it passes into breast milk.

Driving and using machines

WEGOVY® is unlikely to affect your ability to drive and use machines. Some patients may feel dizzy when taking WEGOVY® mainly during the first 4 months of treatment (see section 4). If you feel dizzy be extra careful while driving or using machines. If you need any further information, talk to your doctor, pharmacist or nurse.

People with type 2 diabetes

If you use this medicine in combination with a sulfonylurea or insulin, low blood sugar (hypoglycaemia) may occur which may reduce your ability to concentrate. Avoid driving or using machines if you get any signs of low blood sugar. See section 2, 'Warnings and precautions' for information on increased risk of low blood sugar and section 4 for the warning signs of low blood sugar. Talk to your doctor for further information.

WEGOVY® contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially 'sodium-free'.

3. How to use WEGOVY®

Do not share medicines prescribed for you with any other person.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Always use WEGOVY® exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to use

Your doctor will tell you how long your treatment with WEGOVY® will last. Do not stop treatment early; if you have the impression that the effect of WEGOVY® is too strong or too weak, tell your doctor or pharmacist.

The usual dose is:

Adults

The recommended dose is 2,4 mg once weekly.

Your treatment will start at a low dose which will be gradually increased over 16 weeks of treatment.

- When you first start using WEGOVY®, the starting dose is 0,25 mg once weekly.
- Your doctor will instruct you to gradually increase your dose every 4 weeks until you reach the recommended dose of 2,4 mg once weekly.
- Once you reach the recommended dose of 2,4 mg, do not increase this dose further.
- In case you are feeling very bothered by sickness (nausea) or by being sick (vomiting) talk with your doctor about delaying dose escalation or lowering to the previous dose until symptoms have improved.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Usually, you will be told to follow the table below.

Dose escalation	Weekly dose
Week 1 – 4	0,25 mg
Week 5 – 8	0,5 mg
Week 9 – 12	1 mg
Week 13 – 16	1,7 mg
From week 17	2,4 mg

Your doctor will assess your treatment on a regular basis.

Adolescents (above 12 years of age)

For adolescents, the same dose escalation schedule as for adults should be applied (see above). The dose should be increased until 2,4 mg (maintenance dose) or maximum tolerated dose has been reached. Weekly doses higher than 2,4 mg are not recommended.

How WEGOVY® is given

WEGOVY® is given as an injection under the skin (subcutaneous injection). Do not inject it into a vein or muscle.

- The best places to give the injection are the front of your upper arm, upper legs or stomach.
- Before you use the pen for the first time, your doctor, pharmacist, or nurse will show you how to use it.

Detailed instructions on how to use the pen are on the other side of this leaflet.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

People with type 2 diabetes

Tell your doctor if you have type 2 diabetes. Your doctor may adjust the dose of your diabetes medicines to prevent you from getting low blood sugar.

When to use WEGOVY®

- You should use this medicine once a week and if possible, on the same day each week.
- You can give yourself the injection at any time of the day – regardless of meals.

If necessary, you can change the day of your weekly injection of this medicine as long as it has been at least 3 days since your last injection. After selecting a new dosing day, continue with once-a-week dosing.

If you use more WEGOVY® than you should

You may get side effects such as feeling sick (nausea), being sick (vomiting) or have diarrhoea, which may cause dehydration (loss of fluids). In the event of overdose, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to use WEGOVY®

If you forgot to inject a dose and:

- it is 5 days or less since you should have used WEGOVY®, use it as soon as you remember. Then inject your next dose as usual on your scheduled day.
- it is more than 5 days since you should have used WEGOVY®, skip the missed dose. Then inject your next dose as usual on your next scheduled day.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Do not use a double dose to make up for a forgotten individual dose.

If you stop using WEGOVY®

Do not stop using this medicine without talking to your doctor.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

4. Possible side effects

WEGOVY® can have side effects. Not all side effects reported for WEGOVY® are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking WEGOVY®, please consult your health care provider for advice.

Serious side effects

Common (may affect up to 1 in 10 people)

- Complications of diabetic eye disease (diabetic retinopathy). If you have diabetes, you should inform your doctor if you experience eye problems, such as changes in vision, during treatment with this medicine.

Uncommon (may affect up to 1 in 100 people)

- Inflamed pancreas (acute pancreatitis). Signs of inflamed pancreas may include severe and long-lasting pain in your stomach, the pain may move to your back. You should see your doctor immediately if you experience such symptoms.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Rare (may affect up to 1 in 1,000 people)

- Severe allergic reactions (anaphylactic reactions, angioedema). You should seek immediate medical help and inform your doctor straight away if you get symptoms such as breathing difficulty, swelling, light-headedness, fast heartbeat, sweating and loss of consciousness or rapid swelling under the skin in areas such as the face, throat, arms and legs, which can be life threatening if throat swelling blocks the airway.

These are all very serious side effects. If you have them, you may have had a serious reaction to WEGOVY®. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

- chest pain,
- angina
- changes in the way your heart beats, for example, if you notice it beating faster,
- difficulty breathing,
- signs of recurrent infections such as fever or sore throat,
- less urine than is normal for you,
- yellowing of the skin and eyes, dark urine, and tiredness which may be symptoms of liver problems

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Other side effects

Very common (may affect more than 1 in 10 people)

- headache
- feeling sick (nausea)
- being sick (vomiting)
- diarrhoea
- constipation
- stomach pain
- feeling weak or tired

Common (may affect up to 1 in 10 people)

- feeling dizzy
- upset stomach or indigestion
- burping
- gas (flatulence)
- bloating of the stomach
- inflamed stomach ('gastritis') – the signs include stomach-ache, feeling sick (nausea) or being sick (vomiting)
- reflux or heartburn – also called 'gastro-oesophageal reflux disease'
- gallstones
- hair loss
- injection site reactions
- low blood sugar (hypoglycaemia) in patients with type 2 diabetes.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

The warning signs of low blood sugar may come on suddenly. They can include: cold sweat, cool pale skin, headache, fast heartbeat, feeling sick (nausea) or very hungry, changes in vision, feeling sleepy or weak, feeling nervous, anxious or confused, difficulty concentrating or shaking.

Your doctor will tell you how to treat low blood sugar and what to do if you notice these warning signs.

Low blood sugar is more likely to happen if you also take a sulfonylurea or insulin. Your doctor may reduce your dose of these medicines before you start using this medicine.

Uncommon (may affect up to 1 in 100 people)

- low blood pressure
- feeling dizzy or lightheaded on standing or sitting up because of a drop in blood pressure
- fast heartbeat
- increase of pancreatic enzymes (such as lipase and amylase) shown in blood tests.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the “**6.04 Adverse Drug Reaction Reporting Form**”, found online under SAHPRA's publications: <https://www.sahpra.org.za/Publications/Index/8>. By reporting side effects, you can help provide more information on the safety of WEGOVY®.

5. How to store WEGOVY®

Store all medicines out of reach of children.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Do not use this medicine after the expiry date which is stated on the pen label and carton after 'EXP'. The expiry date refers to the last day of that month.

Before opening

Store at or between 2 °C – 8 °C (in a refrigerator). Do not freeze. Keep away from the cooling element.

During use

- You can keep the pen for 6 weeks when stored at a temperature below 30 °C or in a refrigerator (2 °C – 8 °C) away from the cooling element. Do not freeze WEGOVY® and do not use it if it has been frozen.
- When you are not using the pen, keep the pen cap on in order to protect from light.

Do not use this medicine if you notice that the solution is not clear and colourless.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g., toilets).

6. Contents of the pack and other information

What WEGOVY® contains

The active substance is semaglutide.

WEGOVY® 0,25 mg FlexTouch® solution for injection

Each pre-filled pen contains 1 mg semaglutide in 1,5 mL (0,68 mg/mL).

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

WEGOVY® 0,5 mg FlexTouch® solution for injection

Each pre-filled pen contains 2 mg semaglutide in 1,5 mL (1,34 mg/mL).

WEGOVY® 1 mg FlexTouch® solution for injection

Each pre-filled pen contains 4 mg semaglutide in 3 mL (1,34 mg/mL).

WEGOVY® 1,7 mg FlexTouch® solution for injection

Each pre-filled pen contains 6,8 mg semaglutide in 3 mL (2,27 mg/mL).

WEGOVY® 2,4 mg FlexTouch® solution for injection

Each pre-filled pen contains 9,6 mg of semaglutide in 3 mL (3,2 mg/mL).

The other ingredients are disodium phosphate dihydrate, propylene glycol, phenol, hydrochloric acid/sodium hydroxide (for pH adjustment), water for injections. See also section 2 'WEGOVY® contains sodium' for information on sodium.

What WEGOVY® looks like and contents of the pack

WEGOVY® is a clear and colourless solution for injection in a pre-filled pen.

Each pre-filled pen contains 4 doses.

WEGOVY® 0,25, 0,5, 1 and 1,7 mg FlexTouch® solution for injection is available in the following pack size:

1 pre-filled pen and 4 disposable NovoFine® Plus needles.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

WEGOVY® 2,4 mg FlexTouch® solution for injection is available in the following pack sizes:

1 pre-filled pen and 4 disposable NovoFine® Plus needles.

3 pre-filled pens and 12 disposable NovoFine® Plus needles

Not all pack sizes may be marketed.

Holder of Certificate of Registration

Novo Nordisk (Pty) Ltd

150 Rivonia Road

10 Marion Street Office Park Building C1

Sandton

Johannesburg

2196

This leaflet was last revised in

N/A

Registration numbers

Wegovy® 0,25 mg: 58/21.013/0223

Wegovy® 0,5 mg: 58/21.013/0224

Wegovy® 1 mg: 58/21.013/0225

Wegovy® 1,7 mg: 58/21.013/0226

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Wegovy® 2,4 mg: 58/21.013/0227

Instructions on how to use WEGOVY®

Before you begin using your once-weekly WEGOVY® FlexTouch® pen, **always read these instructions carefully**, and talk to your doctor, nurse or pharmacist about how to inject WEGOVY® correctly.

WEGOVY® pen is a dial-a-dose pen that **contains four of your prescribed doses of WEGOVY®, corresponding to four times of once-weekly use.**

Please use the table inside the lid of the carton to keep track of how many injections you have used and how many doses remain in your pen.

WEGOVY® comes in five different pens, each containing one of the following prescribed doses of semaglutide:

0.25 mg

0.5 mg

1 mg

1.7 mg

2.4 mg

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Always start by checking your pen label to make sure that it contains your prescribed dose of WEGOVY®.

Your pen is designed to be used with 30 G, 31 G, and 32 G disposable needles up to a length of 8 mm.

The pack contains:


- WEGOVY® pen
- 4 NovoFine® Plus needles
- Patient Information Leaflet

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		


1

Wegovy FlexTouch pen (example)

Please note: Your pen may differ in size and your pen label may differ in colour from the example shown in the pictures.
These instructions apply to all Wegovy FlexTouch pens



NovoFine Plus needle (example)




1 Prepare your pen with a new needle

Check the name and dose of your pen to make sure it contains your prescribed dose of WEGOVY®.

Pull off the pen cap.

(See figure A).




Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Check that the solution in your pen is clear and colourless.

Look through the pen window. If WEGOVY® looks cloudy or coloured, do not use the pen.

(See figure B).




Always use a new needle for each injection.

Take a needle when you are ready to take your injection.

Check the paper tab and the outer needle cap for damages that could affect sterility. If any damage is seen, use a new needle.


Tear off the paper tab.

(See figure C).



Push the needle straight onto the pen. Turn until it is on tight.

(See figure D).



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

The needle is covered by two caps. You must remove both caps.


If you forget to remove both caps you will not inject any WEGOVY®.

Pull off the outer needle cap and keep it for later. You will need it to safely remove the needle from the pen after the injection.

Pull off the inner needle cap and dispose of it. A drop of WEGOVY® may appear at the needle tip. You must still check the WEGOVY® flow if you use a new pen for the first time. See **‘Check the flow with each new pen’**.

Never use a bent or damaged needle. For more information about needle handling, see **‘About your needles’** below these instructions.

(See figure E).



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Check the flow with each new pen

If your WEGOVY® pen is already in use, go to '**2 Set your dose**'.

Only check the WEGOVY® flow before your **first injection with each new pen**.


Turn the dose selector until you see the flow check symbol (■ ■ ▴).

(See figure F).



Make sure the flow check symbol lines up with the dose pointer.

(See figure G).



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Check the flow

Hold the pen with the needle pointing up.

Press and hold in the dose button until the dose counter returns to **0**. The **0** must line up with the dose pointer.

A drop of WEGOVY® should appear at the needle tip. This drop indicates that your pen is ready for use.

If a drop does not appear, check the flow again. **This should only be done twice.**

If there is still no drop, **change the needle and check the flow once more.**

Do not use the pen if a drop of WEGOVY® still does not appear.

(See figure H).



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

2 Set your dose

Turn the dose selector until the **dose counter stops**, and it **shows your prescribed dose**.


(See figure I).



The dashed line () in the dose counter will guide you to your dose.

The dose selector clicks differently when turned forward, backwards or past your dose. You will hear a 'click' every time you turn the dose selector. Do not set the dose by counting the number of clicks you hear.

(See figure J).



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		


When your prescribed dose lines up with the dose

pointer, you have selected your dose. In this picture, the

dose **0.25 mg** is shown as an example.

If the dose counter stops before you reach your prescribed dose, see the section ‘**Do you have enough WEGOVY®?**’ below these instructions.


(See figure K).



Choose your injection site

Choose your upper arms, upper legs or stomach (keep a 5 cm distance from your belly button).

You may inject in the same body area each week, but make sure it is not in the same spot as used the last time.




Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

3 Inject your dose

Insert the needle into your skin.

Make sure you can see the dose counter. Do not cover it with your fingers. This could interrupt the injection.

(See figure L).



Press and hold down the dose button until the dose counter shows **0.0**.


(See figure M).

Keep pressing the dose button with the needle in your skin and slowly count to 6. The **0.0** must line up with the dose pointer. You may hear or feel a click when the dose counter returns to **0.0**.



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

(See figure N).




Remove the needle from your skin. If the needle is removed earlier, a stream of WEGOVY® may come from the needle tip and the full dose will not be delivered.

If blood appears at the injection site, press lightly on the area to stop the bleeding.

You may see a drop of WEGOVY® at the needle tip after injecting. This is normal and does not affect your dose.

(See figure O).




Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

4 After your injection

Lead the needle tip into the outer needle cap on a flat surface without touching the needle or the outer needle cap.

Once the needle is covered, carefully push the outer needle cap completely on.

(See figure P).




Unscrew the needle and dispose of it carefully as instructed by your doctor, nurse, pharmacist or local authorities.

Never try to put the inner needle cap back on the needle.

You may stick yourself with the needle.

Always dispose of the needle immediately after each injection to prevent blocked needles, contamination, infection, and inaccurate dosing. **Never store your pen with the needle attached.**

(See figure Q).



Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

Put the pen cap on your pen after each use to protect

WEGOVY® from light.

(See figure R).



When the pen is empty, dispose of the pen without a needle on as instructed by your doctor, nurse, pharmacist, or local authorities.

The pen cap and the empty carton can be disposed of in your household waste.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

About your needles

How to identify a blocked or damaged needle

- If **0** does not appear in the dose counter after continuously pressing the dose button, you may have used a blocked or damaged needle.
- In this case, you have **not** received any WEGOVY® – even though the dose counter has moved from the original dose that you have set.

How to handle a blocked needle

- Change the needle as instructed in '**1 Prepare your pen with a new needle**' and go to '**2 Set your dose**'.

Caring for your pen

Treat your pen with care. Rough handling or misuse may cause inaccurate dosing. If this happens, you might not get the intended effect of WEGOVY®.


- See the back of this leaflet to read the storage conditions for your pen.
- **Do not inject WEGOVY® that has been exposed to direct sunlight.**
- **Do not subject WEGOVY® to frost and never inject WEGOVY® that has been frozen.** Dispose of the pen.
- **Do not drop your pen** or knock it against hard surfaces.
- **Do not try to refill your pen.** Once empty, it must be disposed of.
- **Do not try to repair your pen** or pull it apart.
- **Do not expose your pen to dust, dirt or liquid.**

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		

- **Do not wash, soak or lubricate your pen.** If necessary, clean it with a mild detergent on a moistened cloth.

Do you have enough WEGOVY®?

If the dose counter stops before you reach your prescribed dose, there is not enough WEGOVY® left for a full dose.
Dispose of the pen and use a new WEGOVY® pen.



Important information

- **Only inject one dose of WEGOVY® once weekly.** If you do not use your WEGOVY® as prescribed, you may not get the intended effect of this medicine.
- If you use more than one type of injectable medicine, it is very **important to check the name and dose** of your pen label **before use**.
- **Do not use this pen without help if you have poor eyesight and cannot follow these instructions.** Get help from a person with good eyesight who is trained to use the WEGOVY® pen.
- Always keep pen and needles **out of sight and reach of others, especially children**.
- **Never share** your pen or your needles with other people.
- **Needles are for single use only. Never reuse your needles** as it may lead to blocked needles, contamination, infection and inaccurate dosing.
- Caregivers must **be very careful when handling used needles** to prevent accidental needle stick injuries and infection.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Solution for injection; Semaglutide 0,25 mg Solution for injection; Semaglutide 0,5 mg Solution for injection; Semaglutide 1 mg Solution for injection; Semaglutide 1,7 mg Solution for injection; Semaglutide 2,4 mg
Product name	WEGOVY® 0,25 mg, 0,5 mg, 1 mg, 1,7 mg, 2,4 mg		