

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

SCHEDULING STATUS

S3

PROPRIETARY NAME AND DOSAGE FORM

JANUVIA® 25 Tablets

JANUVIA® 50 Tablets

JANUVIA® 100 Tablets

COMPOSITION

Each film-coated JANUVIA Tablet contains 32,13; 64,25 or 128,5 mg of sitagliptin phosphate monohydrate, which is equivalent to 25, 50 or 100 mg, respectively, of free base.

Excipients: anhydrous dibasic calcium phosphate, croscarmellose sodium, magnesium stearate, microcrystalline cellulose and sodium stearyl fumarate.

In addition, the film-coating contains the following inactive ingredients: polyethylene glycol (macrogol), polyvinyl alcohol, red iron oxide, talc, titanium dioxide and yellow iron oxide.

JANUVIA Tablets are sugar free.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

PHARMACOLOGICAL CLASSIFICATION

A.21.2 Oral Hypoglycaemics

PHARMACOLOGICAL ACTION

Pharmacodynamic properties

Sitagliptin is an orally-active inhibitor of the dipeptidyl peptidase 4 (DPP-4) enzyme. The DPP-4 inhibitors are a class of agents that act as incretin enhancers. By inhibiting the DPP-4 enzyme, sitagliptin increases the levels of two known active incretin hormones, glucagon-like peptide-1 (GLP-1) and glucose-dependent insulintropic peptide (GIP). Incretin hormones physiologically regulate blood glucose levels by increasing insulin response from pancreatic beta cells and suppressing glucagon secretion from pancreatic alpha cells when blood glucose levels are normal or elevated. These effects are not observed when blood glucose levels are low.

Cardiac electrophysiology

In a randomised, placebo-controlled crossover study, 79 healthy subjects were administered a single oral dose of sitagliptin 100 mg, sitagliptin 800 mg (8 times the recommended dose), and placebo. At the recommended dose of 100 mg, there was no effect on the QTc interval obtained at the peak plasma concentration, or at any other time during the study. Following the 800 mg dose, the maximum increase in the placebo-corrected mean change in QTc from baseline at 3 hours post-

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

dose was 8,0 msec. This increase was not considered to be clinically significant. At the 800 mg dose, peak sitagliptin plasma concentrations were approximately 11 times higher than the peak concentrations following a 100 mg dose.

In patients with type 2 diabetes administered sitagliptin 100 mg (n=81) or sitagliptin 200 mg (n=63) daily, there were no meaningful changes in QTc interval based on ECG data obtained at the time of expected peak plasma concentration.

Pharmacokinetic properties

The pharmacokinetics of sitagliptin have been characterised in healthy subjects and patients with type 2 diabetes. After oral administration of a 100 mg dose to healthy subjects, sitagliptin was absorbed, with peak plasma concentrations (median T_{max}) occurring 1 to 4 hours post-dose. Plasma AUC of sitagliptin increased in a dose-proportional manner. Following a single oral 100 mg dose to healthy volunteers, mean plasma AUC of sitagliptin was 8,52 micromolar hours, C_{max} was 950 nanomolar and apparent terminal half-life ($t_{1/2}$) was 12,4 hours. Plasma AUC of sitagliptin increased approximately 14 % following 100 mg doses at steady-state compared to the first dose. The intra-subject and inter-subject coefficients of variation for sitagliptin AUC were small (5,8 % and 15,1 %). The pharmacokinetics of sitagliptin were generally similar in healthy subjects and in patients with type 2 diabetes.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

Absorption

The absolute bioavailability of sitagliptin is approximately 87 %. Co-administration of a high-fat meal with sitagliptin had no effect on the pharmacokinetics (see **DOSAGE AND DIRECTIONS FOR USE**).

Distribution

The mean volume of distribution at steady-state following a single 100 mg intravenous dose of sitagliptin to healthy subjects is approximately 198 litres. The fraction of sitagliptin reversibly bound to plasma proteins is low (38 %).

Metabolism

Sitagliptin is primarily eliminated unchanged in urine (79 %), and metabolism is a minor pathway. Following a radioactively-labelled ¹⁴C sitagliptin oral dose, approximately 16 % of the radioactivity was excreted as metabolites of sitagliptin. Six metabolites were detected at trace levels and are not expected to contribute to the plasma DPP-4 inhibitory activity of sitagliptin. *In vitro* studies indicated that the primary enzyme responsible for the limited metabolism of sitagliptin was CYP3A4, with contribution from CYP2C8.

Elimination

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

Following administration of an oral radioactively-labelled ^{14}C sitagliptin dose to healthy subjects, approximately 100 % of the administered radioactivity was eliminated in faeces (13 %) or urine (87 %) within one week of dosing. The apparent terminal half-life ($t_{1/2}$) following a 100 mg oral dose of sitagliptin was approximately 12,4 hours and renal clearance was approximately 350 mL/min.

Elimination of sitagliptin occurs primarily via renal excretion and involves active tubular secretion. Sitagliptin is a substrate for human organic anion transporter-3 (hOAT-3), which may be involved in the renal elimination of sitagliptin. The clinical relevance of hOAT-3 in sitagliptin transport has not been established. Sitagliptin is also a substrate of p-glycoprotein, which may also be involved in mediating the renal elimination of sitagliptin. However, ciclosporin, a p-glycoprotein inhibitor, did not reduce the renal clearance of sitagliptin.

Characteristics in Patients

Renal insufficiency: A single-dose, open-label study was conducted to evaluate the pharmacokinetics of sitagliptin (50 mg dose) in patients with varying degrees of chronic renal insufficiency compared to normal healthy control subjects. The study included patients with renal insufficiency classified on the basis of creatinine clearance as mild (50 to < 80 mL/min), moderate (30 to < 50 mL/min) and severe (< 30 mL/min), as well as patients with end-stage renal disease on haemodialysis. Creatinine clearance was measured by 24-hour urinary creatinine clearance measurements or estimated from serum creatinine based on the Cockcroft-Gault formula:

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

$$\text{CrCl} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)} \{ \times 1,2 \}}{[\text{serum creatinine } (\mu\text{mol/L})]}$$

For female patients: 0,85 x value calculated for males

Compared to normal healthy control subjects, an approximate 1,6-fold increase in plasma AUC of sitagliptin was observed in patients with mild renal insufficiency. An approximately 2,3-fold increase in the plasma AUC of sitagliptin was observed in patients with moderate renal insufficiency; an approximately 3,8-fold increase was observed in patients with severe renal insufficiency and an approximately 4,5-fold increase was observed in patients with end-stage renal disease on haemodialysis, as compared to normal healthy control subjects. Sitagliptin was not meaningfully removed by haemodialysis (13,5 % over a 3- to 4-hour haemodialysis session starting 4 hours post-dose). To achieve plasma concentrations of sitagliptin similar to those in patients with normal renal function, lower dosages are recommended in patients with moderate and severe renal insufficiency, as well as in end-stage renal disease patients requiring haemodialysis (see **DOSAGE AND DIRECTIONS FOR USE, Patients with renal insufficiency**).

Hepatic insufficiency: In patients with moderate hepatic insufficiency (Child-Pugh score 7 to 9), mean AUC and C_{max} of sitagliptin increased approximately 21 % and 13 %, respectively, compared to healthy matched controls following administration of a single 100 mg dose of sitagliptin. These differences are not considered to be clinically meaningful.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

There is no clinical experience in patients with severe hepatic insufficiency (Child-Pugh score > 9) (see **CONTRAINDICATIONS**).

Elderly: Age did not have a clinically meaningful impact on the pharmacokinetics of sitagliptin, based on a population pharmacokinetic analysis of Phase I and Phase II data. Elderly subjects (65 to 80 years) had approximately 19 % higher plasma concentrations of sitagliptin compared to younger subjects.

Paediatric: No studies with sitagliptin have been performed in paediatric patients.

Race: Race had no clinically meaningful effect on the pharmacokinetics of sitagliptin based on a composite analysis of Phase I pharmacokinetic data and on a population pharmacokinetic analysis of Phase I and Phase II data, including subjects of White, Hispanic, Black, Asian and other racial groups.

Type 2 diabetes: The pharmacokinetics of sitagliptin in patients with type 2 diabetes are generally similar to those in healthy subjects.

INDICATIONS

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

Monotherapy

JANUVIA is indicated as an adjunct to diet and exercise to improve glycaemic control in adult patients with type 2 diabetes mellitus.

Combination therapy

JANUVIA is indicated in patients with type 2 diabetes mellitus to improve glycaemic control in combination with metformin, pioglitazone, a sulphonylurea or combinations thereof, or insulin when diet and exercise, plus the other therapies do not provide adequate glycaemic control.

CONTRAINDICATIONS

JANUVIA is contraindicated in patients with:

- hypersensitivity to sitagliptin, other gliptins (DPP-4 inhibitors) or to any components of JANUVIA or have a history of serious hypersensitivity reactions, such as anaphylaxis and angioedema to JANUVIA or other gliptins (DPP-4 inhibitors).

- type 1 diabetes mellitus

JANUVIA has not been studied in patients with severe hepatic insufficiency (see

Pharmacokinetic properties, Hepatic Insufficiency).

WARNINGS AND SPECIAL PRECAUTIONS

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

Pancreatitis: In post-marketing experience there have been reports of acute pancreatitis, including fatal and non-fatal haemorrhagic or necrotising pancreatitis (see **SIDE EFFECTS, Post-Marketing Experience**), in patients taking JANUVIA. Patients should be informed of the characteristic symptom of acute pancreatitis such as persistent, abdominal pain. Resolution of pancreatitis has been observed after discontinuation of JANUVIA. If pancreatitis is suspected, JANUVIA should be discontinued immediately.

Hypersensitivity Reactions: There have been post-marketing reports of serious hypersensitivity reactions in patients treated with JANUVIA. These reactions include anaphylaxis, angioedema and exfoliative skin conditions including Stevens-Johnson syndrome. Onset of these reactions usually occurred within the first 3 months after initiation of treatment with JANUVIA, with some reports occurring after the first dose. If a hypersensitivity reaction is suspected, discontinue JANUVIA immediately, assess for other potential causes for the event and institute an alternative class of medicines for treatment for diabetes (see **CONTRAINDICATIONS** and **SIDE EFFECTS, Post-Marketing Experience**).

Hypoglycaemia in combination with a sulphonylurea or with insulin: When JANUVIA was used in combination with a sulphonylurea or with insulin, medicines known to cause hypoglycaemia, the incidence of hypoglycaemia was increased over that of placebo (see **SIDE EFFECTS**). Therefore, to

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

reduce the risk of sulphonylurea- or insulin-induced hypoglycaemia, a lower dose of sulphonylurea or insulin may be considered (see **DOSAGE AND DIRECTIONS FOR USE**).

Renal insufficiency: A dosage adjustment is recommended in patients with moderate or severe renal insufficiency and in patients with end-stage renal disease requiring haemodialysis or peritoneal dialysis (see **DOSAGE AND DIRECTIONS FOR USE, Patients with renal insufficiency**).

There have been post-marketing reports of severe and disabling arthralgia in patients taking DPP-4 inhibitors such as JANUVIA. The time to onset of symptoms following initiation of drug therapy varied from one day to years. Patients experienced relief of symptoms upon discontinuation of the drug therapy. Some patients experienced a recurrence of symptoms when restarting JANUVIA or a different DPP-4 inhibitor. Consider DPP-4 inhibitors such as JANUVIA as a possible cause for severe joint pain and discontinue drug therapy if appropriate.

Effects on ability to drive and use machinery

No studies of the effects of JANUVIA on the ability to drive and use machines have been performed.

INTERACTIONS

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

In interaction studies, JANUVIA did not have clinically meaningful effects on the pharmacokinetics of the following: metformin, a thiazolidinedione (glitazone), glyburide, simvastatin, warfarin and oral contraceptives. Based on these data, JANUVIA does not inhibit CYP isozymes CYP3A4, 2C8 or 2C9. Based on *in vitro* data, sitagliptin is also not expected to inhibit CYP2D6, 1A2, 2C19 or 2B6 or to induce CYP3A4. There is limited information on multiple dose co-administration of these medicines.

There was an increase in the area under the curve (AUC, 11 %) and mean peak medicine concentration ($C_{\max}$, 18 %) of digoxin with the co-administration of sitagliptin. Patients receiving digoxin should be monitored appropriately.

The AUC and $C_{\max}$ of JANUVIA were increased approximately 29 % and 68 %, respectively, in subjects with co-administration of a single 100 mg oral dose of JANUVIA and a single 600 mg oral dose of ciclosporin, a potent probe inhibitor of p-glycoprotein. The observed changes in JANUVIA pharmacokinetics are not considered likely to be clinically meaningful. No dosage adjustment for JANUVIA is recommended when co-administered with ciclosporin or other p-glycoprotein inhibitors (e.g. ketoconazole).

HUMAN REPRODUCTION

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

There are no studies in pregnant women; therefore, JANUVIA is not recommended for use in pregnancy.

Lactation

JANUVIA is secreted in the milk of lactating rats. It is not known whether JANUVIA is secreted in human milk. Therefore, JANUVIA should not be used by women who are breastfeeding their infants.

DOSAGE AND DIRECTIONS FOR USE

JANUVIA can be taken with or without food.

The recommended dose of JANUVIA is 100 mg once daily as monotherapy or as combination therapy with metformin, a sulfonylurea, insulin (with or without metformin), a PPAR γ agonist, metformin plus a sulphonylurea or metformin plus a PPAR γ agonist.

When JANUVIA is used in combination with a sulphonylurea or with insulin, a lower dose of sulphonylurea or insulin may be considered to reduce the risk of hypoglycaemia (see **WARNINGS AND SPECIAL PRECAUTIONS, Hypoglycaemia in combination with a sulphonylurea or with insulin.**)

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

If a dose of JANUVIA is missed, it should be taken as soon as the patient remembers. A double dose of JANUVIA should not be taken on the same day.

Patients with renal insufficiency

For patients with mild renal insufficiency (creatinine clearance [CrCl] $\geq$ 50 mL/min, approximately corresponding to serum creatinine levels of $\leq$ 150 μ mol/L in men and $\leq$ 133 μ mol/L in women), no dosage adjustment for JANUVIA is required.

For patients with moderate renal insufficiency (CrCl $\geq$ 30 to $<$ 50 mL/min, approximately corresponding to serum creatinine levels of $>$ 150 μ mol/L to $\leq$ 265 μ mol/L in men and $>$ 133 μ mol/L to not $\leq$ 221 μ mol/L in women), the dose of JANUVIA is 50 mg once daily. This dose should be decreased if CrCl decreases to $<$ 30mL/min.

For patients with severe renal insufficiency (CrCl $<$ 30 mL/min, approximately corresponding to serum creatinine levels of $>$ 265 μ mol/L in men and $>$ 221 μ mol/L in women) or with end-stage renal disease requiring haemodialysis or peritoneal dialysis, the dose of JANUVIA is 25 mg once daily. JANUVIA may be administered without regard to the timing of dialysis.

Patients with hepatic insufficiency

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

No dosage adjustment is necessary for patients with mild to moderate hepatic insufficiency.

JANUVIA has not been studied in patients with severe hepatic insufficiency.

Elderly

No dosage adjustment is necessary for elderly patients.

Paediatric Population

There are no data available on the use of JANUVIA in patients younger than 18 years of age.

Therefore, use of JANUVIA in paediatric patients is not recommended.

SIDE EFFECTS

Adverse reactions considered as medicine related reported in patients treated with sitagliptin occurring in excess ($> 0,2\%$ and difference more than 1 patient) of that in patients treated with placebo are listed below by system organ class and frequency. Frequencies are defined as: very common ($\geq$ to $1/10$); common ($\geq$ to $1/100$, $< 1/10$); uncommon ($\geq 1/1\ 000$, $< 1/100$); rare ($\geq 1/10\ 000$, $< 1/1\ 000$); and very rare ($< 1/10\ 000$).

JANUVIA Monotherapy

Infections and Infestations

Common: upper respiratory tract infection, nasopharyngitis

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

Metabolism and nutrition disorders

Common: hypoglycaemia

Nervous system disorders

Common: headache

Uncommon: dizziness

Gastrointestinal disorders

Uncommon: constipation

Musculoskeletal and connective tissue disorders

Common: osteoarthritis, pain in extremity.

JANUVIA with metformin

Metabolism and nutrition disorders

Common: hypoglycaemia

Nervous system disorders

Uncommon: somnolence

Gastrointestinal disorders

Common: nausea, flatulence, vomiting

Uncommon: diarrhoea, constipation, upper abdominal pain

Investigations

Uncommon: blood glucose.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

JANUVIA with Sulphonylurea

Metabolism and nutrition disorders

Common: hypoglycaemia.

JANUVIA with a Sulphonylurea and Metformin

Metabolism and nutrition disorders

Common: hypoglycaemia

Respiratory, thoracic and mediastinal disorders

Uncommon: rhinorrhoea

Gastrointestinal disorders

Uncommon: nausea.

JANUVIA with a PPAR γ Agonist (pioglitazone)

Metabolism and nutrition disorders

Common: hypoglycaemia

Gastrointestinal disorders

Common: flatulence

General disorders and administration site conditions

Common: peripheral oedema

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

Investigations

Common: blood glucose.

JANUVIA with a PPAR γ Agonist (pioglitazone) and Metformin

Metabolism and nutrition disorders

Common: hypoglycaemia

General disorders and administration site conditions

Common: peripheral oedema.

JANUVIA with Insulin (+/-) Metformin)

Infections and Infestations

Common: influenza

Metabolism and nutrition disorders

Common: hypoglycaemia

Nervous system disorders

Common: headache

Gastrointestinal disorders

Uncommon: dry mouth, constipation.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

In addition, in monotherapy studies of up to 24 weeks in duration of JANUVIA 100 mg once daily alone compared to placebo, adverse reactions considered as medicine related reported in patients treated with JANUVIA in excess (> 0,2 % and difference more than 1 patient) of that in patients receiving placebo are headache, hypoglycaemia, constipation and dizziness.

In addition to the medicine-related adverse experiences described above, adverse experiences reported regardless of causal relationship to medication and occurring in at least 5 % and more commonly in patients treated with JANUVIA included upper respiratory tract infection and nasopharyngitis. Additional adverse experiences reported regardless of causal relationship to medication that occurred more frequently in patients treated with JANUVIA (not reaching the 5 % level but occurring with an incidence of greater than 0,5 % higher with JANUVIA than that in the control group) included osteoarthritis and pain in extremity.

Post-Marketing Experience

During post-marketing experience the following additional side effects have been reported: hypersensitivity reactions including anaphylaxis, angioedema, rash, urticaria, cutaneous vasculitis and exfoliative skin conditions including Stevens-Johnson syndrome (see **WARNINGS AND SPECIAL PRECAUTIONS**); acute pancreatitis, including fatal and non-fatal haemorrhagic and necrotising pancreatitis (see **WARNINGS AND SPECIAL PRECAUTIONS, Pancreatitis**); worsening renal function, including acute renal failure (sometimes requiring dialysis); vomiting,

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

headache, constipation, nasopharyngitis, upper respiratory tract infection; severe and disabling arthralgia (see **WARNINGS AND SPECIAL PRECAUTIONS**); myalgia; pain in extremity; back pain, pemphigoid, pruritus.

Laboratory investigations

Across clinical studies, a small mean increase in uric acid (approximately 12 µmol/L difference from placebo; mean baseline 297 to 327 µmol/L) was found in patients treated with JANUVIA 100 or 200 mg daily. No increase in the incidence of gout was reported. A small mean decrease in total alkaline phosphatase (up to approximately 5IU/l difference from placebo; mean baseline approximately 56 to 62 IU/L) was also observed, partly related to a small decrease in bone alkaline phosphatase. A small increase in white blood cell count (approximately 0,2 x 10⁹/L) difference in WBC versus placebo; mean baseline WBC approximately 6,6 x 10⁹/L was observed due to an increase in neutrophils.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT

In the event of an overdose it is reasonable to employ the usual supportive measures, e.g. remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring (including obtaining an electrocardiogram) and institute supportive therapy if required.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

JANUVIA is modestly dialysable in clinical studies, approximately 13,5 % of the dose was removed over a 3- to 4-hour haemodialysis session. Prolonged haemodialysis may be considered if clinically appropriate. It is not known if JANUVIA is dialysable by peritoneal dialysis.

IDENTIFICATION

JANUVIA 25: A pink, round, biconvex, film-coated tablet with 221 on one side and plain on the other.

JANUVIA 50: A light beige, round, biconvex, film-coated tablet with 112 on one side and plain on the other.

JANUVIA 100: A beige, round, biconvex, film-coated tablet with 277 on one side and plain on the other.

PRESENTATION

JANUVIA 25, 50 and 100 is packaged in opaque PVDC/PE/PVC push-through blisters (referred to as PVDC blister) with push-through aluminium lidding. Packs contain 28 tablets with blisters of 14 tablets. The blisters are opened by pushing the tablets through the foil.

STORAGE INSTRUCTIONS

Store at or below 30 °C.

Keep out of reach of children.

Product Name: JANUVIA 25, 50 and 100 mg	Component: English Professional Information Clinical approval: 17 July 2020 Date Approved: 14 September 2012
--	---

The blister should be kept in the carton until required for use.

REGISTRATION NUMBERS

JANUVIA 25: 41/21.2/0354

JANUVIA 50: 41/21.2/0355

JANUVIA 100: 41/21.2/0356

NAME AND ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION

MSD (Pty) Ltd

117 16th Road

Halfway House

1685

South Africa

DATE OF PUBLICATION OF THE PROFESSIONAL INFORMATION

The date of registration: 14 September 2012

The date of the most recent revision: 17 July 2020

WPC-MK0431-T-042012