

Doc Number: OF-CEM-POST-01C	CLINICAL APPROVAL LETTER: POST REGISTRATION APPLICATIONS	
Revision: 4.0		Effective date: 06 February 2026

Enquiries: variations@sahpra.org.za

CLINICAL APPROVAL LETTER: POST REGISTRATION APPLICATIONS

Clinical approval date	20/05/2026
Applicant's cover letter date	30/11/2025
Applicant	TEVA PHARMACEUTICALS (Pty) Ltd
Proprietary name(s) (including duplicates)	Filengia
API(s)	Fingolimod hydrochloride
Registration number(s)	52/34/0364
Sequence no	0007

Dear Sir/Madam,

Kindly be informed that the Authority has completed the evaluation of the above-mentioned Professional Information (PI) and Patient Information Leaflet (PIL) variation application.

1. The amendments to the PI and PIL are approved.
2. The PI and PIL be accepted as the currently approved PI and PIL
3. The date of revision reflected on this PI and PIL should be: **20/05/2026**
4. The final, dated version of the above PI and PIL must be submitted via docubridge in a closing sequence using the Submission Type "Closing sequence" and Sequence Description "Approved updated PI and PIL within 10 working days.

Note to applicant: Amendments to any quality aspects of the medicine within the PI and PIL are subject to P&A unit approval and may only be implemented in the PI if approved by the P&A unit.

Yours Faithfully,



L. DHLAMINI

MANAGER: CLINICAL POST-REGISTRATION UNIT

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

PROFESSIONAL INFORMATION

SCHEDULING STATUS:

S4

1. NAME OF THE MEDICINE:

Filengia, 0,5 mg (hard capsules)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION:

Each capsule contains 0,5 mg fingolimod (as hydrochloride).

Filengia is sugar free.

For the full list of excipients, see **section 6.1**.

3. PHARMACEUTICAL FORM:

Hard capsules.

Hard gelatin capsule (size 4), filled with white to off-white powder with small agglomerates. Capsule body: white opaque with TV 7820 imprinted. Capsule cap: yellow with TV 7820 imprinted.

4. CLINICAL PARTICULARS:

4.1 Therapeutic indications:

Filengia is indicated as a disease modifying therapy for the treatment of patients with relapsing multiple sclerosis to reduce the frequency of relapses and to delay the progression of disability.

4.2 Posology and method of administration:

Posology:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Do not exceed the recommended dosage.

The recommended dose of Filengia is one 0,5 mg capsule taken orally once daily, which can be taken with or without food. If a dose is missed treatment should be continued with the next dose as planned.

On initiation of Filengia treatment, after the first dose, all patients should be observed, with hourly pulse and blood pressure measurement, for a period of at least 6 hours for signs and symptoms of bradycardia. All patients should have an electrocardiogram performed prior to dosing and at the end of 6-hour monitoring period (see **section 4.4**).

Special populations:

Renal impairment:

No Filengia dose adjustments are needed in patients with renal impairment (see **section 5.2**).

Hepatic impairment:

No Filengia dose adjustments are needed in patients with mild to moderate hepatic impairment. Filengia must not be used in patients with severe hepatic impairment (Child-Pugh class C) (see **section 4.3**).

Paediatric patients:

Filengia is not indicated for use in paediatric patients (see **section 5.2**).

Elderly patients:

Filengia should be used with caution in patients aged 65 years and over (see **section 5.2**).

Diabetic patients:

Filengia should be used with caution in patients with diabetes mellitus due to a potential increased risk of macular oedema (see **section 4.4**).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Method of administration:

For oral use, taken once daily with or without food.

4.3 Contraindications:

- Hypersensitivity to fingolimod or to any of the excipients listed in **section 6.1**.
- Patients who in the last 6 months had myocardial infarction (MI), unstable angina pectoris, stroke/transient ischaemic attack (TIA), decompensated heart failure (requiring inpatient treatment), or New York Heart Association (NYHA) Class III/IV heart failure (see **section 4.4**).
- Patients with severe cardiac dysrhythmias requiring antidysrhythmic treatment with Class Ia (e.g., quinidine, procainamide, disopyramide) or Class III (e.g., amiodarone, sotalol) antidysrhythmic medicines (see **section 4.4**).
- Pregnancy and lactation (see **section 4.6**).
- Known immunodeficiency syndrome.
- Patients with increased risk for opportunistic infections, including immunocompromised patients (including those currently receiving immunosuppressive therapies or those immunocompromised by prior therapies).
- Suspected or confirmed progressive multifocal leukoencephalopathy (PML) (see **section 4.4**).
- Severe active infections, active chronic infections (hepatitis, tuberculosis).
- Known active malignancies.
- Severe liver impairment (Child-Pugh class C).
- Patients with second-degree Mobitz type II atrioventricular (AV) block or third-degree AV block, or sick-sinus syndrome, if they do not have a pacemaker (see **section 4.4**).
- Patients with a baseline QTc interval ≥ 500 msec (see **section 4.4**).
- Women of childbearing potential not using effective contraception (see **sections 4.4 and 4.6**).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

4.4 Special warnings and precautions for use:

Filengia should be used only by neurologists experienced in the treatment of multiple sclerosis (MS).

Filengia induces a reduction in heart rate upon treatment initiation, which can lead to bradydysrhythmia. The effect is usually maximal on Day 1, within the first 6 hours and heart rate usually normalises by 1 month. However, these events may occur at any time.

6-hour monitoring (ECG, heart rate and blood pressure) on Day 1 is mandatory for all patients, in order to determine individual response to treatment initiation. Patients who experience these events or patients with risk factors should have extended monitoring (at least overnight).

If patients develop signs or symptoms related to heart rate reduction, the monitoring should be extended until resolution of the event.

Bradydysrhythmia:

Initiation of Filengia treatment results in a transient decrease in heart rate and may also be associated with atrioventricular conduction delays, including the occurrence of isolated reports of transient, spontaneously resolving complete AV block (see **sections 4.8** and **5.1**).

After the first dose, the decline in heart rate starts within one hour, and is maximal within 6 hours and usually normalises by one month. However individual patients may not return to baseline heart rate by the end of the first month. Conduction abnormalities were typically transient and asymptomatic. They usually did not require treatment and resolved within the first 24 hours on treatment. If necessary, the decrease in heart rate induced by Filengia can be reversed by parenteral doses of atropine or isoprenaline.

All patients should have an ECG and blood pressure measurement performed prior to and 6 hours after the first dose of Filengia. All patients should be monitored for a period of 6 hours for signs and symptoms of bradycardia

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

with hourly heart rate and blood pressure measurement. Continuous (real time) ECG monitoring during this 6-hour period is recommended.

Should post-dose bradydysrhythmia-related symptoms occur, appropriate clinical management should be initiated, and monitoring should be continued until the symptoms have resolved. Should a patient require pharmacological intervention during the first-dose monitoring, overnight monitoring in a medical facility should be instituted and the first-dose monitoring should be repeated after the second dose of Filengia.

If the heart rate at 6 hours is the lowest since the first dose was administered (suggesting that the maximum pharmacodynamic effect on the heart may not yet be manifest), monitoring should be extended by at least 2 hours and until heart rate increases again. Additionally, if after 6 hours, the heart rate is < 45 bpm in adults, or the ECG shows new onset second degree or higher grade AV block or a QTc interval $\geq$ 500 msec, extended monitoring (at least overnight monitoring), should be performed, and until the findings have resolved. The occurrence at any time of third degree AV block should also lead to extended monitoring (at least overnight monitoring).

The effects on heart rate and atrioventricular conduction may recur on re-introduction of Filengia treatment depending on duration of the interruption and time since start of Filengia treatment. The same first dose monitoring as for treatment initiation is recommended when treatment is interrupted for:

- 1 day or more during the first 2 weeks of treatment.
- more than 7 days during weeks 3 and 4 of treatment.
- more than 2 weeks after one month of treatment.

If the treatment interruption is of shorter duration than the above, the treatment should be continued with the next dose as planned.

Cases of T-wave inversion have been reported in adult patients treated with fingolimod as contained in Filengia.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

In case of T-wave inversion, the prescriber should ensure that there are no associated myocardial ischaemia signs or symptoms. If myocardial ischaemia is suspected, it is recommended to seek advice from a cardiologist.

Due to the risk of serious rhythm disturbances or significant bradycardia, Filengia should not be used in patients with sino-atrial heart block, a history of symptomatic bradycardia, recurrent syncope or cardiac arrest, or in patients with significant QT prolongation (QTc > 470 msec (adult female), or > 450 msec (adult male)), uncontrolled hypertension or severe sleep apnoea (see also **section 4.3**). In such patients, if treatment with Filengia is considered, advice from a cardiologist should be sought prior to initiation of treatment in order to determine the most appropriate monitoring. At least overnight extended monitoring is recommended for treatment initiation (see also **section 4.5**).

Filengia has not been studied in patients with dysrhythmias requiring treatment with Class Ia (e.g., quinidine, procainamide, disopyramide) or Class III (e.g., amiodarone, sotalol) anti-dysrhythmic medicines. Class Ia and Class III anti-dysrhythmic medicines have been associated with cases of torsades de pointes in patients with bradycardia.

Since initiation of Filengia treatment results in decreased heart rate, Filengia should not be co-administered with these medicines (see **section 4.3**).

Experience with Filengia is limited in patients receiving concurrent therapy with beta blockers, heart-rate-lowering calcium channel blockers (such as verapamil or diltiazem), or other medicines which may decrease heart rate (e.g., ivabradine, digoxin, anticholinesteratic medicines or pilocarpine). Since the initiation of Filengia treatment is also associated with slowing of the heart rate (see **section 4.8**), concomitant use of these medicines during Filengia initiation may be associated with severe bradycardia and heart block. Because of the potential additive effect on heart rate, treatment with Filengia should not be initiated in patients who are concurrently treated with these medicines (see **section 4.5**). If treatment with Filengia is considered, advice from a cardiologist should be sought

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

regarding the switch to non-heart rate lowering medicines prior to initiation of treatment. If the heartrate lowering medication cannot be stopped, cardiologist's advice should be sought to determine appropriate first dose monitoring, at least overnight extended monitoring is recommended (see **section 4.5**).

QT interval:

In a thorough QT interval study of doses of 1,25 or 2,5 mg fingolimod at steady-state, when a negative chronotropic effect of fingolimod (as contained in Filengia) was still present, fingolimod treatment resulted in a prolongation of QTcI, with the upper limit of the 90 % CI $\leq 13,0$ ms. There is no dose- or exposure-response relationship of fingolimod and QTcI prolongation. There is no consistent signal of increased incidence of QTcI outliers, either absolute or change from baseline, associated with fingolimod treatment.

The clinical relevance of this finding is unknown. Reports have shown that from the multiple sclerosis studies, clinically relevant effects on prolongation of the QTc-interval have not been observed but patients at risk for QT prolongation were not included in clinical studies.

Medicines that may prolong QTc interval are best avoided in patients with relevant risk factors, for example, hypokalaemia, hypomagnesaemia or congenital QT prolongation.

Immunosuppressive effects:

Filengia has an immunosuppressive effect that predisposes patients to an infection risk, including opportunistic infections that can be fatal, and increases the risk of developing lymphomas and other malignancies, particularly those of the skin. Medical practitioners should carefully monitor patients, especially those with concurrent conditions or known factors, such as previous immunosuppressive therapy. If this risk is suspected, discontinuation of treatment should be considered by the medical practitioner on a case by-case basis (see also **section 4.4, Infections and Cutaneous neoplasms** and **section 4.8, Lymphomas**).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Infections:

Filengia causes a dose-dependent reduction of the peripheral lymphocyte count to 20 to 30 % of baseline values. This is due to the reversible sequestration of lymphocytes in lymphoid tissues (see **section 5.1**).

Before initiating treatment with Filengia, a recent complete blood count (CBC) (i.e., within 6 months or after discontinuation of prior therapy) should be available. Assessments of CBC are also recommended periodically during treatment, at month 3 and at least yearly thereafter, and in case of signs of infection. Absolute lymphocyte count $< 0,2 \times 10^9/L$, if confirmed, should lead to treatment interruption until recovery, because in clinical studies, fingolimod, as contained in Filengia, treatment was interrupted in patients with absolute lymphocyte count $< 0,2 \times 10^9/L$.

Initiation of treatment with Filengia should be delayed in patients with severe active infection until resolution. The immune system effects of Filengia may increase the risk of infections, including opportunistic infections (see **section 4.8**). Effective diagnostic and therapeutic strategies should be employed in patients with symptoms of infection while on therapy. When evaluating a patient with a suspected infection that could be serious, referral to a medical practitioner experienced in treating infections should be considered. During treatment, patients receiving Filengia should be instructed to report promptly symptoms of infection to their medical practitioner.

Suspension of Filengia should be considered if a patient develops a serious infection.

Elimination of fingolimod as contained in Filengia, following discontinuation of therapy may take up to two months and vigilance for infection should therefore be continued throughout this period. Patients should be instructed to report symptoms of infection up to 2 months after discontinuation of Filengia.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Herpes viral infection:

Serious, life-threatening, and sometimes fatal cases of encephalitis, meningitis or meningoencephalitis caused by herpes simplex and varicella zoster viruses have occurred with fingolimod as contained in Filengia, at any time during treatment. If herpes encephalitis, meningitis or meningoencephalitis occur, Filengia should be discontinued and appropriate treatment for the respective infection should be administered.

Patients need to be assessed for their immunity to varicella (chickenpox) prior to Filengia treatment. It is recommended that patients without a healthcare professional confirmed history of chickenpox or documentation of a full course of vaccination with varicella vaccine undergo antibody testing to varicella zoster virus (VZV) before initiating Filengia therapy. A full course of vaccination for antibody-negative patients with varicella vaccine is recommended prior to commencing treatment with Filengia (see **section 4.8**). Initiation of treatment with Filengia should be postponed for 1 month to allow full effect of vaccination to occur.

Cryptococcal meningitis:

Cases of cryptococcal meningitis (a fungal infection), sometimes fatal, have been reported in the post-marketing setting after approximately 2 to 3 years of treatment, although an exact relationship with the duration of treatment is unknown (see **section 4.8**). Patients with symptoms and signs consistent with cryptococcal meningitis (e.g., headache accompanied by mental changes such as confusion, hallucinations, and/or personality changes) should undergo prompt diagnostic evaluation. If cryptococcal meningitis is diagnosed, Filengia should be discontinued and appropriate treatment should be initiated. A multidisciplinary consultation (i.e., infectious disease specialist) should be undertaken if re-initiation of Filengia is warranted.

Progressive multifocal leukoencephalopathy:

Progressive multifocal leukoencephalopathy (PML) has been reported under fingolimod treatment since marketing authorisation (see **section 4.8**). PML is an opportunistic infection caused by John Cunningham virus (JCV), which may be fatal or result in severe disability. The majority of PML cases have occurred after 2 or more years of

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

fingolimod treatment. In addition to duration of fingolimod exposure, other potential risk factors for PML include prior therapy with immunosuppressants or immunomodulators, and/or severe lymphopenia ($<0.5 \times 10^9/l$). Patients at increased risk should be closely monitored for any signs or symptoms of PML.

PML can only occur in the presence of a JCV infection. If JCV testing is undertaken, it should be considered that the influence of lymphopenia on the accuracy of anti-JCV antibody testing has not been studied in fingolimod-treated patients. A negative anti-JCV antibody test does not preclude the possibility of subsequent JCV infection.

Before initiating treatment with Filengia, a baseline MRI should be available (usually within 3 months) as a reference. During routine MRI (in accordance with national and local recommendations), medical practitioners should pay attention to PML suggestive lesions. MRI findings may be apparent before clinical signs or symptoms. Annual MRIs may be considered as part of increased vigilance especially in patients at increased risk of PML. Cases of asymptomatic PML based on MRI findings and positive JCV DNA in the cerebrospinal fluid have been reported in patients treated with fingolimod, as contained in Filengia. If PML is suspected, MRI should be performed immediately for diagnostic purposes and treatment with Filengia should be discontinued until PML has been excluded. If PML is confirmed, treatment with fingolimod must be permanently discontinued (see **section 4.3**).

Immune reconstitution inflammatory syndrome (IRIS) has been reported in patients treated with sphingosine 1-phosphate (S1P) receptor modulators, including Filengia, who developed PML and subsequently discontinued treatment. IRIS presents as a clinical decline in the patient's condition that may be rapid, can lead to serious neurological complications or death, and is often associated with characteristic changes on MRI. The time to onset of IRIS in patients with PML was usually from weeks to months after S1P receptor modulator discontinuation. Monitoring for development of IRIS and appropriate treatment of the associated inflammation should be undertaken.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Human papilloma virus infection:

Human papilloma virus (HPV) infection, including papilloma, dysplasia, warts and HPV-related cancer, has been reported under treatment with fingolimod, as contained in Filengia, in the post-marketing setting (see **section 4.8**). Due to the immunosuppressive properties of Filengia, vaccination against HPV should be considered prior to treatment initiation with Filengia taking into account vaccination recommendations. Cancer screening, including Pap test, is recommended as per standard of care.

Vaccination:

Vaccination may be less effective during and for up to two months after stopping treatment with Filengia. The use of live attenuated vaccines may carry a risk of infections and should therefore be avoided (see **sections 4.5 and 4.8**).

Macular oedema:

Macular oedema with or without visual symptoms has been reported in 0,5 % of patients treated with fingolimod 0,5 mg, as contained in Filengia, occurring predominantly in the first 3 to 4 months of therapy (see **section 4.8**). An ophthalmological evaluation is therefore recommended at 3 to 4 months after treatment initiation. If patients report visual disturbances at any time while on therapy, evaluation of the fundus, including the macula, should be carried out.

Patients with history of uveitis and patients with diabetes mellitus are at increased risk of macular oedema (see **section 4.8**). Filengia has not been studied in multiple sclerosis patients with concomitant diabetes mellitus. It is recommended that multiple sclerosis patients with diabetes mellitus or a history of uveitis undergo an ophthalmological evaluation prior to initiating therapy and have follow-up evaluations while receiving Filengia therapy.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Continuation of Filengia in patients with macular oedema has not been evaluated. It is recommended that Filengia be discontinued if a patient develops macular oedema. A decision on whether or not Filengia therapy should be re-initiated after resolution of macular oedema needs to take into account the potential benefits and risks for the individual patient.

Liver injury:

Increased hepatic enzymes, mostly alanine aminotransaminase (ALT) but also gamma glutamyl transferase (GGT) and aspartate transaminase (AST) have been reported in multiple sclerosis patients treated with Filengia. Cases of acute liver failure requiring liver transplant and clinically significant liver injury have also been reported. Signs of liver injury, including markedly elevated serum hepatic enzymes and elevated total bilirubin, have occurred as early as ten days after the first dose and have also been reported after prolonged use. Elevations 3-fold the upper limit of normal (ULN) or greater in ALT occurred in 8,0 % of patients treated with fingolimod as contained in Filengia, and the medicine was discontinued if the elevation exceeded a 5-fold the ULN increase. Recurrence of liver transaminase elevations occurred with rechallenge in some patients, supporting a relationship to fingolimod, as contained in Filengia. Transaminase elevations occurred at any time during treatment although the majority occurred within the first 12 months. Serum transaminase levels returned to normal within approximately 2 months after discontinuation of fingolimod, as contained in Filengia.

Filengia has not been studied in patients with severe pre-existing hepatic injury (Child-Pugh class C) and should not be used in these patients.

Due to the immunosuppressive properties of Filengia, initiation of treatment should be delayed in patients with active viral hepatitis until resolution.

Recent (i.e., within last 6 months) transaminase and bilirubin levels should be available before initiation of

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

treatment with Filengia. In the absence of clinical symptoms, liver transaminases and serum bilirubin should be monitored at months 1, 3, 6, 9 and 12 on therapy and periodically thereafter until 2 months after Filengia discontinuation. In the absence of clinical symptoms, if liver transaminases are greater than 3 but less than 5 times the ULN without increase in serum bilirubin, more frequent monitoring including serum bilirubin and alkaline phosphatase (ALP) measurement should be instituted to determine if further increases occur and in order to discern if an alternative aetiology of hepatic dysfunction is present. If liver transaminases are at least 5 times the ULN or at least 3 times the ULN associated with any increase in serum bilirubin, Filengia should be discontinued. Hepatic monitoring should be continued. If serum levels return to normal (including if an alternative cause of the hepatic dysfunction is discovered), Filengia may be restarted based on a careful assessment of the patient.

Patients who develop symptoms suggestive of hepatic dysfunction, such as unexplained nausea, vomiting, abdominal pain, fatigue, anorexia, or jaundice and/or dark urine, should have liver enzymes and bilirubin checked and Filengia should be discontinued if significant liver injury is confirmed. Treatment should not be resumed unless a plausible alternative aetiology for the signs and symptoms of liver injury can be established.

Although there are no data to establish that patients with pre-existing liver disease are at increased risk of developing elevated liver function tests when taking Filengia, caution in the use of Filengia should be exercised in patients with a history of significant liver disease.

Interference with serological testing:

Since fingolimod, as contained in Filengia, reduces blood lymphocyte counts via re-distribution in secondary lymphoid organs, peripheral blood lymphocyte counts cannot be utilised to evaluate the lymphocyte subset status of a patient treated with Filengia. Laboratory tests involving the use of circulating mononuclear cells require larger blood volumes due to reduction in the number of circulating lymphocytes.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Blood pressure effects:

Patients with hypertension uncontrolled by medication were excluded from participation in premarketing clinical trials with fingolimod as contained in Filengia, and special care is indicated if patients with uncontrolled hypertension are treated with Filengia.

Hypertension was reported as an adverse event and therefore blood pressure should be regularly monitored during treatment with Filengia.

Respiratory effects:

Minor dose-dependent reductions in values for forced expiratory volume (FEV₁) and diffusion capacity for carbon monoxide (DLCO) were observed with fingolimod as contained in Filengia, starting at month 1 and remaining stable thereafter. Filengia should be used with caution in patients with severe respiratory disease, pulmonary fibrosis and chronic obstructive pulmonary disease (see **section 4.8**).

Posterior reversible encephalopathy syndrome:

Cases of posterior reversible encephalopathy syndrome (PRES) have been reported at the 0,5 mg dose (see **section 4.8**). Symptoms reported included sudden onset of severe headache, nausea, vomiting, altered mental status, visual disturbances and seizure. Symptoms of PRES are usually reversible but may evolve into ischaemic stroke or cerebral haemorrhage. Delay in diagnosis and treatment may lead to permanent neurological sequelae. If PRES is suspected, Filengia should be discontinued.

Prior treatment with immunosuppressive or immunomodulatory therapies:

There have been no studies performed to evaluate the efficacy and safety of fingolimod as contained in Filengia, when switching patients from teriflunomide, dimethyl fumarate or alemtuzumab treatment to Filengia. When switching patients from another disease modifying therapy to Filengia, the half-life and mode of action of the other

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

therapy must be considered in order to avoid an additive immune effect whilst at the same time minimising the risk of disease reactivation. A recent complete blood cell count (CBC) is recommended prior to initiating Filengia to ensure that immune effects of the previous therapy (i.e., cytopenia) have resolved.

Filengia can generally be started immediately after discontinuation of interferon or glatiramer acetate.

For dimethyl fumarate, the washout period should be sufficient for CBC to recover before treatment with Filengia is started.

Due to the long half-life of natalizumab, elimination usually takes up to 2 to 3 months following discontinuation. Teriflunomide is also eliminated slowly from the plasma. Without an accelerated elimination procedure, clearance of teriflunomide from plasma can take from several months up to 2 years. An accelerated elimination procedure as defined in the teriflunomide professional information is recommended or alternatively washout period should not be shorter than 3,5 months. Caution regarding potential concomitant immune effects is required when switching patients from natalizumab or teriflunomide to Filengia.

Alemtuzumab has profound and prolonged immunosuppressive effects. As the actual duration of these effects is unknown, initiating treatment with Filengia after alemtuzumab is not recommended.

A decision to use prolonged concomitant treatment with corticosteroids should be taken after careful consideration.

Co-administration with potent CYP450 inducers:

The combination of Filengia with potent CYP450 inducers should be used with caution. Concomitant administration with St John's wort is not recommended (see **section 4.5**).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Malignancies:

Cutaneous malignancies:

Basal cell carcinoma (BCC) and other cutaneous neoplasms, including malignant melanoma, squamous cell carcinoma, Kaposi's sarcoma and Merkel cell carcinoma, have been reported in patients receiving Filengia (see **section 4.8**).

Vigilance for skin lesions is warranted and a medical evaluation of the skin is recommended at initiation, and then every 6 to 12 months taking into consideration clinical judgement. The patient should be referred to a dermatologist in case suspicious lesions are detected.

Since there is a potential risk of malignant skin growths, patients treated with Filengia should be cautioned against exposure to sunlight without protection. These patients should not receive concomitant phototherapy with UV-B-radiation or PUVA-photochemotherapy.

Lymphomas:

There have been cases of lymphoma (see **section 4.8**). The cases reported were heterogeneous in nature, mainly non-Hodgkin's lymphoma, including B-cell and T-cell lymphomas. Cases of cutaneous T-cell lymphoma (mycosis fungoides) have been observed. A fatal case of Epstein-Barr virus (EBV) positive B-cell lymphoma has also been observed. If lymphoma is suspected, Filengia should be discontinued.

Women of childbearing potential:

Due to risk to the foetus, fingolimod, as contained in Filengia, is contraindicated during pregnancy and in women of childbearing potential not using effective contraception. Before initiation of treatment, women of childbearing potential must be informed of this risk to the foetus, must have a negative pregnancy test and must use effective contraception during treatment and for 2 months after treatment discontinuation (see **sections 4.3 and 4.6**).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Tumefactive lesions:

Cases of tumefactive lesions associated with MS relapse have been reported in the post-marketing setting. In case of severe relapses, MRI should be performed to exclude tumefactive lesions. Discontinuation of Filengia should be considered by the medical practitioner on a case-by-case basis taking into account individual benefits and risks.

Return of disease activity (rebound) after Filengia discontinuation:

Cases of severe exacerbation of disease have been reported after stopping fingolimod as contained in Filengia, in the post-marketing setting. This was generally observed within 12 weeks after stopping fingolimod as contained in Filengia, but was also reported up to and beyond 24 weeks after discontinuation. Caution is therefore indicated when stopping fingolimod therapy. If discontinuation of fingolimod is deemed necessary, the possibility of recurrence of exceptionally high disease activity should be considered and patients should be monitored for relevant signs and symptoms and appropriate treatment initiated as required (see **Stopping therapy** below).

Stopping therapy:

If a decision is made to stop treatment with Filengia a 6-week interval without therapy is needed, based on half-life, to clear fingolimod, as contained in Filengia, from the circulation (see **section 5.2**). Lymphocyte counts progressively return to normal range within 1 to 2 months of stopping therapy in most patients (see **section 5.1**) although full recovery can take significantly longer in some patients. Starting other therapies during this interval will result in concomitant exposure to Filengia. Use of immunosuppressants soon after the discontinuation of Filengia may lead to an additive effect on the immune system and caution is therefore indicated.

After stopping Filengia in the setting of PML, it is recommended to monitor patients for development of immune reconstitution inflammatory syndrome (PML-IRIS) (see **Progressive multifocal leukoencephalopathy** above).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Caution is also indicated when stopping Filengia therapy due to the risk of a rebound (see **Return of disease activity (rebound) after Filengia discontinuation** above). If discontinuation of Filengia is deemed necessary, patients should be monitored during this time for relevant signs of a possible rebound.

Paediatric population:

Filengia is not indicated for use in paediatric patients.

4.5 Interaction with other medicines and other forms of interaction:

Anti-neoplastic, immunomodulatory or immunosuppressive therapies:

Anti-neoplastic, immunomodulatory or immunosuppressive therapies should not be co-administered due to the risk of additive immune system effects (see **sections 4.3 and 4.4**).

Caution should also be exercised when switching patients from long-acting therapies with immune effects such as natalizumab, teriflunomide or mitoxantrone (see **section 4.4**). In multiple sclerosis clinical studies, the concomitant treatment of relapses with a short course of corticosteroids was not associated with an increased rate of infection.

Vaccination:

During and for up to two months after treatment with Filengia vaccination may be less effective. The use of live attenuated vaccines may carry a risk of infections and should therefore be avoided (see **sections 4.4 and 4.8**).

Bradycardia-inducing medicines:

Fingolimod, as contained in Filengia, has been studied in combination with atenolol and diltiazem. When fingolimod was used with atenolol in an interaction study in healthy volunteers, there was an additional 15 % reduction of heart rate at fingolimod, as contained in Filengia, treatment initiation, an effect not seen with diltiazem. Treatment with Filengia should not be initiated in patients receiving beta blockers, or other medicines which may decrease heart rate, such as Class Ia and III anti-dysrhythmics, calcium channel blockers (such as

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

verapamil or diltiazem), ivabradine, digoxin, anti-cholinesteratic medicines or pilocarpine because of the potential additive effects on heart rate (see **sections 4.4** and **4.8**). If treatment with Filengia is considered in such patients, advice from a cardiologist should be sought regarding the switch to non-heart rate lowering medicines or appropriate monitoring for treatment initiation, at least overnight monitoring is recommended, if the heart-rate-lowering medication cannot be stopped.

Pharmacokinetic interactions of other substances on fingolimod:

Filengia is metabolised mainly by CYP4F2. Other enzymes like CYP3A4 may also contribute to its metabolism, notably in the case of strong induction of CYP3A4. Potent inhibitors of transporter proteins are not expected to influence fingolimod, as contained in Filengia, disposition. Co-administration of fingolimod, as contained in Filengia, with ketoconazole resulted in a 1,7-fold increase in fingolimod and fingolimod phosphate exposure (AUC) by inhibition of CYP4F2. Caution should be exercised with substances that may inhibit CYP3A4 (protease inhibitors, azole antifungals, some macrolides such as clarithromycin or telithromycin).

Co-administration of carbamazepine 600 mg twice daily at steady-state and a single dose of fingolimod as contained in Filengia, 2 mg reduced the AUC of fingolimod and its metabolite by approximately 40 %. Other strong CYP3A4 enzyme inducers, for example rifampicin, phenobarbital, phenytoin, efavirenz and St. John's Wort, may reduce the AUC of fingolimod and its metabolite at least to this extent. As this could potentially impair the efficacy, their co-administration should be used with caution. Concomitant administration with St. John's Wort is however not recommended (see **section 4.4**).

Pharmacokinetic interactions of fingolimod on other medicines:

Fingolimod, as contained in Filengia, is unlikely to interact with substances mainly cleared by the CYP450 enzymes or by substrates of the main transporter proteins.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Co-administration of fingolimod, as contained in Filengia, with ciclosporin did not elicit any change in the ciclosporin or fingolimod exposure. Therefore, Filengia is not expected to alter the pharmacokinetics of medicines that are CYP3A4 substrates.

Co-administration of fingolimod, as contained in Filengia, with oral contraceptives (ethinylestradiol and levonorgestrel) did not elicit any change in oral contraceptive exposure. No interaction studies have been performed with oral contraceptives containing other progestogens, however an effect of fingolimod, as contained in Filengia, on their exposure is not expected.

4.6 Fertility, pregnancy and lactation:

Women of childbearing potential / Contraception in males and females:

Filengia is contraindicated in women of childbearing potential not using effective contraception (see **section 4.3**). Therefore, before initiation of Filengia treatment in women of childbearing potential, a negative pregnancy result must be available and counselling should be provided regarding the serious risk to the foetus. Women of childbearing potential must use effective contraception during treatment and for 2 months after discontinuation of Filengia since fingolimod takes approximately 2 months to eliminate from the body after treatment discontinuation (see **section 4.4**).

When stopping Filengia treatment for planning a pregnancy the possible return of disease activity should be considered (see **section 4.4**: Return of disease activity (rebound) after Filengia discontinuation and stopping therapy).

Male reproductive toxicity:

Fingolimod is present in seminal ejaculate.

Safety regarding an increased risk of male mediated foetal toxicity has not been demonstrated.

Pregnancy:

Filengia is contraindicated in pregnancy (see **section 4.3**). Safety in pregnancy has not been established.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Based on human experience, post-marketing data suggest that the use of fingolimod is associated with a 2-fold increased risk of major congenital malformation when administered during pregnancy compared with the general population. The following major malformations were most frequently reported:

- Congenital heart disease such as atrial and ventricular septal defects, tetralogy of Fallot
- Renal abnormalities
- Musculoskeletal abnormalities.

Fingolimod as contained in Filengia, is teratogenic in animals.

Breastfeeding:

Filengia is contraindicated in breastfeeding (see **section 4.3**). Fingolimod is excreted in the milk of treated animals during lactation. Due to the potential for serious adverse reactions to fingolimod in nursing infants, women receiving Filengia should not breastfeed.

Fertility:

Data from preclinical studies do not suggest that fingolimod would be associated with an increased risk of reduced fertility.

4.7 Effects on ability to drive and use machines:

Filengia has no or negligible influence on the ability to drive and use machines.

Dizziness or drowsiness may occasionally occur when initiating therapy with Filengia. On initiation of Filengia treatment it is recommended that patients be observed for a period of 6 hours (see **section 4.4, Bradydysrhythmia**).

4.8 Undesirable effects:

Summary of the safety profile:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

The most serious adverse reactions reported for the 0,5 mg recommended therapeutic dose were infections, macular oedema and transient atrio-ventricular blocks on treatment initiation.

The most frequent adverse reactions at the 0,5 mg dose were headache, influenza, diarrhoea, sinusitis, back pain, hepatic enzyme increased and cough. The most frequent reported adverse event for the 0,5 mg dose leading to treatment interruption was ALT elevations.

Tabulated list of adverse reactions:

MedDRA SOC	
Infections and infestations:	
<i>Frequent</i>	Influenza, sinusitis, Herpes viral infections, bronchitis, tinea versicolor
<i>Less frequent</i>	Pneumonia
<i>Unknown frequency</i>	Progressive multifocal leukoencephalopathy (PML)**, cryptococcal infections** Varicella Zoster Virus, John Cunningham Virus causing progressive multifocal leukoencephalopathy (PML), herpes simplex virus (HSV), bacterial (e.g., atypical mycobacterium).
Neoplasms benign, malignant and unspecified (incl. cysts and polyps):	
<i>Frequent</i>	Basal cell carcinoma
<i>Less frequent</i>	Malignant melanoma****, lymphoma***, squamous cell carcinoma****, Kaposi's sarcoma****
<i>Unknown frequency</i>	Merkel cell carcinoma***
Blood and lymphatic system disorders:	
<i>Frequent</i>	Lymphopenia, leukopenia
<i>Less frequent</i>	Thrombocytopenia
<i>Unknown frequency</i>	Autoimmune haemolytic anaemia***, peripheral oedema***
Immune system disorders:	

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

<i>Unknown frequency</i>	Hypersensitivity reactions, including rash, urticaria and angioedema upon treatment initiation**, immune reconstitution inflammatory syndrome (IRIS)
Psychiatric disorders:	
<i>Frequent</i>	Depression
<i>Less frequent</i>	Depressed mood
Nervous system disorders	
<i>Frequent</i>	Headache, dizziness, migraine
<i>Less frequent</i>	Seizure, posterior reversible encephalopathy syndrome (PRES)*
<i>Unknown frequency</i>	Severe exacerbation of disease after Filengia discontinuation***
Eye disorders:	
<i>Frequent</i>	Blurred vision
<i>Less frequent</i>	Macular oedema
Cardiac disorders:	
<i>Frequent</i>	Bradycardia, atrioventricular block
<i>Less frequent</i>	T-wave inversion**
Vascular disorders:	
<i>Frequent</i>	Hypertension
<i>Unknown frequency</i>	Ischaemic and haemorrhagic strokes
Respiratory, thoracic and mediastinal disorders:	
<i>Frequent</i>	Cough, dyspnoea
Gastrointestinal disorders:	
<i>Frequent</i>	Diarrhoea
<i>Less frequent</i>	Nausea**

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Skin and subcutaneous tissue disorders:	
<i>Frequent</i>	Eczema, alopecia, pruritus
Musculoskeletal and connective tissue disorders:	
<i>Frequent</i>	Back pain, myalgia, arthralgia
General disorders and administration site conditions:	
<i>Frequent</i>	Asthenia
Investigations:	
<i>Frequent</i>	Increased hepatic enzyme (increased alanine transaminase (ALT), gamma glutamyl transferase (GGT), aspartate transaminase (AST)), decreased weight***, increased blood triglycerides
<i>Less frequent</i>	Decreased neutrophil count
* The frequency category was based on an estimated exposure of approximately 10 000 patients to fingolimod in all clinical trials. ** PML and cryptococcal infections (including cases of cryptococcal meningitis) have been reported in the post-marketing setting (see section 4.4). *** Adverse drug reactions from spontaneous reports and literature. **** The frequency category and risk assessment were based on an estimated exposure of more than 24 000 patients to fingolimod 0,5 mg in all reported clinical trials.	

Description of selected adverse reactions:

Infections:

Lower respiratory tract infections, primarily bronchitis and to a lesser extent herpes infection and pneumonia were reported more frequently in fingolimod, as contained in Filengia, treated patients.

Some cases of disseminated herpes infection, including fatal cases, have been reported even at the 0,5 mg dose.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Cases of infections with opportunistic pathogens, such as viral (e.g., varicella zoster virus (VZV), John Cunningham virus (JCV) causing Progressive Multifocal Leukoencephalopathy, herpes simplex virus (HSV)), fungal (e.g., cryptococci including cryptococcal meningitis) or bacterial (e.g., atypical mycobacterium), have been reported, some of which have been fatal (see **section 4.4**).

Human papilloma virus (HPV) infection, including papilloma, dysplasia, warts and HPV-related cancer, has been reported under treatment with fingolimod, as contained in Filengia. Due to the immunosuppressive properties of fingolimod, as contained in Filengia, vaccination against HPV should be considered prior to treatment initiation with fingolimod, as contained in Filengia, taking into account vaccination recommendations. Cancer screening, including Pap test, is recommended as per standard of care.

Macular oedema:

Macular oedema has been reported at the recommended dose of 0,5 mg. The incidence of macular oedema increased at a higher dose of 1,25 mg. The majority of cases reported occurred within the first 3 to 4 months of therapy. Blurred vision or decreased visual acuity have been reported. Macular oedema has also been reported in asymptomatic patients during routine ophthalmological examination. Macular oedema generally improved or resolved spontaneously after discontinuation of treatment with fingolimod, as contained in Filengia. The risk of recurrence after re-challenge has not been evaluated.

A higher incidence of macular oedema has been reported in multiple sclerosis patients with a history of uveitis. The increased risk for macular oedema in patients with diabetes mellitus has not been studied (see **section 4.4**). A 2-fold increase in the incidence of macular oedema has been reported at a dose of 2,5 mg and 5 mg in renal transplant patients, including patients with diabetes mellitus.

Bradydysrhythmia:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

A transient decrease in heart rate which may also be associated with atrioventricular conduction delays has been reported at initiation of treatment with fingolimod, as contained in Filengia.

Maximal decline in heart rate has been reported within 6 hours after treatment initiation with fingolimod, as contained in Filengia, with declines in mean heart rate of 12 to 13 beats per minute. Heart rate below 40 beats per minute in adults, has been observed less frequently in patients treated with fingolimod, as contained in Filengia. The average heart rate returns towards baseline within 1 month of chronic treatment. Bradycardia is generally asymptomatic. Mild to moderate symptoms, including hypotension, dizziness, fatigue and/or palpitations have been reported. These symptoms resolve within 24 hours after initiation of treatment (see **sections 4.4** and **5.1**).

First-degree atrioventricular block (prolonged PR interval on ECG) has been detected and reported after treatment initiation in adult patients. Second-degree atrioventricular block has been reported in adults at a dose of 0,5 mg. Less frequent reports of transient, spontaneously resolving complete AV block have been observed during the six-hour monitoring period following the first dose of fingolimod, as contained in Filengia. The conduction abnormalities reported were typically transient, asymptomatic and resolved within the first 24 hours after treatment initiation. Although most patients do not require medical intervention, an asymptomatic second-degree Mobitz I atrioventricular block requiring medical intervention can occur in patients.

Delayed onset events including transient asystole and unexplained death have been reported within 24 hours of the first dose. These reported cases have been confounded by concomitant medications and/or pre-existing disease. The relationship of such events to fingolimod, as contained in Filengia, is uncertain.

Blood pressure:

It has been reported that fingolimod, as contained in Filengia, was associated with an average increase of approximately 3 mmHg in systolic pressure and approximately 1 mmHg in diastolic pressure, manifesting approximately 1 month after treatment initiation. Cases of hypertension have been reported within the first month of treatment initiation and on the first day of treatment that may require treatment with antihypertensive agents or

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

discontinuation of fingolimod, as contained in Filengia (see also **section 4.4, Blood pressure effects**).

Liver function:

Increased hepatic enzymes have been reported in multiple sclerosis patients treated with fingolimod, as contained in Filengia. Asymptomatic elevation in serum levels of ALT of $\geq 3 \times \text{ULN}$ (upper limit of normal) and $\geq 5 \times \text{ULN}$ has been reported. Recurrence of liver transaminase elevations has occurred upon re-challenge in some patients, supporting a relationship to the medicine. It has been reported that transaminase elevations occurred at any time during treatment although the majority occurred within the first 12 months. ALT levels returned to normal within approximately 2 months after discontinuation. ALT elevations $\geq 5 \times \text{ULN}$ returned to normal within approximately 5 months in patients who continued treatment with fingolimod, as contained in Filengia (see **section 4.4, Liver function**).

Nervous system disorders:

Reports have shown that rare events involving the nervous system occurred in patients treated with fingolimod, as contained in Filengia, at higher doses (1,25 or 5,0 mg) including ischaemic and haemorrhagic strokes and neurological atypical disorders, such as acute disseminated encephalomyelitis (ADEM)-like events.

Cases of seizures, including status epilepticus, have been reported with the use of fingolimod, as contained in Filengia.

Vascular disorders:

Cases of ischaemic and haemorrhagic strokes have been reported at the 0,5 mg dose. Cases of peripheral arterial occlusive disease were reported to have occurred in patients treated with fingolimod, as contained in Filengia, at higher doses (1,25 mg).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Respiratory system:

Minor dose-dependent reductions in values for forced expiratory volume (FEV₁) and diffusion capacity for carbon monoxide (DLCO) were observed with fingolimod, as contained in Filengia, treatment starting at month 1 and remaining stable thereafter. At month 24, a reduction from baseline values of predicted FEV₁ has been reported. The difference resolved after treatment discontinuation.

Lymphomas:

Cases of lymphoma of different varieties, including a fatal case of Epstein-Barr virus (EBV) positive B-cell lymphoma have been reported. Non-Hodgkin's lymphoma (B-cell and T-cell) cases, including cases of cutaneous T-cell lymphoma (mycosis fungoides) have been reported.

Haemophagocytic syndrome:

Cases of haemophagocytic syndrome (HPS) with fatal outcome have been reported in patients treated with fingolimod, as contained in Filengia, in the context of an infection. HPS is a condition that occur less frequently and has been described in association with infections, immunosuppression and a variety of autoimmune diseases.

Psychiatric disorders:

Depression and anxiety are known to occur with increased frequency in the multiple sclerosis population. Depression and anxiety have also been reported in patients on fingolimod, as contained in Filengia.

Reporting of suspected adverse reactions:

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Healthcare providers are asked to report any suspected adverse reactions to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

4.9 Overdose:

At 40 mg (i.e., 80-fold above the recommended dose) administered to healthy volunteers, 5 of 6 subjects reported mild chest tightness or discomfort which was clinically consistent with bronchoconstriction.

Filengia can induce bradycardia. The decline in heart rate usually starts within one hour of the first dose and is maximal within 6 hours. There have been reports of slow atrioventricular conduction with isolated reports of transient, spontaneously resolving complete AV block (see **sections 4.4** and **4.8**).

If the overdose constitutes first exposure to Filengia it is important to observe for signs and symptoms of bradycardia, which could include overnight monitoring. Regular measurements of pulse rate and blood pressure are required and electrocardiograms should be performed (see **sections 4.2** and **4.4**).

Neither dialysis nor plasma exchange would result in meaningful removal of Filengia from the body.

5. PHARMACOLOGICAL PROPERTIES:

5.1 Pharmacodynamic properties:

A 34 Other: selective immunosuppressive agents

Pharmacotherapeutic group: Immunosuppressants, selective immunosuppressants, ATC code: L04AA27

Mechanism of action:

Fingolimod is a sphingosine-1-phosphate receptor modulator. Fingolimod is metabolised by sphingosine kinase to the active metabolite fingolimod-phosphate.

Fingolimod-phosphate binds at low nanomolar concentrations to sphingosine-1-phosphate (S1P) receptors 1, 3,

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

and 4 located on lymphocytes, and readily crosses the blood brain barrier to bind to S1P receptors 1, 3, and 5 located on neural cells in the central nervous system. By acting as a functional antagonist of S1PR on lymphocytes, fingolimod-phosphate blocks the capacity of lymphocytes to egress from lymph nodes, causing a redistribution, rather than depletion, of lymphocytes. This redistribution reduces the infiltration of lymphocytes, including pro-inflammatory Th17 cells, into the central nervous system where they would be involved in nerve inflammation and nervous tissue damage.

Animal studies and *in vitro* experiments indicate that fingolimod may also exert beneficial effects in multiple sclerosis via interaction with S1P receptors on neural cells. Fingolimod penetrates the CNS, and has been shown in animals, to reduce astrogliosis, demyelination and neuronal loss. Further, fingolimod treatment increases the levels of brain derived neurotropic factor (BDNF) in the cortex, hippocampus and striatum of the brain of mice to support neuronal survival and improve motor functions.

Pharmacodynamic effects:

Immune system:

Within 4 to 6 hours after the first dose of fingolimod 0,5 mg, the lymphocyte count decreases to approximately 75 % of baseline. With continued daily dosing, the lymphocyte count continues to decrease over a two-week period, reaching a minimal count of approximately 500 cells/ μ L or approximately 30 % of baseline. Eighteen percent of patients reached a minimal count of < 200 cells/ μ L on at least one occasion. Low lymphocyte counts are maintained with chronic daily dosing. Peripheral lymphocyte count increases are evident within days of stopping fingolimod treatment and typically normal counts are reached within one to two months. Chronic fingolimod dosing leads to a decrease in the neutrophil count to approximately 80 % of baseline. Monocytes are unaffected by fingolimod.

Heart rate and rhythm:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Fingolimod causes a transient reduction in heart rate and decrease in atrioventricular conduction at treatment initiation (see **sections 4.4** and **4.8**). The maximal decline in heart rate is seen within 6 hours post dose, with 70 % of the negative chronotropic effect achieved on the first day. With continued administration heart rate returns to baseline within one month. The decrease in heart rate induced by fingolimod can be reversed by parenteral doses of atropine or isoprenaline. Inhaled salmeterol has also been shown to have a modest positive chronotropic effect. With initiation of fingolimod treatment there is an increase in atrial premature contractions, but there is no increased rate of atrial fibrillation/flutter or ventricular dysrhythmias or ectopy. Fingolimod treatment is not associated with a decrease in cardiac output. Autonomic responses of the heart, including diurnal variation of heart rate and response to exercise are not affected by fingolimod treatment.

S1P4 could partially contribute to the effect but was not the main receptor responsible for the lymphoid depletion. The mechanism of action of bradycardia and vasoconstriction were also studied in vitro in guinea pigs and isolated rabbit aorta and coronary artery. It was concluded that bradycardia could be mediated primarily by activation of inward rectifying potassium channel or G-protein activated inwardly rectifying K⁺ channel (IKACH/GIRK) and that vasoconstriction seems to be mediated by a Rho kinase and calcium dependent mechanism.

Potential to prolong the QT interval:

In a QT interval study of doses of 1,25 or 2,5 mg fingolimod at steady-state, when a negative chronotropic effect of fingolimod was still present, fingolimod treatment resulted in a mean prolongation of QTcI, with the upper boundary of the 90 % CI $\leq 13,0$ msec. There is no dose or exposure-response relationship of fingolimod and QTcI prolongation. There is no consistent signal of increased incidence of QTcI outliers, either absolute or change from baseline, associated with fingolimod treatment. In the multiple sclerosis studies, there were no clinically relevant prolongation of the QT interval.

Pulmonary function:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Fingolimod treatment with single or multiple doses of 0,5 and 1,25 mg for two weeks is not associated with a detectable increase in airway resistance as measured by FEV₁ and forced expiratory flow rate (FEF) 25 to 75. However, single fingolimod doses ≥ 5 mg (10-fold the recommended dose) are associated with a dose-dependent increase in airway resistance. Fingolimod treatment with multiple doses of 0,5, 1,25, or 5 mg is not associated with impaired oxygenation or oxygen desaturation with exercise or an increase in airway responsiveness to methacholine. Subjects on fingolimod treatment have a normal bronchodilator response to inhaled beta-agonists.

5.2 Pharmacokinetic properties:

Absorption:

Fingolimod absorption is slow ($t_{\max}$ of 12 to 16 hours) and extensive (≥ 85 %). The apparent absolute oral bioavailability is 93 % (95 % confidence interval: 79 to 111 %). Steady-state-blood concentrations are reached within 1 to 2 months following once-daily administration and steady-state levels are approximately 10-fold greater than with the initial dose.

Food intake does not alter $C_{\max}$ or exposure (AUC) of fingolimod. Fingolimod phosphate $C_{\max}$ was slightly decreased by 34 % but AUC was unchanged. Therefore, Filengia may be taken without regard to meals (see **section 4.2**).

Distribution:

Fingolimod highly distributes in red blood cells, with the fraction in blood cells of 86 %. Fingolimod phosphate has a smaller uptake in blood cells of < 17 %. Fingolimod and fingolimod phosphate are highly protein bound (> 99 %). Fingolimod and fingolimod-phosphate protein binding is not altered by renal or hepatic impairment. Fingolimod is extensively distributed to body tissues with a volume of distribution of about $1\,200 \pm 260$ litres. Fingolimod readily distributes into the brain and low levels are detected in seminal ejaculate.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Biotransformation:

The biotransformation of fingolimod in humans occurs by three main pathways; by reversible stereoselective phosphorylation to the pharmacologically active (S)-enantiomer of fingolimod-phosphate, by oxidative biotransformation catalysed mainly by CYP4F2 and possibly other CYP4F isoenzymes and subsequent fatty acid-like degradation to inactive metabolites, and by formation of pharmacologically inactive non-polar ceramide analogues of fingolimod.

Following single oral administration of [¹⁴C] fingolimod, the major fingolimod-related components in blood, as judged from their contribution to the AUC up to 816 hours post dose of total radiolabelled components, are fingolimod itself (23,3 %), fingolimod-phosphate (10,3 %), and inactive metabolites (M3 carboxylic acid metabolite (8,3 %), M29 ceramide metabolite (8,9 %) and M30 ceramide metabolite (7,3 %)).

Elimination:

Fingolimod blood clearance is $6,3 \pm 2,3$ L/h, and the average apparent terminal half-life ($t_{1/2}$) is 6 to 9 days. Blood levels of fingolimod and fingolimod phosphate decline in parallel in the terminal phase, leading to similar half-lives for both.

After an oral administration, about 81 % of the dose is slowly excreted in the urine as inactive metabolites. Fingolimod and fingolimod-phosphate are not excreted intact in urine but are the major components in the faeces with amounts representing less than 2,5 % of the dose each. After 34 days, the recovery of the administered dose is 89 %.

Linearity:

Fingolimod and fingolimod phosphate concentrations increase in an apparently dose proportional manner after multiple once-daily doses of 0,5 mg or 1,25 mg.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Characteristics in specific groups of subjects or patients:

The pharmacokinetics of fingolimod and fingolimod phosphate do not differ in males and females, in patients of different ethnic origin, or in patients with mild to severe renal impairment.

Renal dysfunction:

Severe renal impairment increases fingolimod C_{max} and AUC by 32 % and 43 %, respectively, and fingolimod-phosphate C_{max} and AUC by 25 % and 14 %, respectively. The apparent elimination half-life is unchanged for both analytes.

Hepatic dysfunction:

In subjects with mild, moderate, or severe hepatic impairment (Child-Pugh class A, B, and C), no change in fingolimod C_{max} was observed, but fingolimod AUC was increased respectively by 12 %, 44 %, and 103 %. In patients with severe hepatic impairment (Child-Pugh class C), fingolimod-phosphate C_{max} was decreased by 22 % and AUC was not substantially changed. The pharmacokinetics of fingolimod-phosphate were not evaluated in patients with mild or moderate hepatic impairment. The apparent elimination half-life of fingolimod is unchanged in subjects with mild hepatic impairment but is prolonged by about 50 % in patients with moderate or severe hepatic impairment.

Fingolimod should not be used in patients with severe hepatic impairment (Child-Pugh class C) (see **section 4.3**).

Fingolimod should be introduced cautiously in mild and moderate hepatic impaired patients (see **section 4.2**).

Elderly:

Clinical experience and pharmacokinetic information in patients aged above 65 years are limited. Filengia should be used with caution in patients aged 65 years and over (see **section 4.2**).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

Paediatric population:

Safety and efficacy of Filengia in paediatric patients below the age of 18 have not been studied. Filengia is not indicated for use in paediatric patients.

6. PHARMACEUTICAL PARTICULARS:

6.1 List of excipients:

Pregelatinised starch

Sodium laurilsulphate

Gelatin

Titanium dioxide (E171)

Yellow iron oxide (E172)

Printing ink:

Black iron oxide (E172)

Potassium hydroxide

Propylene glycol

Shellac

Strong ammonia solution

6.2 Incompatibilities:

Not applicable.

6.3 Shelf life:

2 years

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

6.4 Special precautions for storage:

Store at or below 25 °C.

Keep the capsule in the blister in the outer carton until required for use.

6.5 Nature and contents of container:

The capsules are packed into:

Aluminium – aluminium blister packs

Forming foil: Aluminium soft temper (OPA/Alum/PVC)

Lidding foil: Aluminium foil, heat sealable

Pack sizes: 7, 10, 28, 30, 98 in blisters.

or

Aluminium – aluminium blister packs (peel)

Forming foil: Aluminium soft temper (OPA/Alum/PVC)

Lidding foil: Aluminium foil, peelable (Paper/PET/Alum)

Pack sizes: 7x1, 10x1, 28x1, 30x1, 98x1 and 100x1 in perforated unit-dose blisters.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling:

Any unused medicine or waste material should be disposed of in accordance with local requirements.

7. HOLDER OF CERTIFICATE OF REGISTRATION:

Teva Pharmaceuticals (Pty) Ltd

Maxwell Office Park

Magwa Crescent West

Waterfall City, Midrand

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 27 November 2025 eCTD seq 0007

2090

8. REGISTRATION NUMBER(S):

52/34/0364

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION:

14 June 2022

10. DATE OF REVISION OF THE TEXT:

To be allocated.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S4

Filengia, 0,5 mg hard capsules

Fingolimod (as hydrochloride)

Filengia is sugar free

Read all of this leaflet carefully before you start taking Filengia

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor, pharmacist nurse or other healthcare provider.
- Filengia has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

WHAT IS IN THIS LEAFLET:

1. What Filengia is and what it is used for
2. What you need to know before you take Filengia
3. How to take Filengia
4. Possible side effects
5. How to store Filengia
6. Contents of the pack and other information

1. WHAT FILENGIA IS AND WHAT IT IS USED FOR:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Filingia contains the active substance fingolimod. Fingolimod is a selective immunosuppressant and is used in adults to treat relapsing (returning) multiple sclerosis (MS) (characterised by repeated attacks (relapses) of nervous system symptoms that reflect inflammation within the central nervous system.

Filingia does not cure MS, but it helps to reduce the number of relapses and to slow down the progression of physical disabilities due to MS.

2. WHAT YOU NEED TO KNOW BEFORE YOU TAKE FILENGIA:

Do not take Filingia:

- if you are hypersensitive (allergic) to fingolimod or any of the other ingredients of Filingia (listed in section 6).
- if you are taking or have recently taken medicine for irregular heartbeat such as quinidine, procainamide, disopyramide, amiodarone or sotalol.
- if, in the last 6 months, you have had heart attack, angina, stroke or warning of a stroke or certain types of heart failure.
- if you have certain types of irregular or abnormal heartbeat (dysrhythmia), including patients in whom the electrocardiogram (ECG) shows prolonged QT interval before starting Filingia.
- if you are pregnant or breastfeeding your baby.
- if you can bear children and are not using effective contraception
- if you have a lowered immune response (due to an immunodeficiency syndrome, a disease or to medicines that suppress the immune system).
- if your doctor suspects you may have a rare brain infection called progressive multifocal leukoencephalopathy (PML) or if PML has been confirmed.
- if you have a severe active infection or active chronic infection such as hepatitis or tuberculosis.
- if you have an active cancer.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

- if you have severe liver problems.

Warnings and precautions:

Take special care with Filengia:

Tell your doctor before you start taking Filengia:

- if you have been told you have an abnormal electrocardiogram.
- if you suffer from symptoms of slow heart rate (e.g., dizziness, nausea, or palpitations).
- if you have a history of sudden loss of consciousness or fainting (syncope).
- if you have severe breathing problems during sleep (severe sleep apnoea).
- if you are taking or have recently taken medicines that slow your heart rate (beta blockers (such as atenolol), calcium channel blockers (such as verapamil, diltiazem), ivabradine, digoxin, anticholinesteratic medicines or pilocarpine)
- if you have low levels of potassium or magnesium in your blood as this can increase your risk for heart rhythm disorder.
- if you plan to get vaccinated.
- if you have never had chickenpox or have not been vaccinated against varicella zoster virus.
- if you have or have had visual disturbances or other signs of swelling in the central vision area (macula) at the back of the eye (a condition known as macular oedema, see below), inflammation or infection of the eye (uveitis), or if you have diabetes (which can cause eye problems).
- if you have liver problems.
- if you have high blood pressure that cannot be controlled by medicines.
- if you have severe lung problems.

Slow heart rate (bradycardia) and irregular heartbeat:

Filengia causes the heart rate to slow down. As a result, you may feel dizzy or tired, or be consciously aware of

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

your heartbeat, or your blood pressure may drop. If these effects are pronounced, tell your doctor, because you may need treatment right away. Filengia can also cause an irregular heartbeat, especially after the first dose. Irregular heartbeat usually returns to normal in less than one day. Slow heart rate usually returns to normal within one month.

Your doctor will ask you to stay at the surgery or clinic for at least 6 hours, with hourly pulse and blood pressure measurements, after taking the first dose of Filengia so that appropriate measures can be taken in the event of side effects that occur at the start of treatment. You should have an electrocardiogram performed prior to the first dose of Filengia and after the 6-hour monitoring period. Your doctor may monitor your electrocardiogram continuously during that time. If after the 6-hour period you have a very slow or decreasing heart rate, or if your electrocardiogram shows abnormalities, you may need to be monitored for a longer period (at least 2 more hours and possibly overnight) until these have resolved. The same may apply if you are resuming Filengia after a break in treatment, depending on both how long the break was and how long you had been taking Filengia before the break.

If you have, or if you are at risk for, an irregular or abnormal heartbeat, if your electrocardiogram is abnormal, or if you have heart disease or heart failure, Filengia may not be appropriate for you.

If you have a history of sudden loss of consciousness or decreased heart rate, Filengia may not be appropriate for you. You will be evaluated by a cardiologist (heart specialist) to advise how you should start treatment with Filengia, including overnight monitoring.

If you are taking medicines that can cause your heart rate to decrease, Filengia may not be appropriate for you. You will need to be evaluated by a cardiologist, who will check whether you can be switched to alternative medicine that does not decrease your heart rate in order to allow treatment with Filengia. If such a switch is

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

impossible, the cardiologist will advise how you should start treatment with Filengia, including overnight monitoring.

If you have never had chickenpox or have not been vaccinated against varicella zoster virus:

If you have never had chickenpox, your doctor will check your immunity against the virus that causes it (varicella zoster virus). If you are not protected against the virus, you may need a vaccination before you start treatment with Filengia. If this is the case, your doctor will delay the start of treatment with Filengia until one month after the full course of vaccination is completed.

Infections:

Filengia lowers the white blood cell count (particularly the lymphocyte count). White blood cells fight infection. While you are taking Filengia (and for up to 2 months after you stop taking it), you may get infections more easily. Any infection that you already have may get worse. Infections could be serious and life-threatening. If you think you have an infection, have fever, feel like you have the flu, or have a headache accompanied by stiff neck, sensitivity to light, nausea, and/or confusion (these may be caused by a fungal infection and may be symptoms of meningitis), contact your doctor straight away, because it could be serious and life-threatening. If you believe your MS is getting worse (e.g., weakness or visual changes) or if you notice any new symptoms, talk to your doctor straight away, because these may be the symptoms of a brain disorder caused by infection and called progressive multifocal leukoencephalopathy (PML). PML is a serious condition that may lead to severe disability or death.

Human papilloma virus (HPV) infection, including papilloma, dysplasia, warts and HPV-related cancer, has been reported in patients treated with Filengia. Your doctor will consider whether you need to have a vaccination against HPV before starting treatment. If you are a woman, your doctor will also recommend HPV screening.

PML:

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

PML is a rare brain disorder caused by an infection that may lead to sever disability or death. Your doctor will arrange magnetic resonance imaging (MRI) scans before you start treatment and during treatment to monitor the risk of PML.

If you believe your MS is getting worse or if you notice any new symptoms, for example changes in mood or behaviour, new or worsening weakness on one side of the body, changes in vision, confusion., memory lapses or speech and communication difficulties, talk to our doctor as soon as possible. These may be symptoms of PML. Also speak with your partner or caregivers and inform them about your treatment. Symptoms might arise that you might not become aware of by yourself.

If you get PML it can be treated and your treatment with Filengia will be stopped. Some people get an inflammatory reaction as Filengia is removed from the body. This reaction (known as immune reconstitution inflammatory syndrome or IRIS) may lead to your condition getting worse, including worsening of brain function.

Macular oedema:

Before you start Filengia, if you have or have had visual disturbances or other signs of swelling in the central vision area (macula) at the back of the eye, inflammation or infection of the eye (uveitis) or diabetes, your doctor may want you to undergo an eye examination.

Your doctor may want you to undergo an eye examination 3 to 4 months after starting Filengia treatment.

The macula is a small area of the retina at the back of the eye which enables you to see shapes, colours, and details clearly and sharply. Filengia may cause swelling in the macula, a condition that is known as macular oedema. The swelling usually happens in the first 4 months of Filengia treatment.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Your chance of developing macular oedema is higher if you have diabetes or have had an inflammation of the eye called uveitis. In these cases, your doctor will want you to undergo regular eye examinations in order to detect macular oedema.

If you have had macular oedema, talk to your doctor before you resume treatment with Filengia.

Macular oedema can cause some of the same vision symptoms as an MS attack (optic neuritis). Early on, there may not be any symptoms. Be sure to tell your doctor about any changes in your vision. Your doctor may want you to undergo an eye examination, especially if:

- the centre of your vision gets blurry or has shadows;
- you develop a blind spot in the centre of your vision;
- you have problems seeing colours or fine detail.

Liver function tests:

If you have severe liver problems, you should not take Filengia. Filengia may affect your liver function. You will probably not notice any symptoms but if you notice yellowing of your skin or the whites of your eyes, abnormally dark urine, unexplained nausea and vomiting, abdominal pain, anorexia or fatigue, tell your doctor straight away.

If you get any of these symptoms after starting Filengia, tell your doctor straight away.

Before, during and after treatment with Filengia your doctor will request blood tests to monitor your liver function.

If your test results indicate a problem with your liver, you may have to interrupt treatment with Filengia.

High blood pressure:

As Filengia causes a slight elevation of blood pressure, your doctor may want to check your blood pressure

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

regularly.

Lung problems:

Filengia has a slight effect on the lung function. Patients with severe lung problems or with smoker's cough may have a higher chance of developing side effects.

Posterior reversible encephalopathy syndrome (PRES):

A condition called posterior reversible encephalopathy syndrome (PRES) has been reported in MS patients treated with Filengia. Symptoms may include sudden onset of severe headache, confusion, seizures and vision changes. Tell your doctor straight away if you experience any of these symptoms during your treatment with Filengia, because it could be serious.

Blood count:

The desired effect of Filengia treatment is to reduce the amount of white blood cells in your blood. This will usually go back to normal within 2 months of stopping treatment. If you need to have any blood tests, tell the doctor that you are taking Filengia. Otherwise, it may not be possible for the doctor to understand the results of the test, and for certain types of blood test your doctor may need to take more blood than usual.

Before you start Filengia, your doctor will confirm whether you have enough white blood cells in your blood and may want to repeat a check regularly. In case you do not have enough white blood cells, you may have to interrupt treatment with Filengia.

Skin cancer:

Skin cancers have been reported in MS patients treated with Filengia. Talk to your doctor straight away if you notice any skin nodules (e.g., shiny pearly nodules), patches or open sores that do not heal within weeks. Symptoms

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

of skin cancer may include abnormal growth or changes of skin tissue (e.g., unusual moles) with a change in colour, shape or size over time. Before you start Filengia, a skin examination is required to check whether you have any skin nodules. Your doctor will also carry out regular skin examinations during your treatment with Filengia. If you develop problems with your skin, your doctor may refer you to a dermatologist, who after consultation may decide that it is important for you to be seen on a regular basis.

Exposure to the sun and protection against the sun:

Fingolimod weakens your immune system. This increases your risk of developing cancers, in particular skin cancers. You should limit your exposure to the sun and

UV rays by:

- wearing appropriate protective clothing.
- regularly applying sunscreen with a high degree of UV protection.

Unusual brain lesions associated with MS relapse:

Some cases of unusually large brain lesions associated with MS relapse have been reported in patients treated with Filengia. In case of severe relapse, your doctor will consider performing MRI to evaluate this condition and will decide whether you need to stop taking Filengia.

Switch from other treatments to Filengia:

Your doctor may switch you directly from beta interferon, glatiramer acetate or dimethyl fumarate to Filengia if there are no signs of abnormalities caused by your previous treatment. Your doctor may have to do a blood test in order to exclude such abnormalities. After stopping natalizumab you may have to wait for 2-3 months before starting treatment with Filengia. To switch from teriflunomide, your doctor may advise you to wait for a certain time or to go through an accelerated elimination procedure. If you have been treated with alemtuzumab, a thorough evaluation and discussion with your doctor is required to decide if Filengia is appropriate for you. Should you

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

require combined treatment with a corticosteroid and Filengia, your doctor will need to carefully consider the concomitant treatment if required for a prolonged period.

Children and adolescents:

Filengia is not intended for children and adolescents, as it has not been studied in MS patients aged under 18.

Other medicines and Filengia:

Always tell your healthcare provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

Tell your doctor if you are taking any of the following medicines:

- Medicines that suppress or modulate the immune system, including other medicines used to treat MS, such as beta interferon, glatiramer acetate, natalizumab, mitoxantrone, teriflunomide, dimethyl fumarate or alemtuzumab. You must not use Filengia together with such medicines as this could intensify the effect on the immune system (see also ‘Do not take Filengia’).
- Corticosteroids, due to a possible added effect on the immune system.
- Vaccines. If you need to receive a vaccine, seek your doctor’s advice first. During and for up to 2 months after treatment with Filengia, you should not receive certain types of vaccine (live attenuated vaccines) as they could trigger the infection that they were supposed to prevent. Other vaccines may not work as well as usual if given during this period.
- Medicines that slow the heartbeat for example beta blockers (such as atenolol), calcium channel blockers (such as verapamil, diltiazem), ivabradine, digoxin, anticholinesteratic medicines or pilocarpine. Use of Filengia together with such medicines could intensify the effect on heartbeat in the first days after starting Filengia.
- Medicines for irregular heartbeat, such as quinidine, procainamide, disopyramide, amiodarone or sotalol. You must not use Filengia if you are taking such a medicine because it could intensify the effect on irregular heartbeat (see also ‘Do not take Filengia’).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

- Other medicines:
 - protease inhibitors, anti-infectives such as ketoconazole, azole antifungals, clarithromycin or telithromycin.
 - carbamazepine, rifampicin, phenobarbital, phenytoin, efavirenz or St. John's Wort.

Filengia with food and drink:

Filengia can be taken with or without food.

Pregnancy, breastfeeding and fertility:

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other healthcare provider for advice before taking Filengia.

Before you start treatment with Filengia your doctor may ask you to do a pregnancy test in order to ensure that you are not pregnant. You should avoid becoming pregnant while taking Filengia or in the two months after you stop taking it because there is a risk of harm to the baby. Talk to your doctor about reliable methods of birth control that you should use during treatment and for 2 months after you stop treatment.

Pregnancy:

You should not take Filengia if you are pregnant.

If you become pregnant while taking Filengia tell your doctor right away. You and your doctor will decide what is best for you and the baby.

Breastfeeding:

You should not take Filengia if you are breastfeeding your baby. Filengia can pass into breast milk and there is a risk of serious side effects for a breastfed baby. Talk with your doctor before breastfeeding while you take Filengia.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Fertility:

Data from preclinical studies do not suggest an effect on male and female fertility with Filengia.

Driving and using machines:

At initiation of treatment, you will have to stay at the doctor's surgery or clinic for 6 hours after taking the first dose of Filengia. Your ability to drive and use machines may be impaired during and potentially after this time period.

It is not always possible to predict to what extent Filengia may interfere with your daily activities. You should ensure that you do not engage in driving a vehicle or use machines until you are aware of the measure to which Filengia affects you.

3. HOW TO TAKE FILENGIA:

Do not share medicines prescribed for you with any other person.

Always take Filengia exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose of Filengia is one 0,5 mg capsule once a day by mouth.

Do not exceed the recommended dose.

Take one Filengia capsule (0,5 mg of fingolimod) once a day with a glass of water. Filengia capsules should always be swallowed intact, without opening them. Filengia can be taken with or without food.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Taking Filengia at the same time each day will help you remember when to take your medicine.

Your doctor will tell you how long your treatment with Filengia will last. If you have the impression that the effect of Filengia is too strong or too weak, tell your doctor or pharmacist.

If you take more Filengia than you should:

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison centre.

If you forget to take Filengia:

Do not take a double dose to make up for forgotten individual doses.

If you stop taking Filengia:

Do not stop taking Filengia or change your dose without talking to your doctor first.

Filengia will stay in your body for up to 2 months after you stop taking it. Your white blood cell count (lymphocyte count) may also remain low during this time and the side effects described in this leaflet may still occur. After stopping Filengia you may have to wait for 6-8 weeks before starting a new MS treatment.

If you have stopped Filengia for 1 day or more during the first 2 weeks of Filengia treatment, if you stop for more than 7 days during weeks 3 and 4 of treatment, or if you stop taking Filengia for more than 2 weeks after your first month of Filengia treatment, the initial effect on your heart rate may occur again. When you restart Filengia treatment, your doctor may decide to monitor your heart rate and blood pressure every hour, to run ECG's or to keep you under monitoring overnight. Do not restart Filengia after stopping it for more than two weeks without seeking advice from your doctor.

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Your doctor will decide whether and how you need to be monitored after stopping Filengia. Tell your doctor straight away if at any time you think your MS is getting worse after you have stopped treatment with Filengia. This could be serious.

4. POSSIBLE SIDE EFFECTS:

Filengia can have side effects.

Not all side effects reported for Filengia are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking Filengia, please consult your healthcare provider for advice.

If any of the following happens, stop taking Filengia and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Swelling of the hands, feet, ankles, face, lips and mouth or throat, which may cause difficulty in swallowing or breathing,
- rash or itching,
- fainting (dizziness).

These are all very serious side effects. If you have them, you may have had a serious reaction to Filengia. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following:

Frequent

- Bronchitis with symptoms such as cough with phlegm, chest pain, fever
- Gastroenteritis with symptoms such as vomiting, nausea, diarrhoea, fever
- Herpes virus infection (shingles or herpes zoster) with symptoms such as blisters, burning, itching or pain of

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

the skin, typically on the upper body or the face. Other symptoms may be fever and weakness in the early stages of infection, followed by numbness, itching or red patches with severe pain

- a type of skin cancer called basal cell carcinoma (BCC) which often appears as a pearly nodule, although it can also take other forms
- depression and anxiety are known to occur with increased frequency in the MS population and have also been reported in paediatric patients treated with Filengia
- slow heartbeat (bradycardia), irregular heart rhythm

Less frequent

- pneumonia with symptoms such as fever, cough, difficulty breathing
- malignant melanoma (a type of skin cancer which usually develops from an unusual mole). Possible signs of melanoma include moles which may change size, shape, elevation or colour over time, or new moles. The moles may itch, bleed or ulcerate
- lymphoma (a type of cancer that affects the lymph system)
- squamous cell carcinoma: a type of skin cancer which may present as a firm red nodule, a sore with crust, or a new sore on an existing scar
- tumour related to infection with human herpes virus (Kaposi's sarcoma)
- reduction in blood platelets which increases risk of bleeding or bruising
- convulsion, fits (more frequent in children and adolescents than in adults)
- a condition called posterior reversible encephalopathy syndrome (PRES). Symptoms may include sudden onset of severe headache, confusion, seizures and/or vision disturbances
- macular oedema (swelling in the central vision area of the retina at the back of the eye) with symptoms such as shadows or blind spot in the centre of the vision, blurred vision, problems seeing colours or details
- electrocardiogram anomaly (T-wave inversion)

Unknown frequency

- risk of a brain infection called progressive multifocal leukoencephalopathy (PML). The symptoms of PML

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA_{IN}, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

may be similar to an MS relapse. Symptoms might also arise that you might not become aware of by yourself, such as changes in mood or behaviour, memory lapses, speech and communication difficulties, which your doctor may need to investigate further to rule out PML. Therefore, if you believe your MS is getting worse or if you or those close to you notice any new or unusual symptoms, it is very important that you speak to your doctor as soon as possible

- cryptococcal infections (a type of fungal infection), including cryptococcal meningitis with symptoms such as headache accompanied by stiff neck, sensitivity to light, nausea, and/or confusion
- Merkel cell carcinoma (a type of skin cancer). Possible signs of Merkel cell carcinoma include flesh-coloured or bluish-red, painless nodule, often on the face, head or neck. Merkel cell carcinoma can also present as a firm painless nodule or mass. Long-term exposure to the sun and a weak immune system can affect the risk of developing Merkel cell carcinoma
- after Filengia treatment is stopped, symptoms of MS can return and may become worse compared to before or during treatment
- ischaemic and haemorrhagic strokes. Possible symptoms may include trouble walking, speaking and understanding, as well as paralysis or numbness of the face, arm or leg
- acute disseminated encephalomyelitis (ADEM)-like events where rapid appearance of encephalitis-like symptoms such as fever, fatigue, headache, nausea and vomiting can occur.
- an autoimmune form of anaemia (decreased amount of red blood cells) where red blood cells are destroyed (autoimmune haemolytic anaemia)
- haemophagocytic syndrome (HPS), a life-threatening syndrome where the body's immune system does not work normally. Symptoms include enlargement of the spleen, liver and lymph nodes
- peripheral arterial occlusive disease, a condition that may narrow arteries and reduce blood flow to the legs or arms
- Inflammatory disorder after stopping Filengia treatment (known as immune reconstitution inflammatory syndrome or IRIS)

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Frequent

- infection from flu virus with symptoms such as tiredness, chills, sore throat, aching in the joints or muscles, fever
- feeling of pressure or pain in the cheeks and forehead (sinusitis)
- ringworm, a fungal infection of the skin (tinea versicolor)
- low level of white blood cells (lymphocytes, leucocytes)
- headache, migraine
- blurred vision
- high blood pressure
- cough, breathlessness
- diarrhoea
- itchy, red, burning rash (eczema)
- hair loss
- back pain
- muscle pain
- joint pain
- weakness
- increased blood fat (triglycerides) level
- increased liver enzymes showing in blood tests
- weight loss

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Less frequent

- depressed mood
- nausea
- low level of certain white blood cells (neutrophils)
- swelling of your lower legs or hands (peripheral oedema)

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects:

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website.

By reporting side effects, you can help provide more information on the safety of Filengia.

5. HOW TO STORE FILENGIA:

STORE ALL MEDICINES OUT OF REACH OF CHILDREN.

Store at or below 25 °C.

Keep the capsule in the blister in the outer carton until required for use.

Do not use after the expiry date stated on the label / carton / bottle.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g., toilets).

6. CONTENTS OF THE PACK AND OTHER INFORMATION:

What Filengia contains:

The active substance is fingolimod (as hydrochloride).

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IAIN, C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/III/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

The other ingredients are pregelatinised starch, sodium laurilsulphate, gelatin, titanium dioxide (E171), yellow iron oxide (E172) and printing ink consisting of black iron oxide (E172), potassium hydroxide, propylene glycol, shellac and strong ammonia solution.

What Filengia looks like and contents of the pack:

Hard gelatin capsule (size 4), filled with white to off-white powder with small agglomerates. Capsule body: white opaque with TV 7820 imprinted. Capsule cap: yellow with TV 7820 imprinted.

The capsules are packed into:

Aluminium – aluminium blister packs

Forming foil: Aluminium soft temper (OPA/Alum/PVC)

Lidding foil: Aluminium foil, heat sealable

Pack sizes: 7, 10, 28, 30, 98 in blisters.

or

Aluminium – aluminium blister packs (peel)

Forming foil: Aluminium soft temper (OPA/Alum/PVC)

Lidding foil: Aluminium foil, peelable (Paper/PET/Alum)

Pack sizes: 7x1, 10x1, 28x1, 30x1, 98x1 and 100x1 in perforated unit-dose blisters.

Not all pack sizes may be marketed.

Holder of Certificate of Registration:

Teva Pharmaceuticals (Pty) Ltd

Maxwell Office Park

Magwa Crescent West

Waterfall City

Applicant: Teva Pharmaceuticals (Pty) Ltd	Product name: FILENGIA Dosage form & strength: Each capsule contains 0,5 mg fingolimod (as hydrochloride)
Date of Registration: 14 June 2022	FILENGIA: 52/34/0364
IA _{IN} , C.I.3.a – Implementation of the outcome of a procedure concerning PRAC EMA/PRAC/583624/2024 & EMEA/H/C/002202/II/0090/G & PSUSA/00001393/202302.	Date submitted to SAHPRA: 28 November 2025 eCTD seq 0007

Midrand

Gauteng

2090

Tel No.: + 27 (0) 11 635 0429

This leaflet was last revised in:

To be allocated.

Registration/Application number:

52/34/0364