

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**
Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**
Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

PROFESSIONAL INFORMATION for CETORIN ORAL SOLUTION 2,5 mg/ 5 mL

SCHEDULING STATUS

S1

1 NAME OF THE MEDICINE

CETORIN Oral Solution 2,5 mg/ 5 mL

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 5 mL contains 2,5 mg of Levocetirizine dihydrochloride

Contains sugar

Each 5 mL contains 10 mg of saccharin sodium and 2000 mg maltitol solution

'for full list of excipients, see section 6.1'

3 PHARMACEUTICAL FORM

Oral solution

Clear, colorless liquid, with a grape flavour

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

CETORIN Oral Solution is indicated for the relief of symptoms associated with the following allergic conditions:

- seasonal allergic rhinitis
- perennial allergic rhinitis
- chronic idiopathic urticaria.

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**
 Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**
 Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

4.2 Posology and method of administration

Posology

Adults and adolescents 12 years of age and older:

The daily recommended dose is 5 mg (10 mL) once daily.

Special populations

Elderly

Adjustment of the dose is recommended in elderly patients with moderate to severe renal impairment (see patients with renal impairment below).

Adults with renal impairment:

The dosing interval must be individualised according to renal function. Refer to the following table and adjust the dose as indicated. To use this dosing table, an estimate of the patient's creatinine clearance (Cl_{cr}) in mL/min is needed. The Cl_{cr} (mL/min) may be estimated from serum creatinine ($\mu\text{mol/L}$) determination using the following formula:

$$Cl_{cr} = \frac{(140 - \text{age}) \times \text{body weight in kg}}{\text{serum creatinine } (\mu\text{mol/L})} \quad (0,85 \text{ for women})$$

Dosing adjustments for patients with impaired renal function:

Group	Creatinine clearance (mL/min)	Dosage and frequency
Normal	≥ 80	5 mg once daily
Mild	50-79	5 mg once daily
Moderate	30-49	5 mg once every 2 days
Severe	<30	5 mg once every 3 days
End-stage renal disease – Patients undergoing dialysis	<10	Contraindicated

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

In paediatric patients suffering from renal impairment

The dose will have to be adjusted on an individual basis taking into account the renal clearance of the patient and his/her body weight. There are no specific data for children with renal impairment.

Patients with hepatic impairment

No dose adjustment is needed in patients with solely hepatic impairment. In patients with hepatic impairment and renal impairment, adjustment of the dose is recommended (see Patients with renal impairment above).

Paediatric population

Children aged less than 2 years

The administration of **CETORIN Oral Solution** to infants and toddlers aged less than 2 years is not recommended (see section 4.4).

Children aged 2 to 6 years

The daily recommended dose is 2,5 mg to be administered in 2 intakes of 1,25 mg (2,5 mL of solution twice daily).

Children aged 6 to 12 years

The daily recommended dose is 5 mg (10 mL) once daily.

Duration of use

Intermittent allergic rhinitis (symptoms < 4 days/week or during less than 4 weeks) has to be treated according to the disease and its history; it can be stopped once the symptoms have disappeared and can be restarted again when symptoms reappear. In case of persistent allergic rhinitis (symptoms > 4 days/week or during more than 4 weeks), continuous therapy can be proposed to the patient during the period of exposure to allergens. Clinical experience with **CETORIN Oral Solution** is currently available for a 6-month

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

treatment period.

Method of administration

For oral administration

4.3 Contraindications

CETORIN Oral Solution is contraindicated in

- hypersensitivity to the levocetirizine, to any piperazine derivative or to any of the excipients of **CETORIN Oral Solution** (see section 4.4)
- infants and toddlers aged less than 2 years, as safety and efficacy have not been demonstrated
- during breastfeeding of infants and while pregnant (see section 4.6) and lactation
- patients with end-stage renal disease, at less than 10 mL/min creatinine clearance.

4.4 Special warnings and precautions for use

Alcohol

Precaution is recommended with intake of alcohol. CETORIN Oral Solution lacks significant sedative effects. Patients should, however be warned that a small number of individuals may experience sedation. This effect may be compounded by the simultaneous intake of alcohol or other central nervous system depressants (see section 4.5).

Risk of urinary retention

Caution should be exercised in patients with predisposing factors of urinary retention (e.g. spinal cord lesion, prostatic hyperplasia) as levocetirizine, contained in CETORIN Oral Solution, may increase the risk of urinary retention.

Risk of convulsion

Caution should be taken in patients with epilepsy and patients at risk of convulsion as CETORIN Oral Solution may cause seizure aggravation.

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

Skin reactions

Pruritus may occur when levocetirizine, as in CETORIN Oral Solution, is stopped even if those symptoms were not present before treatment initiation.

The symptoms may resolve spontaneously. In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

Allergic skin tests

Response to allergy skin tests are inhibited by antihistamines and a wash-out period (of 3 days) is required before performing them.

Information on excipients

CETORIN Oral Solution contains maltitol and may have a laxative effect. Patients with rare hereditary problems of fructose intolerance should not take CETORIN Oral Solution.

Paediatric population

Infants and children under 2 years

The administration of CETORIN Oral Solution to infants and toddlers aged less than 2 years is not recommended due to the lack of sufficient data in this age group (see sections 4.2 and 4.3).

4.5 Interaction with other medicines and other forms of interaction

No interaction studies have been performed with levocetirizine (including no studies with CYP3A4 inducers). Studies with the racemate compound cetirizine demonstrated that there were no clinically relevant adverse interactions (with antipyrine, ketoconazole, erythromycin, azithromycin, cimetidine, pseudoephedrine, glipizide and diazepam).

Theophylline

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

A decrease in the clearance of cetirizine (16 %) was observed in a multiple dose study with theophylline (400 mg once a day); while the disposition of theophylline was not altered by concomitant cetirizine administration.

Ritonavir:

In a multiple dose study of ritonavir (600 mg twice daily) and cetirizine (10 mg daily), the extent of exposure to cetirizine was increased by about 40 % while the disposition of ritonavir was decreased (-11 %).

Food:

The extent of absorption of CETORIN Oral Solution is not reduced with food, although the rate of absorption is decreased.

Alcohol:

In sensitive patients the simultaneous administration of CETORIN Oral Solution and alcohol or other central nervous system (CNS) depressants may have effects on the central nervous system. It is advisable to avoid excessive alcohol consumption (see section 4.4).

4.6 Fertility, pregnancy and lactation

Pregnancy

CETORIN Oral Solution is contraindicated in pregnancy as safety has not been demonstrated (see section 4.3 and 5.3).

Breastfeeding

CETORIN Oral Solution is contraindicated in women who are breastfeeding their babies, since the active ingredient is excreted in breast milk.

Fertility

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**

Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

No clinical data relating to levocetirizine effects on fertility is available.

4.7 Effects on ability to drive and use machines

CETORIN Oral Solution lacks significant sedative effects. Nevertheless, some patients could experience somnolence, fatigue, and asthenia under therapy with CETORIN Oral Solution. Therefore, patients intending to drive, engage in potentially hazardous activities or operate machinery should take their response to the CETORIN Oral Solution into account.

4.8 Undesirable effects

a) Summary of the safety profile

Mild to moderate side effects most frequently experienced in adults include, headache, somnolence, dry mouth, fatigue, with asthenia or abdominal pain occurring less frequently.

Tabulated list of adverse reactions

System Organ Class	Frequency	Side effects
Immune system disorders	<i>Frequency unknown</i>	Angioedema, hypersensitivity including anaphylaxis
Metabolism and nutrition disorders	<i>Frequency unknown</i>	Increased weight, increased appetite
Psychiatric disorders	<i>Frequency unknown</i>	Aggression, agitation, hallucination, depression, insomnia, suicidal ideation, nightmare
Nervous system disorders	<i>Frequent</i> <i>Frequency unknown</i>	Somnolence, headache Convulsion, paraesthesia, dizziness, syncope, tremor, dysgeusia
Eye disorders	<i>Frequency unknown</i>	Visual disturbances, blurred vision
Ear and labyrinth disorders	<i>Frequency unknown</i>	Vertigo

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**

Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

Cardiac disorders	<i>Frequency unknown</i>	Palpitation, tachycardia
Respiratory, thoracic and mediastinal disorders	<i>Frequency unknown</i>	Dyspnoea
Gastrointestinal disorders	<i>Frequent</i> <i>Less frequent</i> <i>Frequency unknown</i>	Diarrhoea, constipation, dry mouth Nausea and gastro-intestinal discomfort, abdominal pain Vomiting
Hepato-biliary disorders	<i>Frequency unknown</i>	Hepatitis, abnormal liver function test*
Skin and subcutaneous tissue disorders	<i>Frequency unknown</i>	Rash, urticaria, pruritus, fixed drug eruption
Musculoskeletal, connective tissue disorders and bone disorders	<i>Frequency unknown</i>	Myalgia, arthralgia
Renal and urinary disorders	<i>Frequency unknown</i>	Dysuria, urinary retention
General disorders and administration site conditions	<i>Frequent</i> <i>Less frequent</i> <i>Frequency unknown</i>	Fatigue Asthenia, malaise Oedema

c) Description of selected adverse reactions

After levocetirizine discontinuation, pruritus has been reported.

d) Paediatric population

System Organ Class	Frequency	Side effects
Psychiatric disorders	<i>Frequent</i>	Sleep disorders

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**

Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

Nervous system disorders	<i>Frequent</i>	Somnolence
	<i>Less frequent</i>	Headache
Gastrointestinal disorders	<i>Frequent</i>	Diarrhoea, constipation
	<i>Less frequent</i>	Vomiting

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicine is important. It allows continued monitoring of the benefit/risk balance of the medicine. Health care providers **are asked to report any suspected adverse to report any suspected adverse reactions to SAHPRA via the “6.04 Adverse Drug Reactions Reporting Form”, found online under SAHPRA’s publications: <https://www.sahpra.org.za/publications/Index/8> or to the Holder of certificate of registration through the mail: pvg.cdma@heterogroups.com**

4.9 Overdose

Signs and symptoms

Symptoms of overdose may include drowsiness in adults and initially agitation and restlessness, followed by drowsiness in children.

Management of overdoses

There is no known specific antidote to **CETORIN** Oral Solution. Should overdose occur, symptomatic or supportive treatment is recommended. Levocetirizine is not effectively removed by haemodialysis.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antihistamines for systemic use, piperazine derivatives,

ATC code: R06A E09.

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**

Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

Pharmacological classification: A 5.7.1 Antihistaminics

Mechanism of action

Levocetirizine is the R-enantiomer of cetirizine, is a histamine H₁-receptors. Binding studies revealed that levocetirizine has high affinity for human H₁-receptors (K_i = 3,2 nmol/l). Levocetirizine has an affinity 2-fold higher than that of cetirizine (K_i = 6.3 nmol/l). Levocetirizine dissociates from H₁-receptors with a half-life of 115 ± 38 min. After single administration, levocetirizine shows a receptor occupancy of 90 % at 4 hours and 57 % at 24 hours.

Pharmacodynamic studies in healthy volunteers demonstrate that, at half the dose, levocetirizine has comparable activity to cetirizine, both in the skin and in the nose.

5.2 Pharmacokinetic properties

The pharmacokinetic profile is the same when given as the single enantiomer or when given as cetirizine. No chiral inversion occurs during the process of absorption and elimination.

Absorption

Levocetirizine is rapidly and extensively absorbed following oral administration. In adults, peak plasma concentrations are achieved 0,9 hours after dosing. Steady state is achieved after two days. Peak concentrations are typically 270 ng/ml and 308 ng/ml following a single and a repeated 5 mg o.d. dose, respectively. The extent of absorption is dose-independent and is not altered by food, but the peak concentration is reduced and delayed.

Distribution

No tissue distribution data are available in humans, neither concerning the passage of levocetirizine through the blood-brain-barrier. In rats and dogs, the highest tissue levels are found in liver and kidneys, the lowest in the CNS compartment.

Applicant/PHRC: Hetero Drugs South Africa (Pty) Ltd

Product proprietary name: Ceterin Oral Solution 2,5 mg/ 5 mL

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

In humans, levocetirizine is 90 % bound to plasma proteins. The distribution of levocetirizine is restrictive, as the volume of distribution is 0,4 l/kg.

Biotransformation

The extent of metabolism of levocetirizine in humans is less than 14% of the dose and therefore differences resulting from genetic polymorphism or concomitant intake of enzyme inhibitors are expected to be negligible.

Metabolic pathways include aromatic oxidation, N- and O- dealkylation and taurine conjugation. Dealkylation pathways are primarily mediated by CYP 3A4 while aromatic oxidation involved multiple and/or unidentified CYP isoforms. Levocetirizine had no effect on the activities of CYP isoenzymes 1A2, 2C9, 2C19, 2D6, 2E1 and 3A4 at concentrations well above peak concentrations achieved following a 5 mg oral dose. Due to its low metabolism and absence of metabolic inhibition potential, the interaction of levocetirizine with other substances, or vice-versa, is unlikely.

Elimination

The plasma half-life in adults is approximately 8 hours in adults. The half-life is shorter in small children. The mean apparent total body clearance in adults is 0.63 ml/min/kg. The main route of excretion of levocetirizine and metabolites is via urine, accounting for a mean of 85 % of the dose. Approximately 13 % is excreted in the faeces. Levocetirizine is excreted both by glomerular filtration and active tubular secretion.

Linearity

The pharmacokinetics of levocetirizine are linear with dose- and time-independent with low inter subject variability

Pharmacokinetics in special patient groups

Renal impairment

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

The apparent body clearance of levocetirizine is correlated to the creatinine clearance. It is therefore recommended to adjust the dosing intervals of levocetirizine, based on creatinine clearance in patients with moderate and severe renal impairment. In anuric end stage renal disease patients, the total body clearance is decreased by approximately 80 % when compared to normal patients. The amount of levocetirizine removed during a standard 4-hour hemodialysis procedure was < 10 %.

Elderly

Limited pharmacokinetic data are available in elderly patients. Total body clearance in the elderly is approximately 33 % lower compared to that in younger adults. The disposition of racemic cetirizine has been shown to be dependent on renal function rather than age. This would also be applicable for levocetirizine, as both cetirizine and levocetirizine are both predominantly excreted in urine. Therefore, the dose of levocetirizine should be adjusted in accordance with renal function in elderly patients.

Hepatic impairment

The pharmacokinetics of levocetirizine in hepatically impaired patients have not been tested. Patients with chronic liver diseases (hepatocellular, cholestatic, and biliary cirrhosis) given 10 or 20 mg of the racemic compound cetirizine as a single dose had a 50 % increase in half-life along with a 40 % decrease in clearance compared to healthy patients.

Gender

The same daily dose and dosing intervals are applicable for men and women with normal renal function.

Race

The effect of race on levocetirizine has not been studied. As levocetirizine is primarily renally excreted, and there are no important racial differences in creatinine clearance, pharmacokinetic characteristics of levocetirizine are not expected to be different across races. No race-related differences in the kinetics of racemic cetirizine have been observed.

Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd**

Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL**

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

Paediatric population

In children age 6 to 11 years with body weight ranging between 20 and 40 kg the C_{max} and AUC values are about 2-fold greater than in healthy adults, following oral administration of a single dose of 5 mg levocetirizine. Total body clearance is 30 % greater and the elimination half-life 24 % shorter in the paediatric population than in adults. Administration of 1,25 mg once daily to children 6 months to 5 years of age is expected to result in plasma concentrations similar to those of adults receiving 5 mg once daily.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Methyl paraben
- Propyl paraben
- Glycerin
- Sodium acetate trihydrate
- Glacial acetic acid
- Saccharin sodium
- Maltitol solution
- ART Grape FL#10273 Composition
 - Flavouring ingredients approved for use in a regulation of FDA
 - Propylene glycol

6.2 Incompatibilities

Not applicable

6.3 Shelf life

24 months

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

6.4 Special precautions for storage

Store at or below 25 °C. Protect from light and moisture.

KEEP OUT OF REACH OF CHILDREN.

6.5 Nature and contents of container

White opaque round shaped polypropylene 185 ml container pack filled with 148 mL, child resistant plastic cap with FSE wad, 28 mm

or

Amber color polypropylene 115 mL container pack filled with 100 mL, with child resistant plastic cap with FSE wad, 28 mm

6.6 Special precautions for disposal and other handling

No special requirements

7 HOLDER OF CERTIFICATE OF REGISTRATION

Hetero Drugs South Africa (Pty) Ltd

Waterfall Corporate Campus,

Building No.2, First Floor,

74 Waterfall Drive,

Midrand

2066

Telephone number: 012 644 1220

Fax number: 012 644 1564

e-mail address: nokuthula.n@hetero.com

8 REGISTRATION NUMBER

*Applicant/PHRC: **Hetero Drugs South Africa (Pty) Ltd***

*Product proprietary name: **Cetorin Oral Solution 2,5 mg/ 5 mL***

Dosage form and strength: Oral solution, Each 5 mL contains 2,5 mg Levocetirizine dihydrochloride

52/5.7.1/0935

9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

06 September 2022

10 DATE OF REVISION OF THE TEXT

11 June 2026