

PATIENT INFORMATION LEAFLET FOR
CORYX PAEDIATRIC

Scheduling Status

S2

CORYX PAEDIATRIC

**(Each 5 mL contains 1,25 mg Triprolidine HCl,
25,00 mg pseudoephedrine HCl and 75,0 mg vitamin C. Syrup.)**

Contains sugar: Sorbitol 70% 3150, 00 mg/5 mL

Contains sweeteners: Sodium saccharin 1,00 mg/5 mL; Sodium cyclamate 10,00 mg/5 mL

Read all of this leaflet carefully because it contains important information for you

CORYX PAEDIATRIC is available without a doctor's prescription, for you to treat a mild illness.

Nevertheless, you still need to use CORYX PAEDIATRIC carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share CORYX PAEDIATRIC with any other person.
- Ask your pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or do not improve after 5 days.

What is in this leaflet

1. What CORYX PAEDIATRIC is and what it is used for
2. What you need to know before you take CORYX PAEDIATRIC
3. How to take CORYX PAEDIATRIC
4. Possible side effects

5. How to store CORYX PAEDIATRIC
6. Contents of the pack and other information.

1. What CORYX PAEDIATRIC is and what it is used for

CORYX PAEDIATRIC belongs to a class of medicine known as preparations for the common cold including nasal decongestants and antihistaminics.

CORYX PAEDIATRIC syrup is indicated for the relief of symptoms associated with colds and influenza such as nasal congestion.

2. What you need to know before you take CORYX PAEDIATRIC

Do not take CORYX PAEDIATRIC if you:

- if you are hypersensitive (allergic) to triprolidine HCl, pseudoephedrine HCl, aspirin, vitamin C, or any of the other ingredients of CORYX PAEDIATRIC.
- if you are pregnant or breastfeeding your baby,
- if you are taking monoamine oxidase inhibitors or within 14 days of stopping such treatment,
- if you are a child under the age of 2 years.
- Diabetes (High blood sugar levels)
- Hyperthyroidism (overactive thyroid)
- If you have Pheochromocytoma (rare tumour of the adrenal gland)
- If you have heart disease including high blood pressure
- If you have high pressure within the eye
- If you have severe impairment of the kidneys
- If you are unusually or excessively excitable
- If you have a disease which affects the lungs

- If you are receiving other medicines used for nasal or eye congestion, medicines used to decrease hunger or other stimulant medicines.
- If you have severe high blood pressure or uncontrolled high blood pressure.
- If you have severe disease of the kidney

Take special care with CORYX PAEDIATRIC:

- Cases of posterior reversible encephalopathy syndrome (PRES) and reversible cerebral vasoconstriction syndrome (RCVS) have been reported following use of medicines containing pseudoephedrine. PRES and RCVS are rare conditions that can involve reduced blood supply to the brain. Stop using CORYX PAEDIATRIC SYRUP immediately and seek immediate medical assistance if you develop symptoms that may be signs of PRES or RCVS (see **section 4**).
- if you have heart disease or cardiovascular disease,
- if you have hardening or thinning of the arteries,
- if you have high blood pressure,
- if you have angina pectoris,
- if you have hepatic impairment.
- If you suffer from difficulty of urination or enlargement of the prostate gland
- If you suffer from severe skin reactions
- If you suffer from inflammation of the large intestine.
- If you experience loss of vision or decreased visual acuity.

Please consult your doctor or pharmacist before taking CORYX PAEDIATRIC syrup for advice if you have any of the above.

- CORYX PAEDIATRIC syrup may make you sleepy.

Other medicines and CORYX PAEDIATRIC

Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines).

- Care should be taken if concomitantly taking CORYX PAEDIATRIC syrup with the following: alcohol, pain medicines, sedatives, tranquillisers and antidepressants.
- Care should be taken if concomitantly taking CORYX PAEDIATRIC syrup with certain antibiotics and medicines which can affect the hearing.
- CORYX PAEDIATRIC syrup may interfere with certain skin tests and has to be stopped a few days before the testing.
- Care should be taken if concomitantly taking CORYX PAEDIATRIC syrup with guanethidine, reserpine, methyldopa or atropine, anticoagulants.
- CORYX PAEDIATRIC syrup may interfere with skin testing and should be stopped several days before skin testing.
- CORYX PAEDIATRIC syrup should be used with caution with the concurrent use of desferrioxamine, hormonal contraceptives, hormone replacement therapy, fluphenazine and warfarin.
- The use of CORYX PAEDIATRIC syrup should be stopped before undergoing anaesthesia.

Taking CORYX PAEDIATRIC with food, drink and alcohol

It is preferable to take CORYX PAEDIATRIC syrup after meals.

Pregnancy, breastfeeding and fertility

Safety in pregnancy and lactation has not been established.

If you are pregnant or breastfeeding your baby while taking this medicine, please consult your doctor, pharmacist or other healthcare professional for advice before taking CORYX PAEDIATRIC syrup.

Driving and using machinery

- CORYX PAEDIATRIC syrup may make you feel drowsy/sleepy.
- Do not drive because CORYX PAEDIATRIC syrup could interfere with your ability to drive safely.
- Do not operate any tools or machines.
- Do not climb dangerous heights or perform potentially dangerous tasks where impaired decision making could lead to accidents.

Taking other medicines with CORYX PAEDIATRIC syrup

- Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines.)
- Care should be taken if concomitantly taking CORYX PAEDIATRIC syrup with the following: alcohol, pain medicines, sedatives, tranquillisers and antidepressants.
- Care should be taken if concomitantly taking CORYX PAEDIATRIC syrup with certain antibiotics and medicines which can affect the hearing.

- CORYX PAEDIATRIC syrup may interfere with certain skin tests and has to be stopped a few days before the testing.
- Care should be taken if concomitantly taking CORYX PAEDIATRIC syrup with guanethidine, reserpine, methyldopa or atropine, anticoagulants.
- CORYX PAEDIATRIC syrup may interfere with skin testing and should be stopped several days before skin testing.
- CORYX PAEDIATRIC syrup should be used with caution with the concurrent use of desferrioxamine, hormonal contraceptives, hormone replacement therapy, fluphenazine and warfarin.
- The use of CORYX PAEDIATRIC syrup should be stopped before undergoing anaesthesia.

CORYX PAEDIATRIC contains sugar (Sorbitol)

Sorbitol may cause gastrointestinal discomfort and mild laxative effect.

3. How to take CORYX PAEDIATRIC SYRUP

Do not share medicines prescribed for you with any other person.

Always take CORYX PAEDIATRIC syrup exactly as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure.

Adults and children over 12 years

2 medicine measures (10,0 mL) three times a day

Children 6 to 12 years

1 medicine measure (5,0 mL) three times a day

Children 2 to 5 years

½ medicine measure (2,5 mL) three times a day

Children 6 months to 2 years

Not recommended.

If you have the impression that the effect of CORYX PAEDIATRIC syrup is too strong or too weak, tell your doctor or pharmacist.

If you take CORYX PAEDIATRIC syrup than you should

Overdosage may be fatal especially in children. In the event of overdosage, consult your doctor or pharmacist. If neither is available, seek help at contact the nearest hospital or poison control centre.

If you forget to take a dose of CORYX PAEDIATRIC syrup

Do not take a double dose to make up for forgotten individual doses.

4. Possible Side Effects

CORYX PAEDIATRIC syrup can have side effects.

Less frequent side effects include: feelings of fear, anxiety, restlessness, insomnia, sedation, lassitude, incoordination, confusion, irritability, euphoria or nervousness, dizziness, fainting, blurred vision, uneven heart rhythm, heart pain, high or low blood pressure, dryness of mouth, throat and respiratory passages, tightness of the chest, diarrhoea, gastrointestinal disturbances, nausea, vomiting, loss of appetite, constipation, too much salivation, difficult urination or retention, headache, fatigue, and weakness.

Unknown side effects include: serious conditions affecting blood vessels in the brain known as posterior reversible encephalopathy syndrome (PRES) and reversible cerebral vasoconstriction syndrome (RCVS)

Not all side effects reported for CORYX PAEDIATRIC syrup are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your doctor, pharmacist or other healthcare professional for advice.

If any of the following happens, stop taking CORYX PAEDIATRIC syrup and tell your doctor immediately or go to the casualty department at your nearest hospital:

- severe headache with a sudden onset
- feeling sick
- vomiting
- confusion
- seizures
- changes in vision
- swelling of the hands, feet, ankles, face, lips, mouth or throat, which may cause difficulty in swallowing or breathing,

- rash or itching,
- fainting,
- yellowing of the skin and eyes, also called jaundice.

These are all very serious side effects. If you have them, you may have had a serious allergic reaction to CORYX PAEDIATRIC syrup. You may need urgent medical attention or hospitalisation.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on the SAHPRA website, or to Cipla Medpro (Pty) Ltd. by e-mail: drugsafetysa@cipla.com or telephone: 080 222 6662 (toll free). By reporting side effects, you can help provide more information on the safety of CORYX PAEDIATRIC.

5. How to store CORYX PAEDIATRIC

Store all medicines out of the reach of children.

- Store CORYX PAEDIATRIC in the tube tightly closed at or below 25 °C.
- Store in the original container.
- Keep the container in the outer carton.
- Protect from light.
- Do not use after the expiry date stated on the label.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

6. Contents of the pack and other information

What CORYX PAEDIATRIC contains

The active substances in each 5 mL of syrup are contains: 1,25 mg Triprolidine HCl, 25,00 mg pseudoephedrine HCl and 75,0 mg vitamin C.

The other ingredients are: black currant flavour, brilliant blue colour, carmelose sodium, citric acid anhydrous, glycerol, potassium sorbate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution 70 %, sodium cyclamate, sodium saccharin.

What CORYX PAEDIATRIC looks like and contents of the pack

CORYX PAEDIATRIC syrup is a clear blue to blue-green coloured syrup with a black current smell and taste.

CORYX PAEDIATRIC syrup is packed in cartons containing 70 mL, 100 mL, or 140 mL round amber glass bottles with white plastic caps.

Holder of Certificate of Registration

CIPLA MEDPRO (PTY) LTD.

Building 9

Parc du Cap

Mispel Street

Bellville

7530

Customer Care: 080 222 6662

This leaflet was revised in

31 March 2026

Registration number(s)

31/5.8/0056

Namibia: NS1
04/5.6/1652

Botswana: NS3 BOT0801263 100 mL

Botswana: NS3 BOT0801263A 140 mL

Access to the corresponding Professional Information

To access corresponding Professional Information, scan the QR Code below.

PLACE HOLDER:

The QR Code to
be generated and
included after
approval.