

PATIENT INFORMATION LEAFLET

SCHEDULING STATUS: S5

ETIFOXINE 50 BIOTECH 50 mg capsules, hard

Etifoxine hydrochloride

Contains sugar:

Each tablet contains 110 mg lactose monohydrate.

Read all of this leaflet carefully before you start taking ETIFOXINE 50 BIOTECH

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor, pharmacist, nurse or other health care provider.
- ETIFOXINE 50 BIOTECH has been prescribed for you personally and you should not share your medicine with other people. It may harm them, even if their symptoms are the same as yours.

What is in this leaflet

1. What ETIFOXINE 50 BIOTECH is and what it is used for
2. What you need to know before you take ETIFOXINE 50 BIOTECH
3. How to take ETIFOXINE 50 BIOTECH
4. Possible side effects
5. How to store ETIFOXINE 50 BIOTECH
6. Contents of the pack and other information

1. What ETIFOXINE 50 BIOTECH is and what it is used for

ETIFOXINE 50 BIOTECH is a tranquilliser used to oppose various emotional and bodily reactions, which accompany anxiety.

2. What you need to know before you take ETIFOXINE 50 BIOTECH

Do not take ETIFOXINE 50 BIOTECH

- If you are hypersensitive (allergic) to ETIFOXINE 50 BIOTECH or to any of the other ingredients of ETIFOXINE 50 BIOTECH listed in section 6 of this leaflet.
- If you are in a state of shock.
- If you have severely impaired liver and/or renal function.
- If you suffer from myasthenia gravis (a disease which causes weak muscles).
- If you have had severe liver problems, such as inflammation of the liver (hepatitis) or cytolytic hepatitis, during previous treatment with ETIFOXINE 50 BIOTECH.
- If you have had severe skin reactions during previous treatment with ETIFOXINE 50 BIOTECH.

Warnings and precautions

Talk to your doctor before taking ETIFOXINE 50 BIOTECH:

- If you are at risk of developing liver problems, your doctor will do some tests to check your liver function before starting ETIFOXINE 50 BIOTECH and around one month after you start treatment.

You should stop taking the medicine and seek urgent medical attention if you experience the following events during treatment with ETIFOXINE 50 BIOTECH:

- Severe skin or allergic reactions (see section 4).
- Jaundice (yellowing of the skin and eyes), vomiting, tiredness, abdominal (belly) pain or swelling, swelling in the legs and ankles, dark urine colour, pale stool colour, or bloody or tar-coloured stool. These could be signs of severe liver problems (see section 4).
- Watery diarrhoea (see section 4).

Talk to your doctor or pharmacist if you experience bleeding from the uterus between menstrual periods (metrorrhagia) when on oral contraceptives during treatment with ETIFOXINE 50

BIOTECH.

Other medicines and ETIFOXINE 50 BIOTECH

Always tell your health care provider if you are taking any other medicine. (This includes all complementary or traditional medicines.)

ETIFOXINE 50 BIOTECH should not be used together with other central nervous system depressants (medicines that slow down the brain activity) as the depressant effect may be increased. Examples of other central nervous system depressants are:

- Medicines related to morphine (painkillers, cough mixtures).
- Antidepressants.
- Hypnotic medicines (medicines used to help you sleep).
- Antihistamines (medicines used to relieve allergic symptoms).
- Medicines used in the treatment of high blood pressure.
- Baclofen (a muscle relaxant).
- Thalidomide (a medicine used to treat a certain type of cancer).

ETIFOXINE 50 BIOTECH with food, drink and alcohol

Do not use alcohol or alcohol-containing medicines together with ETIFOXINE 50 BIOTECH as alcohol increases the sedating effect of ETIFOXINE 50 BIOTECH. This may lead to decreased alertness and may make vehicle driving and machine operations dangerous.

Pregnancy and breastfeeding

If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist or other health care provider for advice before taking ETIFOXINE 50 BIOTECH.

ETIFOXINE 50 BIOTECH should not be used during pregnancy or breastfeeding.

Driving and using machines

Do not drive or operate any tools or machinery as ETIFOXINE 50 BIOTECH causes drowsiness, and impaired alertness (when used in combination with central nervous system depressants or alcohol/ alcohol containing medicines), and it may hamper your ability to carry out these tasks.

ETIFOXINE 50 BIOTECH contains azorubine

Azorubine is an azo colourant which may cause allergic reactions.

ETIFOXINE 50 BIOTECH contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking ETIFOXINE 50 BIOTECH.

3. How to take ETIFOXINE 50 BIOTECH

Do not share medicines prescribed for you with any other person.

Always take ETIFOXINE 50 BIOTECH exactly as your doctor or pharmacist has told you. You should check with your doctor or pharmacist if you are not sure.

The usual dose is 3 to 4 capsules (150 mg – 200 mg) daily in two to three divided doses. The duration of treatment is usually from a few days to not more than 8 weeks, depending on your doctor's recommendations.

The capsules should be swallowed with a little water.

Stopping the treatment does not lead to withdrawal.

Your doctor will tell you how long your treatment with ETIFOXINE 50 BIOTECH will last. If you have the impression that the effect of ETIFOXINE 50 BIOTECH is too strong or too weak, tell your doctor or pharmacist.

If you take more ETIFOXINE 50 BIOTECH than you should

In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact your nearest hospital or poison centre.

Taking too much ETIFOXINE 50 BIOTECH may increase the risk of somnolence (drowsiness).

If you forget to take ETIFOXINE 50 BIOTECH

If you forget to take a dose, take it as soon as you remember. However if it is time for the next dose, skip the missed dose. Do not take a double dose to make up for a forgotten dose.

4. Possible side effects

ETIFOXINE 50 BIOTECH can have side effects.

Not all side effects reported for ETIFOXINE 50 BIOTECH are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking ETIFOXINE 50 BIOTECH, please consult your health care provider for advice.

If any of the following happens, stop taking ETIFOXINE 50 BIOTECH and tell your doctor immediately or go to the casualty department at your nearest hospital:

- Anaphylactic shock (severe flushing, itching, nausea, vomiting, swelling of the mouth and tongue and throat).
- Severe allergic reactions of the skin and face (flushing, rashes, itching, swelling).
- Allergic manifestations such as urticaria (nettle rash, hives) and angioedema (weals, swelling of lips, eyes and tongue).
- Allergic reactions with fever, rash, conjunctivitis (red, swollen eyes).
- Jaundice (yellowing of the skin and eyes), vomiting, tiredness, abdominal (belly) pain - these could be signs of severe liver problem.
- Watery diarrhoea.

These are all very serious side effects. If you have them, you may have had a serious reaction to ETIFOXINE 50 BIOTECH. You may need urgent medical attention or hospitalisation.

Tell your doctor if you notice any of the following:

Less frequent side effects:

- Bleeding between the expected menstrual periods, in women taking oral contraceptive pill.
- Slight drowsiness at the start of treatment and disappearing spontaneously with its continuation.
- Feeling of rapidly progressing obstruction in the throat (Quincke's oedema).

The following side effects may occur, but the frequency is unknown:

- Leukocytoclastic vasculitis (is a skin disorder or condition that is caused by the inflammation of small blood vessels) symptoms include: muscle aches, fever, abdominal pain, cough, joint pain, vomiting, bloody stool, numbness and weakness.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects

If you get side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org) found on SAHPRA website. By reporting side effects, you can help provide more information on the safety of ETIFOXINE 50 BIOTECH.

5. How to store ETIFOXINE 50 BIOTECH

- Store at or below 25 °C.
- Keep the blister strips in the outer carton until required for use.
- Store in the original package and do not store in bathrooms.
- Do not use after the expiry date stated on the carton.
- Return all unused medicine to your pharmacist.
- Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

- STORE ALL MEDICINES OUT OF REACH OF CHILDREN.

6. Contents of the pack and other information

What ETIFOXINE 50 BIOTECH contains

The active substance is etifoxine hydrochloride.

Each capsule contains 50 mg etifoxine hydrochloride.

The other ingredients are fumaric acid, lactose monohydrate and stearic acid. The capsule body and cap contain gelatine, Patent blue, azorubine and titanium dioxide.

What ETIFOXINE 50 BIOTECH looks like and contents of the pack

Hard gelatine capsule, with a white-coloured opaque body and blue-coloured opaque cap containing an off-white powder.

Clear PVC/PVDC aluminium blister strips packed in a cardboard carton.

Pack size: 60 capsules.

Holder of certificate of registration

Biotech Laboratories (Pty) Ltd

Ground floor, Block K West, Central Park

400 16th Road, Randjespark

Midrand 1685

South Africa

Tel no: +27 (0)11 848 3050

This leaflet was last revised

Registration number

59/2.6/0168.167

Date of registration

14 July 2026