

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

PROFESSIONAL INFORMATION

SCHEDULING STATUS

S4

WARNING: RISK OF THYROID C-CELL TUMORS

- In rodents, semaglutide causes dose-dependent and treatment-duration-dependent thyroid C-cell tumours at clinically relevant exposures. It is unknown whether RYBELSUS causes thyroid C-cell tumours, including medullary thyroid carcinoma (MTC), in humans as human relevance of semaglutide-induced rodent thyroid C-cell tumours has not been determined.
- RYBELSUS is contraindicated in patients with a personal or family history of MTC or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk for MTC with the use of RYBELSUS and inform them of symptoms of thyroid tumours (e.g. a mass in the neck, dysphagia, dyspnoea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with RYBELSUS.

1 NAME OF THE MEDICINE

Rybelsus 3 mg tablets

Rybelsus 7 mg tablets

Rybelsus 14 mg tablets

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Rybelsus 3 mg tablets

Each tablet contains 3 mg semaglutide*.

Rybelsus 7 mg tablets

Each tablet contains 7 mg semaglutide*.

Rybelsus 14 mg tablets

Each tablet contains 14 mg semaglutide*.

*human glucagon-like peptide-1 (GLP-1) analogue produced in *Saccharomyces cerevisiae* cells by recombinant DNA technology.

Excipient with known effect

Each tablet, regardless of semaglutide strength, contains 23 mg sodium.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tablet

Rybelsus 3 mg tablets

White to light yellow, oval shaped tablet (7,5 mm x 13,5 mm) debossed with '3' on one side and 'novo' on the other side.

Rybelsus 7 mg tablets

White to light yellow, oval shaped tablet (7,5 mm x 13,5 mm) debossed with '7' on one side and 'novo' on the other side.

Rybelsus 14 mg tablets

White to light yellow, oval shaped tablet (7,5 mm x 13,5 mm) debossed with '14' on one side and 'novo' on the other side.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

Rybelsus is indicated for the treatment of adults with insufficiently controlled type 2 diabetes mellitus as an adjunct to diet and exercise.

- as monotherapy when metformin is considered inappropriate due to intolerance or contraindications in combination with other medicinal products for the treatment of diabetes.
- For study results with respect to combinations, effects on glycaemic control and cardiovascular events, and the populations studied, see sections 4.4, 4.5 and 5.1.

4.2 Posology and method of administration

The starting dose of Rybelsus is 3 mg once daily for one month. After one month, the dose should be increased to a maintenance dose of 7 mg once daily. After at least one month on

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

with a dose of 7 mg once daily, the dose can be increased to a maintenance dose of 14 mg once daily to further improve glycaemic control.

The maximum recommended single daily dose of Rybelsus is 14 mg. Taking two 7 mg tablets to achieve the effect of a 14 mg dose has not been studied and is therefore not recommended.

Rybelsus can be used as monotherapy or in combination with one or more glucose-lowering medicinal products (see section 5.1).

When Rybelsus is used in combination with metformin and/or a sodium-glucose co-transporter-2 inhibitor (SGLT2i) or thiazolidinedione, the current dose of metformin and/or SGLT2i/thiazolidinedione can be continued.

When Rybelsus is used in combination with a sulfonylurea or insulin, a reduction in the dose of sulfonylurea or insulin should be considered to reduce the risk of hypoglycaemia (see section 4.4).

Self-monitoring of blood glucose is not needed in order to adjust the dose of semaglutide. Blood glucose self-monitoring is necessary to adjust the dose of sulfonylurea and insulin, particularly when semaglutide is started and insulin is reduced. A stepwise approach to insulin reduction is recommended.

Missed dose

If a dose is missed, the missed dose should be skipped, and the next dose should be taken the following day.

Special populations

Elderly

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

No dose adjustment is required based on age. Therapeutic experience in patients ≥ 75 years of age is limited (see section 5.2).

Renal impairment

No dose adjustment is required for patients with mild, moderate or severe renal impairment.

Experience with the use of semaglutide in patients with severe renal impairment is limited.

Semaglutide is not recommended in patients with end-stage renal disease (see section 5.2).

Hepatic impairment

No dose adjustment is required for patients with hepatic impairment. Experience with the use of semaglutide in patients with severe hepatic impairment is limited. Caution should be exercised when treating these patients with semaglutide (see section 5.2).

Paediatric population

The safety and efficacy of Rybelsus in children and adolescents below 18 years have not been established. No data are available.

Method of administration

Rybelsus is a tablet for once-daily oral use.

- This medicinal product should be taken on an empty stomach at any time of the day.
- It should be swallowed whole with a sip of water (up to half a glass of water equivalent to 120 ml). Tablets should not be split, crushed or chewed, as it is not known whether this impacts absorption of semaglutide.
- Patients should wait at least 30 minutes before eating or drinking or taking other oral medicinal products. Waiting less than 30 minutes decreases the absorption of semaglutide (see sections 4.5 and 5.2).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

4.4 Special warnings and precautions for use

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

General

Semaglutide should not be used in patients with type 1 diabetes mellitus or for the treatment of diabetic ketoacidosis. Diabetic ketoacidosis has been reported in insulin-dependent patients whom had rapid discontinuation or dose reduction of insulin when treatment with a GLP-1 receptor agonist is started (see section 4.2).

There is no therapeutic experience in patients with congestive heart failure New York Heart Association (NYHA) class IV and semaglutide is therefore not recommended in these patients.

There is no therapeutic experience with semaglutide in patients with bariatric surgery.

Gastrointestinal effects

Use of GLP-1 receptor agonists may be associated with gastrointestinal adverse reactions that can cause dehydration, which in rare cases can lead to a deterioration of renal function (see section 4.8). Patients treated with semaglutide should be advised of the potential risk of dehydration in relation to gastrointestinal side effects and take precautions to avoid fluid depletion.

Acute pancreatitis

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Acute pancreatitis has been observed with the use of GLP-1 receptor agonists. Patients should be informed of the characteristic symptoms of acute pancreatitis. If pancreatitis is suspected, semaglutide should be discontinued; if confirmed, semaglutide should not be restarted. Caution should be exercised in patients with a history of pancreatitis.

Hypoglycaemia

Patients treated with semaglutide in combination with a sulfonylurea or insulin may have an increased risk of hypoglycaemia (see section 4.8). The risk of hypoglycaemia can be lowered by reducing the dose of sulfonylurea or insulin when initiating treatment with semaglutide (see section 4.2).

Diabetic retinopathy

Rapid improvement in glucose control has been associated with a temporary worsening of diabetic retinopathy, but other mechanism cannot be excluded. Long-term glycaemic control decreases the risk of diabetic retinopathy. Patients with a history of diabetic retinopathy should be monitored for worsening and treated according to clinical guidelines.

Treatment response

Compliance with the dosing regimen is recommended for optimal effect of semaglutide. If the treatment response with semaglutide is lower than expected, the treating physician should be aware that the absorption of semaglutide is highly variable and may be minimal (2 – 4 % of patients will not have any exposure), and that the absolute bioavailability of semaglutide is low.

Sodium content

This medicinal product contains 23 mg sodium per tablet, equivalent to 1 % of the WHO recommended maximum daily intake of 2 g sodium for an adult.

4.5 Interaction with other medicines and other forms of interaction

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Semaglutide delays gastric emptying which may influence the absorption of other oral medicinal products.

Effects of semaglutide on other medicinal products

Thyroxine

Total exposure (AUC) of thyroxine (adjusted for endogenous levels) was increased by 33 % following administration of a single dose of levothyroxine. Maximum exposure (C_{max}) was unchanged. Monitoring of thyroid parameters should be considered when treating patients with semaglutide at the same time as levothyroxine.

Warfarin

Semaglutide did not change the AUC or C_{max} of R- and S-warfarin following a single dose of warfarin, and the pharmacodynamic effects of warfarin as measured by the international normalised ratio (INR) were not affected in a clinically relevant manner. However, upon initiation of semaglutide treatment in patients on warfarin or other coumarin derivatives, frequent monitoring of INR is recommended.

Rosuvastatin

AUC of rosuvastatin was increased by 41 % [90 % CI: 24; 60] when co-administered with semaglutide. Based on the wide therapeutic index of rosuvastatin the magnitude of changes in the exposure is not considered clinically relevant.

Digoxin, oral contraceptives, metformin, furosemide

No clinically relevant change in AUC or C_{max} of digoxin, oral contraceptives (containing ethinylestradiol and levonorgestrel), metformin or furosemide was observed when concurrently administered with semaglutide.

Interactions with medicinal products with very low bioavailability (F: 1 %) have not been evaluated.

Effects of other medicinal products on semaglutide

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Omeprazole

No clinically relevant change in AUC or C_{max} of semaglutide was observed when taken with omeprazole.

In a trial investigating the pharmacokinetics of semaglutide co-administered with five other tablets, the AUC of semaglutide decreased by 34 % and C_{max} by 32 %. This suggests that the presence of multiple tablets in the stomach influences the absorption of semaglutide if co-administered at the same time. After administering semaglutide, the patients should wait 30 minutes before taking other oral medicinal products (see section 4.2).

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Women of childbearing potential are recommended to use contraception when treated with semaglutide.

Pregnancy

Studies in animals have shown reproductive toxicity (see section 5.3). There are limited data from the use of semaglutide in pregnant women. Therefore, semaglutide should not be used during pregnancy. If a patient wishes to become pregnant, or pregnancy occurs, semaglutide should be discontinued. Semaglutide should be discontinued at least 2 months before a planned pregnancy due to the long half-life (see section 5.2).

Breastfeeding

In lactating rats, semaglutide, salcaprozate sodium and/or its metabolites were excreted in milk. As a risk to a breast-fed child cannot be excluded, Rybelsus should not be used during breast-feeding.

Fertility

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

The effect of semaglutide on fertility in humans is unknown. Semaglutide did not affect male fertility in rats. In female rats, an increase in oestrous length and a small reduction in number of ovulations were observed at doses associated with maternal body weight loss (see section 5.3).

4.7 Effects on ability to drive and use machines

Semaglutide has no or negligible influence on the ability to drive or use machines. When it is used in combination with a sulfonylurea or insulin, patients should be advised to take precautions to avoid hypoglycaemia while driving and using machines (see section 4.4).

4.8 Undesirable effects

Summary of the safety profile

In 10 phase 3a trials, 5 707 patients were exposed to semaglutide alone or in combination with other glucose-lowering medicinal products. The duration of the treatment ranged from 26 weeks to 78 weeks. The most frequently reported adverse reactions in clinical trials were gastrointestinal disorders, including nausea (very common), diarrhoea (very common) and vomiting (common).

Tabulated list of adverse reactions

[Table 1](#) lists adverse reactions identified in all phase 3a trials in patients with type 2 diabetes mellitus (further described in section 5.1). The frequencies of the adverse reactions are based on a pool of the phase 3a trials excluding the cardiovascular outcomes trial.

The reactions are listed below by system organ class and absolute frequency. Frequencies are defined as: very common: ($\geq 1/10$); common: ($\geq 1/100$ to $< 1/10$); uncommon: ($\geq 1/1,000$ to $< 1/100$); rare: ($\geq 1/10,000$ to $< 1/1,000$) and very rare: ($< 1/10,000$). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Table 1 Adverse reactions from controlled phase 3a trials

MedDRA system organ class	Very common	Common	Uncommon	Rare
Immune system disorders				Anaphylactic reaction
Metabolism and nutrition disorders	Hypoglycaemia when used with insulin or sulfonylurea ^a	Hypoglycaemia when used with other oral antidiabetic products ^a Decreased appetite		
Eye disorders		Diabetic retinopathy complications ^b		
Cardiac disorders			Increased heart rate	
Gastrointestinal disorders	Nausea Diarrhoea	Vomiting Abdominal pain Abdominal distension Constipation Dyspepsia Gastritis Gastro-oesophageal reflux disease Flatulence	Eructation	Acute pancreatitis

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Hepatobiliary disorders			Cholelithiasis	
General disorders and administration site conditions		Fatigue		
Investigations		Increased lipase Increased amylase	Weight decreased	

a) Hypoglycaemia defined as blood glucose < 3,0 mmol/L or < 54 mg/dL

b) Diabetic retinopathy complications is a composite of retinal photocoagulation, treatment with intravitreal agents, vitreous haemorrhage and diabetes-related blindness (uncommon). Frequency is based on the cardiovascular outcomes trial with s.c. semaglutide, but it cannot be excluded that the risk of diabetic retinopathy complications identified also applies to Rybelsus.

Description of selected adverse reactions

Hypoglycaemia

Severe hypoglycaemia was primarily observed when semaglutide was used with a sulfonylurea (< 0,1 % of subjects, < 0,001 events/patient year) or insulin (1,1 % of subjects, 0,013 events/patient year). Few episodes (0,1 % of subjects, 0,001 events/patient year) were observed with semaglutide in combination with oral antidiabetics other than sulfonylurea.

Gastrointestinal adverse reactions

Nausea occurred in 15 %, diarrhoea in 10 %, and vomiting in 7 % of patients when treated with semaglutide. Most events were mild to moderate in severity and of short duration. The events led to treatment discontinuation in 4 % of subjects. The events were most frequently reported during the first months on treatment.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Acute pancreatitis confirmed by adjudication has been reported in phase 3a trials, semaglutide (< 0,1 %) and comparator (0,2 %). In the cardiovascular outcomes trial the frequency of acute pancreatitis confirmed by adjudication was 0,1 % for semaglutide and 0,2 % for placebo (see section 4.4)

Diabetic retinopathy complications

A 2-year clinical trial with s.c. semaglutide investigated 3 297 patients with type 2 diabetes, with high cardiovascular risk, long duration of diabetes and poorly controlled blood glucose. In this trial, adjudicated events of diabetic retinopathy complications occurred in more patients treated with s.c. semaglutide (3,0 %) compared to placebo (1,8 %). This was observed in insulin-treated patients with known diabetic retinopathy. The treatment difference appeared early and persisted throughout the trial. Systematic evaluation of diabetic retinopathy complication was only performed in the cardiovascular outcomes trial with s.c. semaglutide.

In clinical trials with Rybelsus of up to 18 months duration involving 6 352 patients with type 2 diabetes, adverse events related to diabetic retinopathy were reported in similar proportions in subjects treated with semaglutide (4,2 %) and comparators (3,8 %).

Immunogenicity

Consistent with the potential immunogenic properties of medicinal products containing proteins or peptides, patients may develop antibodies following treatment with semaglutide. The proportion of subjects tested positive for anti-semaglutide antibodies at any time point after baseline was low (0,5 %) and no subjects had neutralising anti-semaglutide antibodies or anti-semaglutide antibodies with neutralising effect on endogenous GLP-1 at end-of-trial.

Heart rate increase

Increased heart rate has been observed with GLP-1 receptor agonists. In the phase 3a trials, mean changes of 0 to 4 beats per minute (bpm) from a baseline of 69 to 76 were observed in patients treated with Rybelsus.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Reporting of suspected adverse reactions:

Reporting suspected adverse reactions after authorisation of Rybelsus is important. It allows continued monitoring of the benefit/risk balance of Rybelsus. Healthcare providers are asked to report any suspected adverse reactions to SAHPRA via the “**6.04 Adverse Drug Reactions Reporting Form**”, found online under SAHPRA’s publications:

<https://www.sahpra.org.za/Publications/Index/8>

Discontinuation due to an adverse event

Discontinuation of treatment due to adverse events was 9% for patients treated with Rybelsus. The most frequent adverse events leading to discontinuation were gastrointestinal.

4.9 Overdose

Effects of overdose with semaglutide in clinical studies may be associated with gastrointestinal disorders. In the event of overdose, appropriate supportive treatment should be initiated according to the patient’s clinical signs and symptoms. A prolonged period of observation and treatment of the symptoms may be necessary, taking into account the long half-life of semaglutide of approximately 1 week (see section 5.2). There is no specific antidote for overdose with semaglutide.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Drugs used in diabetes, Glucagon-like peptide-1 (GLP 1) analogues, ATC code: A10BJ06

Mechanism of action

Semaglutide is a GLP 1 analogue with 94 % sequence homology to human GLP 1.

Semaglutide acts as a GLP 1 receptor agonist that selectively binds to and activates the GLP 1 receptor, the target for native GLP 1.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

GLP 1 is a physiological hormone that has multiple actions in glucose and appetite regulation, and in the cardiovascular system. The glucose and appetite effects are specifically mediated via GLP 1 receptors in the pancreas and the brain.

Semaglutide reduces blood glucose in a glucose-dependent manner by stimulating insulin secretion and lowering glucagon secretion when blood glucose is high. The mechanism of blood glucose lowering also involves a minor delay in gastric emptying in the early postprandial phase. During hypoglycaemia, semaglutide diminishes insulin secretion and does not impair glucagon secretion. The mechanism of semaglutide is independent of the route of administration.

Semaglutide reduces body weight and body fat mass through lowered energy intake, involving an overall reduced appetite. In addition, semaglutide reduces the preference for high fat foods.

GLP 1 receptors are expressed in the heart, vasculature, immune system and kidneys. Semaglutide has a beneficial effect on plasma lipids, lowers systolic blood pressure and reduces inflammation in clinical studies. In animal studies, semaglutide attenuates the development of atherosclerosis by preventing aortic plaque progression and reducing inflammation in the plaque.

Pharmacodynamic effects

Rybelsus lowers fasting glucose and self-measured plasma glucose. The onset happens early with a lowering of fasting glucose within the first week of treatment.

The pharmacodynamic evaluations described below were performed with orally administered semaglutide after 12 weeks of treatment.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Fasting and postprandial glucose

Semaglutide reduces fasting and postprandial glucose concentrations. In patients with type 2 diabetes, treatment with semaglutide resulted in a relative reduction compared to placebo of 22 % [13; 30] for fasting glucose and 29 % [19; 37] for postprandial glucose.

Beta-cell function and insulin secretion

Semaglutide improves beta-cell function. Compared to placebo, semaglutide improved both first- and second-phase insulin response, with a 3- and 2-fold increase, respectively, and increased maximal beta-cell secretory capacity after an arginine stimulation test in patients with type 2 diabetes. In addition, semaglutide treatment increased fasting insulin concentrations compared to placebo.

Glucagon secretion

Semaglutide lowers the postprandial glucagon concentrations. In patients with type 2 diabetes, semaglutide resulted in the following relative reductions in glucagon compared to placebo: postprandial glucagon response of 29 % [15; 41].

Glucose-dependent insulin and glucagon secretion

Semaglutide reduces blood glucose in a glucose-dependent manner by stimulating insulin secretion and lowering glucagon secretion when blood glucose is high. The mechanism of blood glucose lowering also involves a minor delay in gastric emptying in the early postprandial phase. During hypoglycaemia, semaglutide diminishes insulin secretion and does not impair glucagon secretion. The mechanism of semaglutide is independent of the route of administration.

Gastric emptying

Semaglutide causes a minor delay in early postprandial gastric emptying, with paracetamol exposure (AUC 0 – 1 h) 31 % [13; 46] lower in the first hour after the meal, thereby reducing the rate at which glucose appears in the circulation postprandially.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Body weight and body composition

A greater reduction in body weight was observed with Rybelsus compared to studied comparators (placebo, sitagliptin, empagliflozin and liraglutide). The body weight loss with semaglutide was predominantly from fat tissue with loss of fat mass being 3-fold larger than loss of lean mass.

Appetite, energy intake and food choice

Semaglutide compared to placebo lowered the energy intake of 3 consecutive ad libitum meals by 18 – 35 %. This was supported by a semaglutide-induced suppression of appetite in the fasting state as well as postprandially, improved control of eating, less food cravings and a relative lower preference for high fat food.

Fasting and postprandial lipids

Semaglutide compared to placebo lowered fasting triglyceride and very low density lipoproteins (VLDL) cholesterol concentrations by 19 % [8; 28] and 20 % [5; 33], respectively. The postprandial triglyceride and VLDL cholesterol response to a high fat meal was reduced by 24 % [9; 36] and 21 % [7; 32], respectively. ApoB48 was reduced both in fasting and postprandial state by 25 % [2; 42] and 30 % [15; 43], respectively.

Clinical efficacy and safety

The efficacy and safety of Rybelsus have been evaluated in eight global randomised controlled phase 3a trials. In seven trials, the primary objective was the assessment of the glycaemic efficacy; in one trial, the primary objective was the assessment of cardiovascular outcomes.

The trials included 8 842 randomised patients with type 2 diabetes (5 169 treated with semaglutide), including 1 165 patients with moderate renal impairment. Patients had an average age of 61 years (range 18 to 92 years), with 40 % of patients ≥ 65 years of age and

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

8 % ≥ 75 years of age. The efficacy of semaglutide was compared with placebo or active controls (sitagliptin, empagliflozin and liraglutide).

The efficacy of semaglutide was not impacted by baseline age, gender, race, ethnicity, body weight, BMI, diabetes duration, upper gastrointestinal disease and level of renal function.

PIONEER 1 – Monotherapy

In a 26-week double-blind trial, 703 patients with type 2 diabetes inadequately controlled with diet and exercise were randomized to Rybelsus 3 mg, Rybelsus 7 mg, Rybelsus 14 mg or placebo once daily.

The mean age of the study population was 55 years, and the mean duration of type 2 diabetes was 3,5 years. Overall, 75 % were White, 5 % were Black or African American and 17 % were Asian. Hispanic or Latino patients comprised 26 % (n = 180) of the population.

The mean body weight at baseline was 88 kg.

Monotherapy with Rybelsus 7 mg and 14 mg once daily was superior at week 26 in reducing HbA_{1c} compared with placebo. Rybelsus 14 mg was superior in reducing body weight compared with placebo ([Table 2](#)).

Table 2 Results of a 26-week monotherapy trial comparing semaglutide with placebo (PIONEER 1)

	Semaglutide 7 mg	Semaglutide 14 mg	Placebo
Full analysis set (N)	175	175	178
HbA_{1c} (%)			
Baseline	8,0	8,0	7,9
Change from baseline ¹	-1,2	-1,4	-0,3
Difference from placebo ¹ [95% CI]	-0,9 [-1,1; -0,6]*	-1,1 [-1,3; -0,9]*	-
Patients (%) achieving HbA_{1c} <7.0%	69 ^s	77 ^s	31
FPG (mmol/L)			

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Baseline	9,0	8,8	8,9
Change from baseline ¹	-1,5	-1,8	-0,2
Difference from placebo ¹ [95% CI]	-1,4 [-1,9; -0,8] [§]	-1,6 [-2,1; -1,2] [§]	-
Body weight (kg)			
Baseline	89,0	88,1	88,6
Change from baseline ¹	-2,3	-3,7	-1,4
Difference from placebo ¹ [95% CI]	-0,9 [-1,9; 0,1]	-2,3 [-3,1; -1,5]*	-

¹ Irrespective of treatment discontinuation or initiation of rescue medication (pattern

mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity. § p < 0,05, not controlled for multiplicity; for 'Patients achieving HbA_{1c} < 7,0 %', the p-value is for the odds ratio.

PIONEER 2 – Semaglutide vs. empagliflozin, both in combination with metformin

In a 52-week open-label trial (26-week primary endpoint), 822 patients with type 2 diabetes were randomized to semaglutide 14 mg or empagliflozin 25 mg once daily, both in combination with metformin.

The mean age of the trial population was 58 years, and the mean duration of type 2 diabetes was 7,4 years. Overall, 86 % were White, 7 % were Black or African American and 6 % were Asian. Hispanic or Latino patients comprised 24 % (n = 199) of the population. The mean body weight at baseline was 92 kg.

Treatment with semaglutide 14 mg once daily was superior at week 26 in reducing HbA_{1c} compared to empagliflozin 25 mg once daily ([Table 3](#)).

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Table 3 Results of a 52-week trial comparing semaglutide with empagliflozin

(PIONEER 2)

	Semaglutide 14 mg	Empagliflozin 25 mg
Full analysis set (N)	411	410
Week 26		
HbA_{1c} (%)		
Baseline	8,1	8,1
Change from baseline ¹	-1,3	-0,9
Difference from empagliflozin ¹ [95 % CI]	-0,4 [-0,6; -0,3]*	-
Patients (%) achieving HbA_{1c} < 7,0 %	67 [§]	40
FPG (mmol/L)		
Baseline	9,5	9,7
Change from baseline ¹	-2,0	-2,0
Difference from empagliflozin ¹ [95 % CI]	0,0 [-0,2; 0,3]	-
Body weight (kg)		
Baseline	91,9	91,3
Change from baseline ¹	-3,8	-3,7
Difference from empagliflozin ¹ [95 % CI]	-0,1 [-0,7; 0,5]	-
Week 52		
HbA_{1c} (%)		
Change from baseline ¹	-1,3	-0,9
Difference from empagliflozin ¹ [95 % CI]	-0,4 [-0,5; -0,3] [§]	-
Patients (%) achieving HbA_{1c} < 7,0 %	66 [§]	43
Body weight (kg)		
Change from baseline ¹	-3,8	-3,6
Difference from empagliflozin ¹ [95% CI]	-0,2 [-0,9; 0,5]	-

¹ Irrespective of treatment discontinuation or initiation of rescue medication (pattern mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity. § p < 0,05, not controlled for multiplicity; for 'Patients achieving HbA_{1c} < 7,0%', the p-value is for the odds ratio.

PIONEER 3 – Semaglutide vs. sitagliptin, both in combination with metformin or metformin with sulfonylurea

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

In a 78-week, double-blind, double-dummy trial (26-week primary endpoint), 1 864 patients with type 2 diabetes were randomized to semaglutide 3 mg, semaglutide 7 mg, semaglutide 14 mg or sitagliptin 100 mg once daily, all in combination with metformin alone or metformin and sulfonylurea.

The mean age of the trial population was 58 years, and the mean duration of type 2 diabetes was 8,6 years. Overall, 71 % were White, 9 % were Black or African American and 13 % were Asian. Hispanic or Latino patients comprised 17 % (n = 321) of the population. The mean body weight at baseline was 91 kg.

Treatment with Rybelsus 7 mg and 14 mg once daily was superior at week 26 in reducing HbA_{1c} and body weight compared to sitagliptin 100 mg once daily ([Table 4](#)).

Table 4 Results of a 78-week trial comparing semaglutide with sitagliptin (PIONEER 3)

	Semaglutide 7 mg	Semaglutide 14 mg	Sitagliptin 100 mg
Full analysis set (N)	465	465	467
Week 26			
HbA_{1c} (%)			
Baseline	8,4	8,3	8,3
Change from baseline ¹	-1,0	-1,3	-0,8
Difference from sitagliptin ¹ [95% CI]	-0,3 [-0,4; -0,1] [*]	-0,5 [-0,6; -0,4] [*]	-
Patients (%) achieving HbA_{1c} < 7,0 %	44 [§]	56 [§]	32
FPG (mmol/L)			
Baseline	9,4	9,3	9,5
Change from baseline ¹	-1,2	-1,7	-0,9
Difference from sitagliptin ¹ [95% CI]	-0,3 [-0,6; 0,0] [§]	-0,8 [-1,1; -0,5] [§]	-
Body weight (kg)			
Baseline	91,3	91,2	90,9
Change from baseline ¹	-2,2	-3,1	-0,6
Difference from sitagliptin ¹ [95 % CI]	-1,6 [-2,0; -1,1] [*]	-2,5 [-3,0; -2,0] [*]	-
Week 78			

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

HbA_{1c} (%)			
Change from baseline ¹	-0,8	-1,1	-0,7
Difference from sitagliptin ¹ [95 % CI]	-0,1 [-0,3; 0,0]	-0,4 [-0,6; -0,3] [§]	-
Patients (%) achieving HbA_{1c} < 7,0 %	39 [§]	45 [§]	29
Body weight (kg)			
Change from baseline ¹	-2,7	-3,2	-1.0
Difference from sitagliptin ¹ [95 % CI]	-1,7 [-2,3; -1,0] [§]	-2,1 [-2,8; -1,5] [§]	-

¹ Irrespective of treatment discontinuation or initiation of rescue medication (pattern mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity. § p<0.05, not controlled for multiplicity; for 'Patients achieving HbA_{1c} < 7,0 %', the p-value is for the odds ratio.

PIONEER 4 – Semaglutide vs. liraglutide and placebo, all in combination with metformin or metformin with an SGLT2 inhibitor

In a 52-week double-blind, double-dummy trial (26-week primary endpoint), 711 patients with type 2 diabetes were randomized to semaglutide 14 mg, liraglutide 1,8 mg s.c. injection or placebo once daily, all in combination with metformin or metformin and an SGLT2 inhibitor.

The mean age of the trial population was 56 years, and the mean duration of type 2 diabetes was 7,6 years. Overall, 73 % were White, 4 % were Black or African American and 13 % were Asian. Hispanic or Latino patients comprised 6 % (n = 40) of the population. The mean body weight at baseline was 94 kg.

Treatment with semaglutide 14 mg once daily was superior at week 26 in reducing HbA_{1c} and body weight compared with placebo. Treatment with semaglutide 14 mg was non-inferior in reducing HbA_{1c} and superior in reducing body weight at week 26 compared with liraglutide 1,8 mg ([Table 5](#)).

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Table 5 Results of a 52-week trial comparing semaglutide with liraglutide and placebo

(PIONEER 4)

	Semaglutide 14 mg	Liraglutide 1.8 mg	Placebo
Full analysis set (N)	285	284	142
Week 26			
HbA_{1c} (%)			
Baseline	8.0	8.0	7.9
Change from baseline ¹	-1.2	-1.1	-0.2
Difference from liraglutide ¹ [95 % CI]	-0.1 [-0.3; 0.0]	-	-
Difference from placebo ¹ [95 % CI]	-1.1 [-1.2; -0.9]*	-	-
Patients (%) achieving HbA_{1c} < 7.0 %	68 ^{§,a}	62	14
FPG (mmol/L)			
Baseline	9.3	9.3	9.2
Change from baseline ¹	-2.0	-1.9	-0.4
Difference from liraglutide ¹ [95 % CI]	-0.1 [-0.4; 0.1]	-	-
Difference from placebo ¹ [95 % CI]	-1.6 [-2.0; -1.3] [§]	-	-
Body weight (kg)			
Baseline	92.9	95.5	93.2
Change from baseline ¹	-4.4	-3.1	-0.5
Difference from liraglutide ¹ [95 % CI]	-1.2 [-1.9; -0.6]*	-	-
Difference from placebo ¹ [95 % CI]	-3.8 [-4.7; -3.0]*	-	-
Week 52			
HbA_{1c} (%)			
Change from baseline ¹	-1.2	-0.9	-0.2
Difference from liraglutide ¹ [95 % CI]	-0.3 [-0.5; -0.1] [§]	-	-

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Difference from placebo ¹ [95 % CI]	-1,0 [-1,2; -0,8] [§]	-	-
Patients (%) achieving HbA_{1c} < 7,0 %	61 ^{§,a}	55	15
Body weight (kg)			
Change from baseline ¹	-4,3	-3,0	-1,0
Difference from liraglutide ¹ [95 % CI]	-1,3 [-2,1; -0,5] [§]	-	-
Difference from placebo ¹ [95 % CI]	-3,3 [-4,3; -2,4] [§]	-	-

¹ Irrespective of treatment discontinuation or initiation of rescue medication (pattern mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity. § p < 0.05, not controlled for multiplicity; for 'Patients achieving HbA_{1c} < 7,0 %', the p-value is for the odds ratio. a vs placebo.

PIONEER 5 – Semaglutide vs. placebo, both in combination with basal insulin alone, metformin and basal insulin or metformin and/or sulfonylurea, in patients with moderate renal impairment

In a 26-week double-blind trial, 324 patients with type 2 diabetes and moderate renal impairment (eGFR 30 - 59 mL/min/1,73 m²) were randomized to semaglutide 14 mg or placebo once daily. Trial product was added to the patient's stable pre-trial antidiabetic regimen.

The mean age of the trial population was 70 years, and the mean duration of type 2 diabetes was 14,0 years. Overall, 96 % were White, 4 % were Black or African American and less than 1 % were Asian. Hispanic or Latino patients comprised 6 % (n = 21) of the population.

The mean body weight at baseline was 91 kg.

Treatment with semaglutide 14 mg once daily was superior at week 26 in reducing HbA_{1c} and body weight compared with placebo.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Table 6 Results of a 26-week trial comparing semaglutide with placebo in patients with type 2 diabetes and moderate renal impairment (PIONEER 5)

	Semaglutide 14 mg	Placebo
Full analysis set (N)	163	161
HbA_{1c} (%)		
Baseline	8,0	7,9
Change from baseline ¹	-1,0	-0,2
Difference from placebo ¹ [95 % CI]	-0,8 [-1,0; -0,6]*	-
Patients (%) achieving HbA_{1c} < 7,0 %	58 [§]	23
FPG (mmol/L)		
Baseline	9,1	9,1
Change from baseline ¹	-1,5	-0,4
Difference from placebo ¹ [95 % CI]	-1,2 [-1,7; -0,6] [§]	-
Body weight (kg)		
Baseline	91,3	90,4
Change from baseline ¹	-3,4	-0,9
Difference from placebo ¹ [95 % CI]	-2,5 [-3,2; -1,8]*	-

¹ Irrespective of treatment discontinuation or initiation of rescue medication (pattern mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity. § p < 0,05, not controlled for multiplicity; for 'Patients achieving HbA_{1c} <7,0 %', the p-value is for the odds ratio.

PIONEER 7 – Semaglutide vs. sitagliptin, both in combination with metformin, SGLT2 inhibitors, sulfonylurea or thiazolidinediones. Flexible-dose-adjustment trial

In a 52-week open-label trial, 504 patients with type 2 diabetes were randomized to semaglutide (flexible dose adjustment of 3 mg, 7 mg, and 14 mg once daily) or sitagliptin 100 mg once daily, all in combination with 1 - 2 oral glucose-lowering medications (metformin, SGLT2 inhibitors, sulfonylurea or thiazolidinediones). The dose of semaglutide was adjusted every 8 weeks based on patient's glycaemic response and tolerability. The sitagliptin 100 mg dose was fixed. The efficacy and safety of semaglutide were evaluated at

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

week 52. At the end of 52 weeks, the percentage of patients on-treatment with semaglutide 7 mg were 30,2 % and semaglutide 14 mg were 59,4 %.

The mean age of the trial population was 57 years, and the mean duration of type 2 diabetes was 8,8 years. Overall, 76 % were White, 9 % were Black or African American and 14 % were Asian. Hispanic or Latino patients comprised 21 % (n = 105) of the population. The mean body weight at baseline was 89 kg.

After 52 weeks of treatment, 58,3% of the patients achieved the target of $HbA_{1c} < 7,0$ % with adjustable dosing of semaglutide compared to 25,2 % of patients treated with sitagliptin 100 mg. Rybelsus® was superior to sitagliptin at week 52 in enabling patients to achieve $HbA_{1c} < 7,0$ % and in reducing body weight (Table 7).

Table 7 Results of a 52-week flexible-dose-adjustment trial comparing semaglutide with sitagliptin (PIONEER 7)

	Semaglutide Flexible dose	Sitagliptin 100 mg
Full analysis set (N)	253	251
HbA_{1c} (%)		
Baseline	8,3	8,3
Patients (%) achieving $HbA_{1c} < 7,0$ % ¹	58*	25
Body weight (kg)		
Baseline	88,9	88,4
Change from baseline ¹	-2,6	-0,7
Difference from sitagliptin ¹ [95 % CI]	-1,9 [-2,6; -1,2]*	-

¹ Irrespective of treatment discontinuation (16,6 % of the patients with semaglutide flexible dose and 9,2 % with sitagliptin, where 8,7 % and 4,0 %, respectively, were due to AEs) or initiation of rescue medication (pattern mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity (for 'Patients achieving $HbA_{1c} < 7,0$ %', the p-value is for the odds ratio).

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

PIONEER 8 – Semaglutide vs. placebo, both in combination with insulin with or without metformin

In a 52-week double-blind trial, 731 patients with type 2 diabetes inadequately controlled on insulin (basal, basal/bolus or premixed) with or without metformin were randomized to semaglutide 3 mg, semaglutide 7 mg, semaglutide 14 mg or placebo once daily. All patients reduced their insulin dose by 20 % at randomization to reduce the risk of hypoglycemia. For the first 26 weeks, patients were allowed to increase the insulin dose only up to the starting insulin dose prior to randomization. After the 26 weeks, patients were allowed to adjust the insulin dose as needed. At randomization, the total daily insulin dose were 55 U, 63 U, and 53 U for placebo, semaglutide 7 mg and semaglutide 14 mg, respectively.

The mean age of the trial population was 61 years, and the mean duration of type 2 diabetes was 15,0 years. Overall, 51 % were White, 7 % were Black or African American and 36 % were Asian. Hispanic or Latino patients comprised 13 % (n = 97) of the population. The mean body weight at baseline was 86 kg.

Treatment with semaglutide 7 mg and 14 mg once daily was superior in reducing HbA_{1c} and body weight compared with placebo ([Table 8](#)).

Table 8 Results of a 52-week trial comparing semaglutide with placebo in combination with insulin (PIONEER 8)

	Semaglutide 7 mg	Semaglutide 14 mg	Placebo
Full analysis set (N)	182	181	184
Week 26 (insulin dose capped to baseline level)			
HbA_{1c} (%)			
Baseline	8,2	8,2	8,2
Change from baseline ¹	-0,9	-1,3	-0,1
Difference from placebo ¹ [95 % CI]	-0,9 [-1,1; -0,7]*	-1,2 [-1,4; -1,0]*	-

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Patients (%) achieving HbA_{1c} < 7,0 %	43 [§]	58 [§]	7
FPG (mmol/L)			
Baseline	8,5	8,3	8,3
Change from baseline ¹	-1,1	-1,3	0,3
Difference from placebo ¹ [95 % CI]	-1,4 [-1,9; -0,8] [§]	-1,6 [-2,2; -1,1] [§]	-
Body weight (kg)			
Baseline	87,1	84,6	86,0
Change from baseline ¹	-2,4	-3,7	-0,4
Difference from placebo ¹ [95 % CI]	-2,0 [-3,0; -1,0]*	-3,3 [-4,2; -2,3]*	-
Week 52 (uncapped insulin dose)⁺			
HbA_{1c} (%)			
Change from baseline ¹	-0,8	-1,2	-0,2
Difference from placebo ¹ [95 % CI]	-0,6 [-0,8; -0,4] [§]	-0,9 [-1,1; -0,7] [§]	-
Patients (%) achieving HbA_{1c} < 7,0 %	40 [§]	54 [§]	9
Body weight (kg)			
Change from baseline ¹	-2,0	-3,7	0,5
Difference from placebo ¹ [95% CI]	-2,5 [-3,6; -1,4] [§]	-4,3 [-5,3; -3,2] [§]	-

¹ Irrespective of treatment discontinuation or initiation of rescue medication (pattern mixture model using multiple imputation). * p < 0,001 (unadjusted 2-sided) for superiority, controlled for multiplicity. § p<0.05, not controlled for multiplicity; for 'Patients achieving HbA_{1c} < 7,0 %', the p-value is for the odds ratio. + The total daily insulin dose was statistically significantly lower with semaglutide than with placebo at week 52.

The mean changes from baseline in total daily insulin dose at week 26 were -1 U, -8 U and -9 U for placebo, Rybelsus® 7 mg and Rybelsus® 14 mg, respectively. The difference from placebo for Rybelsus® 7 mg and Rybelsus® 14 mg was -8 [-12; -3]_{95 % CI} and -8 [-13; -3]_{95 % CI}, respectively. The mean changes from baseline in daily insulin dose at week

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

52 were 10 U, -6 U and -7 U for placebo, Rybelsus® 7 mg and Rybelsus® 14 mg, respectively. The difference from placebo for Rybelsus® 7 mg and Rybelsus® 14 mg was -16 [-25; -8]_{95 % CI} and -17 [-25; -9]_{95 % CI}, respectively.

Cardiovascular evaluation


In a double-blind trial (PIONEER 6), 3 183 patients with type 2 diabetes at high cardiovascular risk were randomised to Rybelsus 14 mg once daily or placebo in addition to standard-of-care. The median observation period was 16 months.

The primary endpoint was time from randomisation to first occurrence of a major adverse cardiovascular event (MACE): cardiovascular death, non-fatal myocardial infarction or non-fatal stroke.

Patients eligible to enter the trial were: 50 years of age or older and with established cardiovascular disease and/or chronic kidney disease, or 60 years of age or older and with cardiovascular risk factors only. In total, 1 797 patients (56,5 %) had established cardiovascular disease without chronic kidney disease, 354 (11,1 %) had chronic kidney disease only and 544 (17,1 %) had both cardiovascular disease and kidney disease. 488 patients (15,3 %) had cardiovascular risk factors only. The mean age at baseline was 66 years, and 68 % of the patients were men. The mean duration of diabetes was 14,9 years and the mean BMI was 32,3 kg/m². Medical history included stroke (11,7 %) and myocardial infarction (36,1 %).

The total number of first MACE was 137: 61 (3,8 %) with semaglutide and 76 (4,8 %) with placebo. The analysis of time to first MACE resulted in a HR of 0.79 [0.57; 1.11]_{95% CI}.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)



Cumulative incidence plot of primary outcome (a composite of cardiovascular death, nonfatal myocardial infarction, or nonfatal stroke) with non-cardiovascular death as competing risk.

Abbreviations: CI: Confidence interval, HR: Hazard ratio

Figure 1 Cumulative incidence of first occurrence of MACE in PIONEER 6

The treatment effect for the primary composite endpoint and its components in the PIONEER 6 trial is shown in Figure 2.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)




Figure 2 Treatment effect for the primary composite endpoint, its components and all cause death (PIONEER 6)

Body weight

By end-of-treatment, 27 - 65,7 % of the patients had achieved a weight loss of ≥ 5 % and 6 - 34,7 % had achieved a weight loss of ≥ 10 % with semaglutide, compared with 12 – 39 % and 2 – 8 %, respectively, with the active comparators.

Blood pressure

Treatment with semaglutide had reduced systolic blood pressure by 2 - 7 mmHg.

Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Rybelsus in one or more subsets of the paediatric population in type 2 diabetes (see section 4.2 for information on paediatric use).

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

5.2 Pharmacokinetic properties

Absorption

Orally administered semaglutide has a low absolute bioavailability and a variable absorption.

Daily administration according to the recommended posology in combination with a long half-life reduces day-to-day fluctuation of the exposure.

The pharmacokinetics of semaglutide have been extensively characterised in healthy subjects and patients with type 2 diabetes. Following oral administration, maximum plasma concentration of semaglutide occurred 1 hour post dose. Steady-state exposure was reached after 4 – 5 weeks of once-daily administration. In patients with type 2 diabetes, the average steady-state concentrations were approximately 6,7 nmol/L and 14,6 nmol/L with semaglutide 7 mg and 14 mg, respectively; with 90% of subjects treated with semaglutide 7 mg having an average concentration between 1,7 and 22,7 nmol/L and 90 % of subjects treated with semaglutide 14 mg having an average concentration between 3,7 and 41,3 nmol/L. Systemic exposure of semaglutide increased in a dose-proportional manner.

Based on in vitro data, salcaprozate sodium facilitates absorption of semaglutide. The absorption of semaglutide predominantly occurs in the stomach.

The estimated bioavailability of semaglutide is approximately 1 % following oral administration. The between-subject variability in absorption was high (coefficient of variation was approximately 100 %). The estimation of the within-subject variability in bioavailability was not reliable.

Absorption of semaglutide is decreased if taken with food or large volumes of water. A longer post-dose fasting period results in higher absorption.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Distribution

The estimated absolute volume of distribution is approximately 8 L in subjects with type 2 diabetes. Semaglutide is extensively bound to plasma proteins (> 99 %).

Biotransformation

Semaglutide is metabolised through proteolytic cleavage of the peptide backbone and sequential beta-oxidation of the fatty acid sidechain. The enzyme neutral endopeptidase (NEP) is expected to be involved in the metabolism of semaglutide.

Elimination

The primary excretion routes of semaglutide-related material are via the urine and faeces. Approximately 3 % of the absorbed dose is excreted as intact semaglutide via the urine.

With an elimination half-life of approximately 1 week, semaglutide will be present in the circulation for about 5 weeks after the last dose. The clearance of semaglutide in patients with type 2 diabetes is approximately 0,04 L/h.

Switching between oral and subcutaneous (s.c.) administration

The effect of switching between oral and s.c. semaglutide cannot easily be predicted because of the high pharmacokinetic variability of oral semaglutide. Exposure after oral semaglutide 14 mg once daily is comparable to s.c. semaglutide 0,5 mg once weekly. An oral dose equivalent to 1,0 mg of s.c. semaglutide has not been established.

Special populations

Elderly

Age had no effect on the pharmacokinetics of semaglutide based on data from clinical trials, which studied patients up to 92 years of age.

Gender

Gender had no clinically meaningful effects on the pharmacokinetics of semaglutide.

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

Race and ethnicity

Race (White, Black or African-American, Asian) and ethnicity (Hispanic or Latino, not Hispanic or Latino) had no effect on the pharmacokinetics of semaglutide.

Body weight

Body weight had an effect on the exposure of semaglutide. Higher body weight was associated with lower exposure. Semaglutide provided adequate systemic exposure over the body weight range of 40 - 212 kg evaluated in the clinical trials.

Renal impairment

Renal impairment did not impact the pharmacokinetics of semaglutide in a clinically relevant manner. The pharmacokinetics of semaglutide were evaluated in patients with mild, moderate or severe renal impairment and patients with end-stage renal disease on dialysis compared with subjects with normal renal function in a study with 10 consecutive days of once-daily doses of semaglutide. This was also shown for subjects with type 2 diabetes and renal impairment based on data from phase 3a studies.

Hepatic impairment

Hepatic impairment did not impact the pharmacokinetics of semaglutide in a clinically relevant manner. The pharmacokinetics of semaglutide were evaluated in patients with mild, moderate or severe hepatic impairment compared with subjects with normal hepatic function in a study with 10 consecutive days of once-daily doses of semaglutide.

Upper GI tract disease

Upper GI tract disease (chronic gastritis and/or gastroesophageal reflux disease) did not impact the pharmacokinetics of semaglutide in a clinically relevant manner. The pharmacokinetics were evaluated in patients with type 2 diabetes with or without upper GI tract disease dosed for 10 consecutive days with once-daily doses of semaglutide. This was

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

also shown for subjects with type 2 diabetes and upper GI tract disease based on data from phase 3a studies.

Paediatric population

Semaglutide has not been studied in paediatric patients.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity or genotoxicity.

Non-lethal thyroid C-cell tumours observed in rodents are a class effect for GLP 1 receptor agonists. In 2-year carcinogenicity studies in rats and mice, semaglutide caused thyroid C-cell tumours at clinically relevant exposures. No other treatment-related tumours were observed. The rodent C-cell tumours are caused by a non-genotoxic, specific GLP 1 receptor mediated mechanism to which rodents are particularly sensitive. The relevance for humans is considered to be low, but cannot be completely excluded.

In fertility studies in rats, semaglutide did not affect mating performance or male fertility. In female rats, an increase in oestrous cycle length and a small reduction in corpora lutea (ovulations) were observed at doses associated with maternal body weight loss.

In embryo-foetal development studies in rats, semaglutide caused embryotoxicity below clinically relevant exposures. Semaglutide caused marked reductions in maternal body weight and reductions in embryonic survival and growth. In foetuses, major skeletal and visceral malformations were observed, including effects on long bones, ribs, vertebrae, tail, blood vessels and brain ventricles. Mechanistic evaluations indicated that the embryotoxicity involved a GLP 1 receptor mediated impairment of the nutrient supply to the embryo across the rat yolk sac. Due to species differences in yolk sac anatomy and function, and due to the lack of GLP 1 receptor expression in the yolk sac of non-human primates, this mechanism is

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

considered unlikely to be of relevance to humans. However, a direct effect of semaglutide on the foetus cannot be excluded.

In developmental toxicity studies in rabbits and cynomolgus monkeys, increased pregnancy loss and slightly increased incidence of foetal abnormalities were observed at clinically relevant exposures. The findings coincided with marked maternal body weight loss of up to 16 %. Whether these effects are related to the decreased maternal food consumption as a direct GLP 1 effect is unknown.

Postnatal growth and development were evaluated in cynomolgus monkeys. Infants were slightly smaller at delivery but recovered during the lactation period.

In juvenile rats, semaglutide caused delayed sexual maturation in both males and females. These delays had no impact upon fertility and reproductive capacity of either sex, or on the ability of the females to maintain pregnancy.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Salcaprozate sodium

Povidone K90

Cellulose, microcrystalline

Magnesium stearate

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 mg: 24 months

7 mg: 30 months

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

14 mg: 30 months.

Store at or below 30 °C.

6.4 Special precautions for storage

Store in the original blister package in order to protect from light and moisture.

6.5 Nature and contents of container

Alu/Alu blisters.

Pack sizes of 3 mg tablets: 10, 30, 60 and 90 tablets. The 3 mg blister foil is bright green in colour.

Pack sizes of 7 mg tablets: 30, 60 and 90 tablets. The 7 mg blister foil is bright red in colour.

Pack sizes of 14 mg tablets: 30, 60 and 90 tablets. The 14 mg blister foil is bright blue in colour.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. HOLDER OF CERTIFICATE OF REGISTRATION

Novo Nordisk (Pty) Ltd

150 Rivonia Road

10 Marion Street Office Park

Building C1

Sandton

Johannesburg 2196

Applicant	Novo Nordisk (Pty) Ltd	Dosage form and strength	Tablet; 3 mg Semaglutide Tablet; 7 mg Semaglutide Tablet ; 14 mg Semaglutide
Product name	Rybelsus 3 mg Rybelsus 7 mg Rybelsus 14 mg	Application number	560337 560338 560339
Date of approval	03 September 2024	Module 1	1.3.1.1.1 PI (approved)

8. REGISTRATION NUMBERS

Rybelsus 3 mg: 56/21.13/0337

Rybelsus 7 mg: 56/21.13/0338

Rybelsus 14 mg: 56/21.13/0339

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

3 September 2024

10. DATE OF REVISION OF THE TEXT

Not applicable